


# Introduction to Machine Learning

## Maximum Likelihood Estimation vs. Empirical Risk Minimization



### Learning goals

- Learn the correspondence of between a Laplacian error distributions and the L1 loss
- Learn that there is no error distribution for the Huber loss
- Learn the correspondence between Bernoulli-distributed targets and the Bernoulli loss

# LAPLACE ERRORS - L1-LOSS

Let us assume that errors are Laplacian, i.e.  $\epsilon$  follows a Laplace distribution which has the density

$$\frac{1}{2\sigma} \exp\left(-\frac{|x|}{\sigma}\right), \sigma > 0.$$

Then

$$y = f_{\text{true}}(\mathbf{x}) + \epsilon$$

follows a Laplace distribution with mean  $f(\mathbf{x}^{(i)} | \theta)$  and scale parameter  $\sigma$ .

# LAPLACE ERRORS - L1-LOSS

The likelihood is then

$$\begin{aligned}\mathcal{L}(\boldsymbol{\theta}) &= \prod_{i=1}^n p\left(y^{(i)} \mid f\left(\mathbf{x}^{(i)} \mid \boldsymbol{\theta}\right), \sigma\right) \\ &\propto \exp\left(-\frac{1}{\sigma} \sum_{i=1}^n \left|y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \boldsymbol{\theta}\right)\right|\right).\end{aligned}$$



The negative log-likelihood is

$$-\ell(\boldsymbol{\theta}) \propto -\sum_{i=1}^n \left|y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \boldsymbol{\theta}\right)\right|.$$

Minimizing the negative log-likelihood for Laplacian error terms corresponds to empirical risk minimization with L1-loss.



# LAPLACE ERRORS - L1-LOSS

- Distribution of empirical residuals and their comparison to the theoretical quantiles of a Laplace-distribution.



# OTHER ERROR DISTRIBUTIONS

- There are losses that do not correspond to “real” error densities, like the Huber loss. (In the QQ-plot below we show residuals against quantiles of a normal. )



# OTHER ERROR DISTRIBUTIONS

However, intuitively, we see that a certain type of loss function corresponds to a certain error distribution.

Loss function	Error Distribution
$L_2$ -Loss	Gaussian Errors
$L_1$ -Loss	Laplace Errors
Huber Loss	“Huber Errors”

# MAXIMUM LIKELIHOOD IN CLASSIFICATION

Let us assume the outputs  $y^{(i)}$  to be Bernoulli-distributed, i.e.

$$y^{(i)} \sim \text{Ber}(\pi(\mathbf{x}))$$

with probability  $\pi(\mathbf{x})$  that depends on  $\mathbf{x}$ .

The maximization of the negative log-likelihood is based on

$$\begin{aligned} -\ell(\boldsymbol{\theta}) &= -\sum_{i=1}^n \log p\left(y^{(i)} \mid \mathbf{x}^{(i)}, \boldsymbol{\theta}\right) \\ &= \sum_{i=1}^n -y^{(i)} \log[\pi(\mathbf{x}^{(i)})] - (1 - y^{(i)}) \log[1 - \pi(\mathbf{x}^{(i)})]. \end{aligned}$$

# MAXIMUM LIKELIHOOD IN CLASSIFICATION

This gives rise to the following loss function

$$L_{0,1}(y, \pi(\mathbf{x})) = -y \ln(\pi(\mathbf{x})) - (1 - y) \ln(1 - \pi(\mathbf{x}))$$

which we introduced as **Bernoulli** loss.

