


Introduction to Machine Learning

k -Nearest Neighbors






Learning goals

- Understand the basic idea of k -NN
- Know different distance measures for different scales of feature variables
- Understand that k -NN has no optimization step

K-NEAREST-NEIGHBORS

- k -**NN** can be used for regression and classification
- It generates predictions \hat{y} for a given \mathbf{x} by comparing the k observations that are closest to \mathbf{x}
- "Closeness" requires a distance or similarity measure (usually: Euclidean).
- The set containing the k closest points $\mathbf{x}^{(i)}$ to \mathbf{x} in the training data is called the k -**neighborhood** $N_k(\mathbf{x})$ of \mathbf{x} .




DISTANCE MEASURES

How to calculate distances?

- Most popular distance measure for numerical features:
Euclidean distance

- For two data points \mathbf{x} and $\tilde{\mathbf{x}}$ with p features $\in \mathbb{R}$

- the Euclidean distance is $d_{Euclidean}(\mathbf{x}, \tilde{\mathbf{x}}) = \sqrt{\sum_{j=1}^p (x_j - \tilde{x}_j)^2}$.
- the Manhattan distance is $d_{manhattan}(\mathbf{x}, \tilde{\mathbf{x}}) = \sum_{j=1}^p |x_j - \tilde{x}_j|$.



K-NN REGRESSION

Predictions for regression:

$$\hat{y} = \frac{1}{k} \sum_{i: \mathbf{x}^{(i)} \in N_k(\mathbf{x})} y^{(i)}$$

$$\hat{y} = \frac{1}{\sum_{i: \mathbf{x}^{(i)} \in N_k(\mathbf{x})} w_i} \sum_{i: \mathbf{x}^{(i)} \in N_k(\mathbf{x})} w_i y^{(i)}$$

with neighbors weighted according to their distance to \mathbf{x} : $w_i = \frac{1}{d(\mathbf{x}^{(i)}, \mathbf{x})}$

K-NN SUMMARY

- k -NN has no optimization step and is a very local model.
- We cannot simply use least-squares loss on the training data for picking k , because we would always pick $k = 1$.
- k -NN makes no assumptions about the underlying data distribution.
- The smaller k , the less stable, less smooth and more “wiggly” the decision boundary becomes.
- Accuracy of k -NN can be severely degraded by the presence of noisy or irrelevant features, or when the feature scales are not consistent with their importance.

DISTANCE MEASURES - EXTENSIONS

Categorical variables, missing data and mixed space:

The Gower distance $d_{gower}(\mathbf{x}, \tilde{\mathbf{x}})$ is a weighted mean of $d_{gower}(x_j, \tilde{x}_j)$:

$$d_{gower}(\mathbf{x}, \tilde{\mathbf{x}}) = \frac{\sum_{j=1}^p \delta_{x_j, \tilde{x}_j} \cdot d_{gower}(x_j, \tilde{x}_j)}{\sum_{j=1}^p \delta_{x_j, \tilde{x}_j}}.$$

- $\delta_{x_j, \tilde{x}_j}$ is 0 or 1. It becomes 0 when the j -th variable is **missing** in at least one of the observations (\mathbf{x} or $\tilde{\mathbf{x}}$), or when the variable is asymmetric binary (where “1” is more important/distinctive than “0”, e. g., “1” means “color-blind”) and both values are zero. Otherwise it is 1.

DISTANCE MEASURES - EXTENSIONS

- $d_{gower}(x_j, \tilde{x}_j)$, the j -th variable contribution to the total distance, is a distance between the values of x_j and \tilde{x}_j . For nominal variables the distance is 0 if both values are equal and 1 otherwise. The contribution of other variables is the absolute difference of both values, divided by the total range of that variable.

DISTANCE MEASURES - EXTENSIONS

Example of Gower distance with data on sex and income:

index	sex	salary
1	m	2340
2	w	2100
3	NA	2680

$$d_{gower}(\mathbf{x}, \tilde{\mathbf{x}}) = \frac{\sum_{j=1}^p \delta_{x_j, \tilde{x}_j} \cdot d_{gower}(x_j, \tilde{x}_j)}{\sum_{j=1}^p \delta_{x_j, \tilde{x}_j}}$$

$$d_{gower}(\mathbf{x}^{(1)}, \mathbf{x}^{(2)}) = \frac{1 \cdot 1 + 1 \cdot \frac{|2340 - 2100|}{|2680 - 2100|}}{1 + 1} = \frac{1 + \frac{240}{580}}{2} = \frac{1 + 0.414}{2} = 0.707$$

$$d_{gower}(\mathbf{x}^{(1)}, \mathbf{x}^{(3)}) = \frac{0 \cdot 1 + 1 \cdot \frac{|2340 - 2680|}{|2680 - 2100|}}{0 + 1} = \frac{0 + \frac{340}{580}}{1} = \frac{0 + 0.586}{1} = 0.586$$

$$d_{gower}(\mathbf{x}^{(2)}, \mathbf{x}^{(3)}) = \frac{0 \cdot 1 + 1 \cdot \frac{|2100 - 2680|}{|2680 - 2100|}}{0 + 1} = \frac{0 + \frac{580}{580}}{1} = \frac{0 + 1.000}{1} = 1$$

DISTANCE MEASURES - EXTENSIONS

Weights:

Weights can be used to address two problems in distance calculation:

- **Standardization:** Two features may have values with a different scale. Many distance formulas (not Gower) would place a higher importance on a feature with higher values, leading to an imbalance. Assigning a higher weight to the lower-valued feature can combat this effect.
- **Importance:** Sometimes one feature has a higher importance (e. g., more recent measurement). Assigning weights according to the importance of the feature can align the distance measure with known feature importance.

For example:

$$d_{Euclidean}^{weighted}(\mathbf{x}, \tilde{\mathbf{x}}) = \sqrt{\sum_{j=1}^p w_j (x_j - \tilde{x}_j)^2}$$