




Figure 1. Historical locations and number of accepted papers for International Machine Learning Conferences (ICML 1993 – ICML 2008) and International Workshops on Machine Learning (ML 1988 – ML 1992). At the time this figure was produced, the number of accepted papers for ICML 2008 was unknown and instead estimated.

Algorithm 1 Bubble Sort

Input: data x_i , size m

repeat

Initialize $noChange = true$.

for $i = 1$ **to** $m - 1$ **do**

if $x_i > x_{i+1}$ **then**

Swap x_i and x_{i+1}

$noChange = false$

end if

end for

until $noChange$ is $true$

column, and you may set wide figures across both columns (use the environment `figure*` in $\LaTeX$). Always place two-column figures at the top or bottom of the page.

2.7. Algorithms

If you are using $\LaTeX$, please use the “algorithm” and “algorithmic” environments to format pseudocode. These require the corresponding stylefiles, `algorithm.sty` and `algorithmic.sty`, which are supplied with this package. Algorithm 1 shows an example.

2.8. Tables

You may also want to include tables that summarize material. Like figures, these should be centered, legible, and numbered consecutively. However, place the title *above* the

Table 1. Classification accuracies for naive Bayes and flexible Bayes on various data sets.

DATA SET	NAIVE	FLEXIBLE	BETTER?
BREAST	95.9 ± 0.2	96.7 ± 0.2	✓
CLEVELAND	83.3 ± 0.6	80.0 ± 0.6	×
GLASS2	61.9 ± 1.4	83.8 ± 0.7	✓
CREDIT	74.8 ± 0.5	78.3 ± 0.6	
HORSE	73.3 ± 0.9	69.7 ± 1.0	×
META	67.1 ± 0.6	76.5 ± 0.5	✓
PIMA	75.1 ± 0.6	73.9 ± 0.5	
VEHICLE	44.9 ± 0.6	61.5 ± 0.4	✓

table with at least 0.1 inches of space before the title and the same after it, as in Table 1. The table title should be set in 9 point type and centered unless it runs two or more lines, in which case it should be flush left.

Tables contain textual material, whereas figures contain graphical material. Specify the contents of each row and column in the table’s topmost row. Again, you may float tables to a column’s top or bottom, and set wide tables across both columns. Place two-column tables at the top or bottom of the page.

2.9. Theorems and such

The preferred way is to number definitions, propositions, lemmas, etc. consecutively, within sections, as shown below.

Definition 2.1. A function $f : X \rightarrow Y$ is injective if for any $x, y \in X$ different, $f(x) \neq f(y)$.

Using Definition 2.1 we immediately get the following result:

Proposition 2.2. If f is injective mapping a set X to another set Y , the cardinality of Y is at least as large as that of X

Proof. Left as an exercise to the reader. $\square$

Lemma 2.3 stated next will prove to be useful.

Lemma 2.3. For any $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ injective functions, $f \circ g$ is injective.

Theorem 2.4. If $f : X \rightarrow Y$ is bijective, the cardinality of X and Y are the same.

An easy corollary of Theorem 2.4 is the following:

Corollary 2.5. If $f : X \rightarrow Y$ is bijective, the cardinality of X is at least as large as that of Y .

Assumption 2.6. The set X is finite.

Remark 2.7. According to some, it is only the finite case (cf. Assumption 2.6) that is interesting.