

MA2252 Introduction to Computing

Lecture 15 Taylor series

Sharad Kumar Keshari

School of Computing and Mathematical Sciences

University of Leicester

Learning outcomes

At the end of lecture, students will be able to

- understand Taylor polynomials
- understand Taylor series
- find Taylor series approximations and estimate error

Many functions in Mathematics can be approximated by polynomials upto desired accuracy.

To perform such approximations, we need to understand **Taylor polynomials** of a function.

Taylor polynomials

A Taylor polynomial of a function $f(x)$ centered at $x = a$ is a polynomial approximation of $f(x)$.




Figure: Brook Taylor

Taylor polynomials (contd.)

Question: Find a quadratic polynomial to approximate $f(x)$ near $x = a$.

$$\begin{aligned} p_2(x) &= \frac{f''(a)}{2} (x-a)^2 + f'(a)(x-a) + f(a) \\ &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2} (x-a)^2 \end{aligned}$$

Taylor polynomials (contd.)

A n th order Taylor Polynomial is given as

$$p_n(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^n(a)}{n!}(x-a)^n. \quad (1)$$

- $p_0(x) = f(a)$ (constant function)
 - $p_1(x) = f(a) + \frac{f'(a)}{1!}(x-a)$ (linear approximation)
 - $p_2(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2$
(quadratic approximation)
- Do it as an exercise*

Taylor polynomials (contd.)

Example: The Taylor polynomials of $f(x) = \cos x$ upto degree 6:

- $n = 0$: $p_0(x) = 1$
- $n = 2$: $p_2(x) = 1 - \frac{x^2}{2}$
- $n = 4$: $p_4(x) = 1 - \frac{x^2}{2} + \frac{x^4}{24}$
- $n = 6$: $p_6(x) = 1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720}$

at $n=0$
($a=0$)

~~$p_0(x)$~~
 $p_0(x) = f(0) = \cos 0 = 1$

$p_1(x) = f(0) + \frac{f'(0)}{1!}(x-0)$

$= 1 + f'(0)x$

$f'(x) = -\sin x$
 $f'(0) = 0$

$p_2(x) = f(0) + \frac{f'(0)}{1!}(x-0) + \frac{f''(0)}{2!}(x-0)^2$

Note: It is possible that a n th order Taylor polynomial is not of degree n .


$= 1 - \frac{x^2}{2}$

Taylor polynomials (contd.)

Example: Write a script file to plot the Taylor polynomials of $\cos x$ upto degree 6.

Demo

Taylor polynomials (contd.)



Taylor series

A Taylor series is an infinite series expansion of a function at a given point in its domain.

Taylor series of a real-valued function $f(x)$ at $x = a$ is defined as

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots \quad (2)$$

or

$$f(x) = \sum_{n=0}^{\infty} \frac{f^n(a)}{n!}(x-a)^n \quad (3)$$

Taylor series (contd.)

→ special case of
Taylor series

When $a = 0$, we obtain a **Maclaurin series**.

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \dots \quad (4)$$

Taylor series (contd.)

Example: Find the Taylor series of $\cos x$ at $x = 0$.

$$\begin{aligned} f(x) &= f(0) + f'(0)x + \frac{f''(0)x^2}{2!} \\ &\quad + \frac{f'''(0)x^3}{3!} + \frac{f^{(4)}(0)x^4}{4!} + \dots \\ \sin x &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \end{aligned}$$

Taylor's theorem

Theorem

Suppose a function $f(x)$ has $n + 1$ continuous derivatives in an open interval I containing $x = a$ then $\forall n$ and $\forall x \in I$,

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + R_n(x) \quad (5)$$

where

$$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!}(x-a)^{n+1} \quad (6)$$

for some c between a and x .

\downarrow n th order
Taylor
polynomial
 $R_n(x) \rightarrow 0$ as $n \rightarrow \infty$

Here, $R_n(x)$ is called the Taylor remainder or error term.

Error estimation

We don't know the exact value of $R_n(x)$ since exact value of c is unknown. However, an upper bound on the error term can still be found.

If $|f^{n+1}(x)| \leq M$ for all t between a and x , then

$$|R_n(x)| \leq \frac{M}{(n+1)!} |x - a|^{n+1} \quad (7)$$

✓
Lagrange
upper
bound on remainder

Error estimation (contd.)

$$\cos x = \underbrace{1 - \frac{x^2}{2!}}_{P_2(x)} + \underbrace{\frac{x^4}{4!} + \dots}_{R_2(x)}$$

2nd order Taylor polynomial near $x=0$
 $1 - \frac{(x-0)^2}{2}$

Example: Find the maximum error if $p_2(x) = 1 - \frac{x^2}{2}$ is used to estimate the value of $\cos(x)$ at $x = 0.3$. Verify that the error estimate in MATLAB is less than the maximum error.

$$\begin{aligned} f(x) &= \cos x \\ f'(x) &= -\sin x \\ f''(x) &= -\cos x \\ f'''(x) &= \sin x \end{aligned} \quad \& \quad R_2(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x-a)^{n+1}$$
$$= \frac{f^{(3)}(c) x^3}{3!} = \frac{\sin c \cdot x^3}{6}$$

Error estimation (contd.)

$$-1 \leq \text{sinc} \leq 1$$
$$|\text{sinc}| \leq 1 \quad m=1$$

$$|R_2(u)| = |\text{sinc}| |u|^3$$

$$\leq \frac{|u|^3}{6} = \frac{(0.3)^3}{6}$$
$$= \frac{0.027}{6}$$

$$= 0.0045$$

↓
max
error

$$0 < u < 0.3$$
$$\text{sinc} < \text{sinc}(0.3)$$

Demo

$$\sin(0.3)$$

Some applications:

- Small-angle approximations e.g. $\sin \theta \approx \theta$ for small values of θ .
- Finding non-elementary definite integrals e.g. $\int_1^2 \frac{\sin x}{x} dx$
- Deriving formulas for numerical differentiation and integration (we'll study later in this course)

End of Lecture 15

Please provide your feedback [▶ here](#)