

MA1014 CALCULUS AND ANALYSIS TUTORIAL 18

L

Dr. Andrew Tonks: apt12@le.ac.uk

Ben Smith: bjs30@le.ac.uk

SUBSEQUENCES

If $(a_n)_{n \geq 1}$ is a sequence, then a subsequence of a_n is a sequence

$$(b_k)_{k \geq 1} : \forall k \in \mathbb{N}, \exists n_k \in \mathbb{N} : n_1 < n_2 < \dots < n_k \text{ with } b_k = a_{n_k}$$

e.g. $a_n = n^2 : n_k = 2k, b_k = a_{n_k} = a_{2k} = (2k)^2 = 4k^2$

Bolzano-Weierstrass Theorem:

If $(a_n)_{n \geq 1}$ is a bounded sequence, then there exists a convergent subsequence $(a_{n_k})_{k \geq 1}$

CAUCHY SEQUENCES

- A sequence $(a_n)_{n \geq 1}$ is Cauchy if

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} : \forall n, m \geq N \Rightarrow |a_n - a_m| < \varepsilon$$

- Every Cauchy sequence is bounded
- A sequence is convergent iff it is Cauchy


EXERCISE

Consider the sequence $(a_n)_{n \geq 1} = (-1)^n$

- a) Is a_n bounded? Is it monotonic? Explain your answers.
- b) Show that a_n diverges.
- c) Does there exist a convergent subsequence $(b_k)_{k \geq 1} = a_{n_k}$?
If so, explain why and give an example of such a subsequence.
- d) Show that your chosen b_k is a Cauchy Sequence. Why is a_n not a Cauchy sequence?

EXERCISE

- 1) Consider $x^3 = 3$.
 - a) Write down Newton's method to solve this equation.
 - b) By choosing $x_0 = 3$, prove by induction that the sequence is monotonic decreasing and bounded below.
 - c) Hence, why does the sequence converge, and find the limit.
- 2) Let $(a_n)_{n \geq 1}$ be a sequence, and $(a_{n_k})_{k \geq 1}$ be a subsequence. Prove by induction that $n_k \geq k$, $\forall k \geq 1$.
- 3) Show that all bounded divergent sequences possess at least two convergent subsequences.



$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

$$\int_a^b f(x) dx = F(b) - F(a)$$

1

L

ANY QUESTIONS?

$$m \frac{d^2 x}{dt^2} = -kx$$

$$\int \frac{dx}{1+x^2} = \tan^{-1}(x) + C$$

