

Solutions for Tutorial Problem Sheet 12, December 15.

(Infinite Sequences and Series)

Problem 1.

- a) Use the Ratio Test to determine if the series $\sum_{n=1}^{\infty} (-1)^n \frac{n^2(n+2)!}{n! 3^{2n}}$ converges absolutely or diverges.

Solution:

$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} \frac{(n+1)^2((n+1)+2)!}{(n+1)! 3^{2(n+1)}}}{(-1)^n \frac{n^2(n+2)!}{n! 3^{2n}}} \right| &= \lim_{n \rightarrow \infty} \left(\frac{(n+1)^2(n+3)(n+2)!}{(n+1) \cdot n! 3^{2n} \cdot 3^2} \cdot \frac{n! 3^{2n}}{n^2(n+2)!} \right) = \lim_{n \rightarrow \infty} \left(\frac{n^3 + 5n^2 + 7n + 3}{9n^3 + 9n^2} \right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{3n^2 + 15n + 7}{27n^2 + 18n} \right) = \lim_{n \rightarrow \infty} \left(\frac{6n + 15}{54n + 18} \right) = \lim_{n \rightarrow \infty} \left(\frac{6}{54} \right) = \frac{1}{9} < 1 \Rightarrow \sum_{n=1}^{\infty} (-1)^n \frac{n^2(n+2)!}{n! 3^{2n}} \text{ converges} \end{aligned}$$

- b) Use the Root Test to determine if the series $\sum_{n=1}^{\infty} \left(\frac{4n+3}{3n-5} \right)^n$ converges absolutely or diverges.

Solution:

$$\lim_{n \rightarrow \infty} \sqrt[n]{\left| \left(\frac{4n+3}{3n-5} \right)^n \right|} = \lim_{n \rightarrow \infty} \left(\frac{4n+3}{3n-5} \right) = \lim_{n \rightarrow \infty} \left(\frac{4}{3} \right) = \frac{4}{3} > 1 \Rightarrow \sum_{n=1}^{\infty} \left(\frac{4n+3}{3n-5} \right)^n \text{ diverges}$$

- c) Use any method to determine if the series $\sum_{n=1}^{\infty} \frac{n^{10}}{10^n}$ converges or diverges. Give reasons for your answer.

Solution:

$$\text{converges by the Ratio Test: } \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{\left(\frac{(n+1)^{10}}{10^{n+1}} \right)}{\left(\frac{n^{10}}{10^n} \right)} = \lim_{n \rightarrow \infty} \frac{(n+1)^{10}}{10^{n+1}} \cdot \frac{10^n}{n^{10}} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^{10} \left(\frac{1}{10} \right) = \frac{1}{10} < 1$$

Problem 2. Determine if the series converges absolutely, converges, or diverges? Give reasons for your answers.

a) $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3+n}{5+n},$

Solution:

diverges by the n th-Term Test since $\lim_{n \rightarrow \infty} \frac{3+n}{5+n} = 1 \neq 0$

b) $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} (n!)^2}{(2n)!}.$

Solution:

converges absolutely by the Ratio Test: $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{((n+1)!)^2}{((2n+2)!)^2} \cdot \frac{(2n)!}{(n!)^2} = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{(2n+2)(2n+1)} = \frac{1}{4} < 1$

Problem 3. Use any method to determine whether the series converges or diverges. Give reasons for your answer.

a) $\sum_{n=0}^{\infty} (-1)^n \frac{(n+2)!}{(2n)!}$,

Solution:

converges by the Ratio Test: $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{(n+3)!}{(2(n+1))!} \cdot \frac{(2n)!}{(n+2)!} = \lim_{n \rightarrow \infty} \frac{n+3}{(2n+2)(2n+1)} = 0 < 1$

b) $\sum_{n=2}^{\infty} \frac{3}{10+n^{4/3}}$.

Solution:

converges by the Direct Comparison Test: $\frac{3}{10+n^{4/3}} < \frac{3}{n^{4/3}}$, which is the n th term of a convergent p -series

Problem 4. Given a series $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{n!}$

a) find the series' radius and interval of convergence.

For what values of x does the series converge b) absolutely, c) conditionally?

Solution:

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| < 1 \Rightarrow \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n} \right| < 1 \Rightarrow |x| \lim_{n \rightarrow \infty} \left(\frac{1}{n+1} \right) < 1 \text{ for all } x$$

- (a) the radius is ∞ ; the series converges for all x
- (b) the series converges absolutely for all x
- (c) there are no values for which the series converges conditionally

Problem 5.

- a) Find the interval of convergence of the power series $\sum_{n=0}^{\infty} \frac{8}{4^{n+2}} x^n$.
b) Represent the power series in part a) as a power series about $x = 3$ and identify the interval of convergence of the new series.

Solution:

We can write the given series as $\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{x}{4}\right)^n$ which shows that the interval of convergence is $-4 < x < 4$.

The function represented by the series in (a) is $\frac{2}{4-x}$ for $-4 < x < 4$. If we rewrite this function as

$\frac{2}{1-(x-3)}$ we can represent it by the geometric series $\sum_{n=0}^{\infty} 2(x-3)^n$ which will converge only for $|x-3| < 1$ or $2 < x < 4$.