

Lecture 20: Partial Derivatives.

MA2032 Vector Calculus

Lecturer: Larissa Serdukova

School of Computing and Mathematical Science
University of Leicester

November 4, 2022

Taylor's Formula for Two Variables

- The use of **Taylor's formula** leads to an extension of the formula that provides **polynomial approximations** of all orders for functions of two independent variables.

Taylor's Formula for $f(x, y)$ at the Point (a, b)

Suppose $f(x, y)$ and its partial derivatives through order $n + 1$ are continuous throughout an open rectangular region R centered at a point (a, b) . Then, throughout R ,

$$\begin{aligned} f(a + h, b + k) = & f(a, b) + (hf_x + kf_y) \Big|_{(a, b)} + \frac{1}{2!} (h^2 f_{xx} + 2hk f_{xy} + k^2 f_{yy}) \Big|_{(a, b)} \\ & + \frac{1}{3!} (h^3 f_{xxx} + 3h^2 k f_{xxy} + 3hk^2 f_{xyy} + k^3 f_{yyy}) \Big|_{(a, b)} + \cdots + \frac{1}{n!} \left(h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y} \right)^n f \Big|_{(a, b)} \\ & + \frac{1}{(n + 1)!} \left(h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y} \right)^{n+1} f \Big|_{(a+ch, b+ck)}. \end{aligned} \quad (7)$$

- The **first n derivative terms** are evaluated at (a, b) . The **last term** is evaluated at some point $(a + ch, b + ck)$ on the line segment joining (a, b) and $(a + h, b + k)$.

Taylor's Formula for Two Variables

- If $(a, b) = (0, 0)$ and we treat h and k as independent variables (denoting them now by x and y), then Equation (7) assumes the following form.

Taylor's Formula for $f(x, y)$ at the Origin

$$\begin{aligned} f(x, y) = & f(0, 0) + xf_x + yf_y + \frac{1}{2!}(x^2f_{xx} + 2xyf_{xy} + y^2f_{yy}) \\ & + \frac{1}{3!}(x^3f_{xxx} + 3x^2yf_{xxy} + 3xy^2f_{xyy} + y^3f_{yyy}) + \cdots + \frac{1}{n!}\left(x^n\frac{\partial^n f}{\partial x^n} + nx^{n-1}y\frac{\partial^n f}{\partial x^{n-1}\partial y} + \cdots + y^n\frac{\partial^n f}{\partial y^n}\right) \\ & + \frac{1}{(n+1)!}\left(x^{n+1}\frac{\partial^{n+1} f}{\partial x^{n+1}} + (n+1)x^ny\frac{\partial^{n+1} f}{\partial x^n\partial y} + \cdots + y^{n+1}\frac{\partial^{n+1} f}{\partial y^{n+1}}\right)\Bigg|_{(cx, cy)} \end{aligned} \quad (8)$$

- The first n derivative terms are evaluated at $(0, 0)$. The last term is evaluated at a point on the line segment joining the origin and (x, y) .
- Taylor's formula provides **polynomial approximations** of two-variable functions. The first n derivative terms give the polynomial; the **last term** gives the **approximation error**.
- The first three terms of Taylor's formula give the **function's linearization**. To improve on the linearization, we add higher-power terms.

Taylor's Formula for Two Variables

Example 1

Find a quadratic approximation to $f(x, y) = \sin x \sin y$ near the origin. How accurate is the approximation if $|x| \leq 0.1$ and $|y| \leq 0.1$?

Solution We take $n = 2$ in Equation (8):

$$\begin{aligned} f(x, y) = f(0, 0) + (xf_x + yf_y) + \frac{1}{2}(x^2f_{xx} + 2xyf_{xy} + y^2f_{yy}) \\ + \frac{1}{6}(x^3f_{xxx} + 3x^2yf_{xxy} + 3xy^2f_{xyy} + y^3f_{yyy}) \Big|_{(cx, cy)}. \end{aligned}$$

Calculating the values of the partial derivatives,

$$f(0, 0) = \sin x \sin y \Big|_{(0, 0)} = 0, \quad f_{xx}(0, 0) = -\sin x \sin y \Big|_{(0, 0)} = 0,$$


$$f_x(0, 0) = \cos x \sin y \Big|_{(0, 0)} = 0, \quad f_{xy}(0, 0) = \cos x \cos y \Big|_{(0, 0)} = 1,$$

$$f_y(0, 0) = \sin x \cos y \Big|_{(0, 0)} = 0, \quad f_{yy}(0, 0) = -\sin x \sin y \Big|_{(0, 0)} = 0,$$

we have the result

$$\sin x \sin y \approx 0 + 0 + 0 + \frac{1}{2}(x^2(0) + 2xy(1) + y^2(0)), \quad \text{or} \quad \sin x \sin y \approx xy.$$

Example 1



The error in the approximation is

$$E(x, y) = \frac{1}{6} (x^3 f_{xxx} + 3x^2 y f_{xxy} + 3xy^2 f_{xyy} + y^3 f_{yyy}) \Big|_{(cx, cy)}.$$

The third derivatives never exceed 1 in absolute value because they are products of sines and cosines. Also, $|x| \leq 0.1$ and $|y| \leq 0.1$. Hence

$$|E(x, y)| \leq \frac{1}{6} ((0.1)^3 + 3(0.1)^3 + 3(0.1)^3 + (0.1)^3) = \frac{8}{6} (0.1)^3 \leq 0.00134$$

(rounded up). The error will not exceed 0.00134 if $|x| \leq 0.1$ and $|y| \leq 0.1$.



Taylor's Formula for Two Variables

Example 2

Use Taylor's formula for $f(x, y) = \ln(2x + y + 1)$ at the origin to find quadratic and cubic approximations of f near the origin.

$$f(x, y) = \ln(2x + y + 1) \Rightarrow f_x = \frac{2}{2x+y+1}, f_y = \frac{1}{2x+y+1}, f_{xx} = \frac{-4}{(2x+y+1)^2}, f_{xy} = \frac{-2}{(2x+y+1)^2}, f_{yy} = \frac{-1}{(2x+y+1)^2}$$

$$\Rightarrow f(x, y) \approx f(0, 0) + xf_x(0, 0) + yf_y(0, 0) + \frac{1}{2} \left[x^2 f_{xx}(0, 0) + 2xy f_{xy}(0, 0) + y^2 f_{yy}(0, 0) \right]$$

$$= 0 + x \cdot 2 + y \cdot 1 + \frac{1}{2} \left[x^2 \cdot (-4) + 2xy \cdot (-2) + y^2 \cdot (-1) \right] = 2x + y + \frac{1}{2} (-4x^2 - 4xy - y^2)$$

$$= (2x + y) - \frac{1}{2} (2x + y)^2, \text{ quadratic approximation;}$$

$$f_{xxx} = \frac{16}{(2x+y+1)^3}, f_{xxy} = \frac{8}{(2x+y+1)^3}, f_{xyy} = \frac{4}{(2x+y+1)^3}, f_{yyy} = \frac{2}{(2x+y+1)^3}$$


$$\Rightarrow f(x, y) \approx \text{quadratic} + \frac{1}{6} \left[x^3 f_{xxx}(0, 0) + 3x^2 y f_{xxy}(0, 0) + 3xy^2 f_{xyy}(0, 0) + y^3 f_{yyy}(0, 0) \right]$$

$$= (2x + y) - \frac{1}{2} (2x + y)^2 + \frac{1}{6} \left(x^3 \cdot 16 + 3x^2 y \cdot 8 + 3xy^2 \cdot 4 + y^3 \cdot 2 \right)$$


$$= (2x + y) - \frac{1}{2} (2x + y)^2 - \frac{1}{3} (8x^3 + 12x^2 y + 6xy^2 + y^3)$$

$$= (2x + y) - \frac{1}{2} (2x + y)^2 + \frac{1}{3} (2x + y)^3, \text{ cubic approximation}$$

Example 2



Example 2



Taylor's Formula for Two Variables

Example 3

Use Taylor's formula for $f(x, y) = \frac{1}{1-x-y+xy}$ at the origin to find quadratic and cubic approximations of f near the origin.

$$f(x, y) = \frac{1}{1-x-y+xy} \Rightarrow f_x = \frac{1-y}{(1-x-y+xy)^2}, f_y = \frac{1-x}{(1-x-y+xy)^2}, f_{xx} = \frac{2(1-y)^2}{(1-x-y+xy)^3}, f_{xy} = \frac{1}{(1-x-y+xy)^2},$$

$$f_{yy} = \frac{2(1-x)^2}{(1-x-y+xy)^3} \Rightarrow f(x, y) \approx f(0, 0) + x f_x(0, 0) + y f_y(0, 0) + \frac{1}{2} \left[x^2 f_{xx}(0, 0) + 2xy f_{xy}(0, 0) + y^2 f_{yy}(0, 0) \right]$$

$$= 1 \cdot x \cdot 1 + y \cdot 1 + \frac{1}{2} \left(x^2 \cdot 2 + 2xy \cdot 1 + y^2 \cdot 2 \right) = 1 + x + y + x^2 + xy + y^2, \text{ quadratic approximation;}$$


$$f_{xxx} = \frac{6(1-y)^3}{(1-x-y+xy)^4}, f_{xxy} = \frac{[-4(1-x-y+xy) + 6(1-y)(1-x)](1-y)}{(1-x-y+xy)^4}, f_{xyy} = \frac{[-4(1-x-y+xy) + 6(1-x)(1-y)](1-x)}{(1-x-y+xy)^4},$$

$$f_{yyy} = \frac{6(1-x)^3}{(1-x-y+xy)^4} \Rightarrow f(x, y) \approx \text{quadratic} + \frac{1}{6} \left[x^3 f_{xxx}(0, 0) + 3x^2 y f_{xxy}(0, 0) + 3xy^2 f_{xyy}(0, 0) + y^3 f_{yyy}(0, 0) \right]$$


$$= 1 + x + y + x^2 + xy + y^2 + \frac{1}{6} \left[x^3 \cdot 6 + 3x^2 y \cdot 2 + 3xy^2 \cdot 2 + y^3 \cdot 6 \right]$$

$$= 1 + x + y + x^2 + xy + y^2 + x^3 + x^2 y + xy^2 + y^3, \text{ cubic approximation}$$

Example 3



Example 3



Taylor's Formula for Two Variables

Example 4

Use Taylor's formula to find a quadratic approximation of $f(x, y) = e^x \sin y$ at the origin. Estimate the error in the approximation if $|x| \leq 0.1$ and $|y| \leq 0.1$.

$$\begin{aligned} f(x, y) = e^x \sin y &\Rightarrow f_x = e^x \sin y, f_y = e^x \cos y, f_{xx} = e^x \sin y, f_{xy} = e^x \cos y, f_{yy} = -e^x \sin y \\ &\Rightarrow f(x, y) \approx f(0, 0) + xf_x(0, 0) + yf_y(0, 0) + \frac{1}{2} \left[x^2 f_{xx}(0, 0) + 2xy f_{xy}(0, 0) + y^2 f_{yy}(0, 0) \right] \\ &= 0 + x \cdot 0 + y \cdot 1 + \frac{1}{2} (x^2 \cdot 0 + 2xy \cdot 1 + y^2 \cdot 0) = y + xy, \text{ quadratic approximation.} \end{aligned}$$

Now, $f_{xxx} = e^x \sin y$, $f_{xxy} = e^x \cos y$, $f_{xyy} = -e^x \sin y$, and $f_{yyy} = -e^x \cos y$.

Since $|x| \leq 0.1$, $|e^x \sin y| \leq e^{0.1} \sin 0.1 \approx 0.11$ and $|e^x \cos y| \leq e^{0.1} \cos 0.1 \approx 1.11$.

Therefore, $E(x, y) \leq \frac{1}{6} \left[(0.11)(0.1)^3 + 3(1.11)(0.1)^3 + 3(0.11)(0.1)^3 + (1.11)(0.1)^3 \right] \leq 0.000814$.

Example 4

