

Nepal PreTST 2026 Solutions

PRAKRIT GAJUREL

28 June 2026

§1 Problems

Problem 1.1. Let $a_1, a_2, \dots, a_{2n+2}$ be positive real numbers. Prove that they can be permuted into numbers $u_0, v_0, u_1, v_1, \dots, u_n, v_n$ such that the polynomial

$$P(x) = u_0 - v_0x + u_1x^2 - v_1x^3 + \dots + u_nx^{2n} - v_nx^{2n+1}$$

has at least one real root, and every real root of P is greater than or equal to 1.

Problem 1.2. Find all functions $f : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$ such that for all $m \in \mathbb{N}$,

$$f(m-1) \mid m \quad \text{and} \quad m+1 \mid T_{2026}(m),$$

where $T_1(m) = f(m)$ and $T_{k+1}(m) = f(m + T_k(m))$ for all $k \geq 1$.

Problem 1.3. Let ABC be an acute scalene triangle with $\angle BAC = 60^\circ$. Denote by O and H the circumcenter and orthocenter of $\triangle ABC$ respectively. Let M be the midpoint of arc BC on (ABC) not containing A , and let ray $\overrightarrow{MH}$ intersect (ABC) a second time at P . Show that the lines AP , OH and BC pass through a common point.

Problem 1.4. Consider the standard triangular grid of unit equilateral triangles forming a larger equilateral triangle of side length n . Initially, only the top unit triangle is colored black, and all other unit triangles are colored white.

A **move** consists of choosing a black unit triangle and simultaneously flipping the colors of **all** unit triangles that share an edge with it; that is, each such neighboring triangle changes from white to black or from black to white. Determine all positive integers n for which it is possible, after a sequence of moves, for all unit triangles in the grid to be black at the same time.

Problem 1.5. Let $\sigma(n)$ denote the divisors sum of n , and let $d(n)$ denote the number of divisors of n . Prove that for any nonnegative integer k , the equation $\sigma(n) = d(n)^k$ has finitely many solutions.

Problem 1.6. Let ω_1 and ω_2 be two circles intersecting at distinct points X and Y . Let ℓ be their common tangent closer to Y , intersecting ω_1 at A and ω_2 at B . The tangent to ω_1 at Y meets ω_2 again at C with $C \neq Y$. Let the line XB meet ω_1 again at K with $K \neq X$. Let $D = BC \cap AY$, and let $E = YK \cap CX$.

Prove that D, Y, X, E are concyclic. (Four points are concyclic if there exists a circle containing all four points.)

Problem 1.7. For each positive integer n , let $\kappa(n)$ denote the unique squarefree positive integer such that $\frac{n}{\kappa(n)}$ is a perfect square. Let G be a connected simple graph whose

edges are labeled by positive integers w_e . Assume that for every cycle C in G , the product $\prod_{e \in C} w_e$ is a perfect square. Prove that there exist squarefree positive integers s_v assigned to the vertices $v \in V(G)$ such that for every edge $uv \in E(G)$ there exists a positive integer t_{uv} with $w_{uv} = \kappa(s_u s_v) t_{uv}^2$.

Problem 1.8. For all positive reals a, b, c , prove the inequality

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2} + \frac{(a-b)^2 + (b-c)^2 + (c-a)^2}{2(a+b+c)^2}.$$

Determine when equality holds.

§2 Solutions

Problem 2.1. Let $a_1, a_2, \dots, a_{2n+2}$ be positive real numbers. Prove that they can be permuted into numbers $u_0, v_0, u_1, v_1, \dots, u_n, v_n$ such that the polynomial

$$P(x) = u_0 - v_0x + u_1x^2 - v_1x^3 + \dots + u_nx^{2n} - v_nx^{2n+1}$$

has at least one real root, and every real root of P is greater than or equal to 1.

Proof. Obviously P has one real root due to it having an odd degree. We then group the numbers into $n + 1$ pairs like $(u_0, v_0), (u_1, v_1), \dots, (u_n, v_n)$ such that for each $0 \leq k \leq n$, $u_k \geq v_k$. We now observe the behavior of the following expression:

$$u_kx^{2k} - v_kx^{2k+1} = x^{2k}((u_k - v_k) + v_k(1 - x))$$

Which, for $x < 1$, is nonnegative. Moreover, consider the case when $x = 0$, we get for $k = 0$ that $u_0 - v_0(0) = u_0 > 0$ so at least one of these pairs is strictly positive for all $x < 1$ and we conclude that

$$P(x) > 0 \text{ for all } x < 1.$$

FTSOC, if $r < 1$ was a real root of P , we would have $P(r) = 0$ and $P(r) > 0$, contradiction. Therefore every real root r of P satisfies $r \geq 1$. $\square$