

# Functional Equations Solutions

PRAKRIT GAJUREL

5 July 2026

## §1 Walkthrough Problems

**Problem 1.1** (USAMO 2002). Determine all functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f(x^2 - y^2) = xf(x) - yf(y)$$

for all pairs of real numbers  $x$  and  $y$ .

*Proof.* Let  $P(x, y)$  denote the given assertion. Note that  $P(0, 0)$  gives us  $f(0) = 0$ . Then,

$$\begin{aligned} P(x, -x) &\implies f(x^2 - x^2) = xf(x) + xf(-x) \\ 0 &= x(f(x) + f(-x)) \end{aligned}$$

For nonzero  $x$ , we have  $f(-x) = -f(x)$ , but when  $x = 0$  we have  $f(0) = 0$  anyway, so we have  $f$  is odd for all real numbers  $x$ .

$$P(x, 0) \implies f(x^2) = xf(x)$$

So now we can rewrite the original condition as:

$$f(x^2 - y^2) = f(x^2) - f(y^2)$$

We let  $a = x^2$ , and  $b = y^2$  such that, for nonnegative  $a, b$  we have:

$$f(a - b) = f(a) - f(b)$$

Again, we can set  $a = x_0 + y_0$  and  $b = y_0$  to get:

$$f(x_0 + y_0 - y_0) = f(x_0 + y_0) - f(y_0) \implies f(x_0 + y_0) = f(x_0) + f(y_0)$$

for nonnegative  $x_0, y_0$ . However, since  $f$  is odd, we get that  $f$  is additive on the negatives too.

Finally, we let  $x = a + b$  in one of the previous equations  $f(x^2) = xf(x)$  to get:

$$f(a^2 + 2ab + b^2) = (a + b)(f(a) + f(b))$$

$$f(2ab) = af(b) + bf(a)$$

$$a = 1 \implies f(2b) = f(b) + f(1)b = 2f(b)$$

Which gives  $f(x) = cx$  for all  $x$ , which upon checking works.  $\square$

**Problem 1.2** (IMO 2017). Solve over  $\mathbb{R}$  the functional equation

$$f(f(x)f(y)) + f(x+y) = f(xy).$$

*Proof.* We claim that  $f(x) = 0$ ,  $f(x) = 1 - x$ , and  $f(x) = x - 1$  are the only solutions. It's easy to check that these solutions indeed work.

Let  $P(x, y)$  denote the given assertion. First, note:

$$P(0, 0) \implies f(f(0)^2 = 0) \implies \text{there exists a } z \text{ such that } f(z) = 0.$$

Also,  $P(x, 0)$  gives that  $f(0) = 0 \implies f \equiv 0$ . So from now assume that  $f(0) \neq 0$ . We take, for  $z \neq 1$ ,

$$P\left(z, \frac{z}{z-1}\right) \implies f\left(f(z)f\left(\frac{z}{z-1}\right)\right) = 0$$

giving  $f(0) = 0$ , which is a contradiction, giving  $z = f(0)^2 = 1 \implies f(0) = \pm 1$ . One can now take  $P(1, x)$  to give  $f(x+1) = f(x) - 1$ , then  $P(x, 0)$  to get  $f(f(x)) = 1 - f(x)$ . Applying  $f$  on this equation and equating some terms gives us:

$$f(f(f(x))) = f(1 - f(x)) = 1 - f(f(x)) = f(x)$$

Before we move on, I'd like to prove something that would help us in our final claim.

**Claim 1.3** — For sufficiently large  $N_1$  and  $N_2$ , we can find  $x, y$  satisfying  $x + y = N_1 + 1$  and  $xy = N_2$ .

*Proof.* First we observe that we can think of  $x, y$  as roots of some quadratic equation. Of course, comparing it with Vieta's gives us that  $x$  and  $y$  are possible roots of the equation

$$x_0^2 - (N_1 + 1)x_0 + N_2 = 0$$

Obviously distinct solutions  $x, y$  to this equation exist if and only if the discriminant,  $\Delta$  is greater than zero. So we have:

$$\begin{aligned} \Delta &= (N_1 + 1)^2 - 4N_2 \\ &= N_1^2 + 2N_1 + 1 - 4N_2 > 0 \end{aligned}$$

So we can just pick sufficiently large  $N_1, N_2$  that satisfies the condition (or maybe something like  $N_1^2 \geq 4N_2$ ) so we have solutions for  $x, y$ .  $\square$

Now, we prove the final claim.

**Claim 1.4** —  $f$  is injective.

*Proof.* Assume for some sufficiently large  $a, b$ , we have  $f(a) = f(b)$  and also,  $x + y = a + 1$  and  $xy = b$  (from the above claim, we can do this). Then, the original assertion gives us

$$\begin{aligned} f(f(x)f(y)) &= f(b) - f(a+1) \\ &= f(b) - (f(a) - 1) \\ &= 1 \end{aligned}$$

Shifting  $f$  we get that  $f(f(x)f(y) + 1) = 0 \implies f(x)f(y) + 1 = 1$  so  $f(x)f(y) = 0$ , giving that  $1 \in \{x, y\}$ . However, we are done as if  $x = 1$  we get  $y + 1 = a + 1 \implies y = a$  and  $xy = b \implies b = y = a$ . This follows similarly for  $y = 1$ , and so  $f$  is injective.  $\square$

From the injectivity, and our earlier calculations we get  $f(1 - f(x)) = f(x) \implies f(x) = 1 - x$  and since  $-f$  must be a solution too,  $f(x) = x - 1$  is another solution. Therefore, our solutions are:

$$\boxed{f(x) = 0, 1 - x, x - 1}$$

□

**Problem 1.5 (USAMO 2018).** Find all functions  $f : (0, \infty) \rightarrow (0, \infty)$  such that

$$f\left(x + \frac{1}{y}\right) + f\left(y + \frac{1}{z}\right) + f\left(z + \frac{1}{x}\right) = 1$$

for all  $x, y, z > 0$  with  $xyz = 1$ .

*Proof.* We begin by substituting the values  $x = \frac{a}{b}$ ,  $y = \frac{b}{c}$ ,  $z = \frac{c}{a}$  to get the following:

$$\sum_{\text{cyc}} f\left(\frac{a+c}{b}\right) = 1$$

We then provide another substitution, this time for the function. Define a function  $g : (0, 1) \rightarrow (0, 1)$  such that  $g(x) = f(\frac{1-x}{x})$ , then for  $a + b + c = 1$  it follows that

$$g(a) + g(b) + g(c) = 1$$

**Claim 1.6 —**  $g$  satisfies Jensen's functional equation over  $(0, \frac{1}{2})$ .

*Proof.* Define variables  $x, y \in (0, \frac{1}{2})$  such that  $\frac{x+y}{2} \in (0, \frac{1}{2}) \subset (0, 1)$  and also  $0 < x + y < 1$ . Therefore, we can make the following substitutions:

$$a = b = \frac{x + y}{2}$$

$$c = 1 - (x + y)$$

Which satisfy  $a + b + c = 1$ , then,

$$g\left(\frac{x+y}{2}\right) + g\left(\frac{x+y}{2}\right) + g(1 - (x+y)) = 1$$

and

$$g(x) + g(y) + g(1 - (x+y)) = 1$$

Give us that

$$g(x) + g(y) = 2g\left(\frac{x+y}{2}\right),$$

as desired. □

We now define a new function  $h : [0, 1] \rightarrow \mathbb{R}$  as:

$$h(t) = g\left(\frac{2t+1}{8}\right) - (1-t)g\left(\frac{1}{8}\right) - tg\left(\frac{3}{8}\right)$$

Notice that  $h(0) = h(1) = 0$ . Also,

$$h\left(\frac{1}{2}\right) = g\left(\frac{2}{8}\right) - \frac{1}{2}g\left(\frac{1}{8}\right) - \frac{1}{2}g\left(\frac{3}{8}\right)$$

$$\begin{aligned} &= g\left(\frac{1}{8}\right) + g\left(\frac{1}{8}\right) - \frac{1}{2}g\left(\frac{1}{8}\right) - \frac{1}{2}g\left(\frac{1}{8}\right) - \frac{1}{2}g\left(\frac{1}{8}\right) \\ &= 0 \end{aligned}$$

So now we have  $h(0) = h(1) = h(\frac{1}{2}) = 0$ .

□

## §2 Problems

**Problem 2.1 (IMO 2008).** Find all functions  $f$  from the positive reals to the positive reals such that

$$\frac{f(w)^2 + f(x)^2}{f(y^2) + f(z^2)} = \frac{w^2 + x^2}{y^2 + z^2}$$

for all positive real numbers  $w, x, y, z$  satisfying  $wx = yz$ .

*Proof.* We let  $P(w, x, y, z)$  represent the given assertion. We claim that  $f(x) = x$  and  $f(x) = \frac{1}{x}$  are the only solutions, it is easy to check that these work.

First, note

$$P(x, x, x, x) \implies \frac{f(x)^2}{f(x^2)} = 1 \implies f(x^2) = f(x)^2$$

Which also gives  $f(1) = 1$ . Further,

$$P(x, 1, \sqrt{x}, \sqrt{x}) \implies (f(x) - x)(xf(x) - 1) = 0$$

Which gives us that  $f(x) = x$  or  $\frac{1}{x}$ . However, we now need to prove that  $f$  is not a piecewise function.

We assume  $f(a) = a$  and  $f(b) = \frac{1}{b}$  for some distinct  $a, b$ . We note:

$$P(\sqrt{a}, \sqrt{b}, \sqrt{ab}, 1) \implies \frac{a + \frac{1}{b}}{f(ab) + 1} = \frac{a + b}{ab + 1}$$

If we take  $f(ab) = ab$ , we get  $b = 1$  and the case where  $f(ab) = \frac{1}{ab}$  gives  $a = 1$ , contradicting that  $a$  and  $b$  are distinct. Thus,  $f$  is not piecewise, and the only solutions are  $f(x) = x$  and  $f(x) = \frac{1}{x}$  for all  $x \in \mathbb{R}^+$ .  $\square$