

A First Lean Formalization Audit of the IUT III Corollary 3.12 Corridor

IUT Lean formalization project

May 29, 2026

Abstract

This note records the present state of a Lean formalization of a narrow part of Mochizuki’s inter-universal Teichmüller theory: the passage from IUT III, Theorem 3.11 to Corollary 3.12, with attention to the Scholze–Stix objection. The current Lean code proves conditional route theorems, obstruction lemmas, and the final ordered-real algebra in Step (xi-f). It does not prove Corollary 3.12 and does not refute it. The main formal result so far is that a supplied hull/log-volume q -to- Θ comparison gives $C_\Theta \geq -1$, and that a nonarchimedean (Ind3) entry together with explicit Hodge/SHE square-weight transport data feeds such a comparison, while several simplified collapse mechanisms are rejected by Lean.

1 Scope

The review used the local Markdown conversions of IUT I–IV, Mochizuki’s 2026 formalization note, and the Scholze–Stix note. The relevant source passages are:

- IUT I: initial Θ -data, prime strips, Hodge theaters, and non-scheme-theoretic Θ -links.
- IUT II: Hodge–Arakelov evaluation, theta values of shape q^{j^2} , multiradiality, and Gaussian monoids.
- IUT III: Theorem 3.11; Corollary 3.12, especially Steps (x) and (xi); Remarks 3.12.2 and 3.12.3.
- IUT IV: later explicit log-volume estimates and Diophantine consequences.
- Scholze–Stix: the ordered real-line identification problem in their Section 2.2.
- Mochizuki 2026: the “3.11.5 $\Rightarrow$ 3.12” fourth-triangle decomposition.

2 Mathematical Target

IUT III Step (x) treats procession-normalized log-volumes as invariants for (Ind1) and (Ind2), and turns (Ind3) into an upper inequality. Step (xi) then uses the multiradial construction, IPL, SHE, holomorphic hulls, determinants, and log-volume to place the q -pilot log-volume in an upper ray controlled by the Θ -pilot log-volume:

$$-|\log(q)| \in \mathbb{R}_{\leq -|\log(\Theta)|} \quad \Rightarrow \quad -|\log(q)| \leq -|\log(\Theta)|.$$

If C_Θ is any real number satisfying

$$-|\log(\Theta)| \leq C_\Theta |\log(q)|, \quad |\log(q)| > 0,$$

then Step (xi-f) concludes $C_\Theta \geq -1$. This last inference is now a Lean theorem. Mochizuki’s 2026 note isolates the final comparison as:

$$\text{Theorem 3.11} \xrightarrow{\text{APT+IPL}} \text{“3.11.5”} \xrightarrow{\text{SHE+IPL}} \text{Corollary 3.12.}$$

The present formalization targets only this corridor, not the full construction of Theorem 3.11.

3 Dispute Interface

Scholze–Stix require explicit bookkeeping for all ordered one-dimensional real vector spaces. Their pressure point is that the Θ -side concrete objects carry j^2 -scaled degrees, while a consistent identification of the real lines appears to leave no room for inserting all these scalars.

The formalization therefore separates:

representative j^2 data		raw real equality
balanced sign-compatible data		transported equality
aggregate averaged data	and	matched transport scales
final hull/log-volume data		cancellable ordered-line comparison.

This separation is the main correctness guard in the current code.

4 Lean Coverage

The relevant implementation is in

`Iut/Stage1/IUTStage1Source.lean` and `Iut/Stage1/IUTStage1Experiments.lean`.

The code currently covers:

- labeled real-line copies and positive-scale transports;
- a finite $\mathbb{F}_\ell$ -label model with representative square weights $j \mapsto j^2$;
- a representative pilot-degree profile $\deg(\Theta_j) = j^2 \deg(q)$;
- proof that one label-independent scale cannot absorb all representative j^2 scales in this model;
- proof that, for nonzero $\deg(q)$, one label-independent scale cannot account for all representative Θ_j -degrees;
- proof that the raw representative j^2 theta-degree rule does not descend to the sign quotient $|F_\ell| = F_\ell / \{\pm 1\}$ when $\deg(q) \neq 0$;
- an absolute-label exponent on $|F_\ell| = \{0\} \cup F_\ell^\times / \{\pm 1\}$ which agrees with j^2 for the canonical representatives $0 \leq j \leq (\ell - 1)/2$, and the corresponding theta-degree profile;
- the degree-level Gaussian evaluation $q \mapsto (q^{j^2})_{j \in |F_\ell|}$, including identity at $j = 1$;

- a canonical representative square-weight profile on F_ℓ : Lean proves the total j^2 -mass is positive from the $j = 1$ contribution and therefore constructs the profile without an external positivity premise; it also proves $6 \sum_{j \in \mathbb{Z}/\ell\mathbb{Z}} j_{\text{rep}}^2 = (\ell - 1)\ell(2(\ell - 1) + 1)$;
- the square-weighted constant-average check: a constant procession-normalized log-volume remains equal to the same constant after averaging with the canonical j^2 -weights;
- the same constant-average check inside the $F_\ell = \mathbb{Z}/\ell\mathbb{Z}$ cusp-label corridor: if the full-label log-volume is constant, the square-weighted average used in the Θ -route is that constant, including for the canonical weights;
- the constant-target structured-SHE subcase: the formal route now proves equality of the square-weighted average with the Θ -source average, not merely an upper bound;
- a log-volume shadow of the IUT III splitting-monoid action: the generator attached to a procession label contributes $j^2 \deg(q)$, contributes 0 at the core label, and adds its finite generator sum to the tensor-packet log-volume;
- the finite procession containers $S_{j+1}^\pm = \{0, \dots, j\}$, their nested inclusions, core-label stability, stage cardinality $j + 1$, and total procession ambiguity $(\ell^\pm)!$ rather than $(\ell^\pm)^{\ell^\pm}$;
- the map from the final procession container to $|F_\ell|$, sending the core label to 0 and recovering the exponent j^2 ;
- the log-volume shadow of the tensor packet over $S_{j+1}^\pm$, with each factor a finite-fiber direct sum over places above $v_\mathbb{Q}$, total log-volume given by the finite sum over the container, normalized denominator $j + 1$, and invariance under container permutations;
- the F_ℓ -label average specialization: for labels represented as $\mathbb{Z}/\ell\mathbb{Z}$, Lean rewrites the procession-normalized average denominator as ℓ ;
- the IUT III Step (x) averaged-log-volume corridor: Lean records that (Ind1) and (Ind2) preserve the procession-normalized averaged log-volume, while (Ind3) supplies an upper bound;
- base-valuation products of tensor packets, the finite $(F_{\text{mod}}^\times)_j$ -action shadow, and a coric transfer shadow $F \rightarrow D \rightarrow D^\Gamma$ preserving product log-volume without identifying Hodge-theater histories;
- the Step (xi-d)–(xi-g) shadow: determinant normalization, upper-ray membership, and equality of the two q -pilot log-volume computations;
- the Corollary 3.12 finiteness endpoint: the statement-level value $-|\log(\Theta)| \in \mathbb{R} \cup \{+\infty\}$ is represented by an explicit finite/ $+\infty$ split, and the hull/determinant upper-ray datum forces the finite real branch;
- the Corollary 3.12 statement boundary: the Θ -pilot log-volume is typed as subject to (Ind1), (Ind2), (Ind3), while the q -pilot log-volume is typed as not subject to these indeterminacies;
- the extended statement endpoint: from the finite Θ -branch, $0 < -|\log(q)|$, the upper-ray comparison, and the displayed hypothesis $-|\log(\Theta)| \leq C_\Theta |\log(q)|$, Lean proves $C_\Theta \geq -1$;

- the Step (xi-f) final algebra: from $-|\log(q)| \leq -|\log(\Theta)|$, $|\log(q)| > 0$, and $-|\log(\Theta)| \leq C_\Theta |\log(q)|$, Lean proves $C_\Theta \geq -1$;
- the signed-payload bridge: the existing Stage 1 comparison record $q_{\text{signed}} \leq \Theta_{\text{signed}}$, together with $\Theta_{\text{signed}} \leq C_\Theta(-q_{\text{signed}})$, feeds this Step (xi-f) algebra directly;
- the Step (xi-g) endpoint bridge: the two-computation q-pilot record produces the signed comparison payload with q_{signed} read from the input prime-strip log-volume and Θ_{signed} read from the hull/determinant upper-ray boundary;
- the opening sign reduction in the proof of Corollary 3.12: from $0 < -q_{\text{signed}}$, Lean proves that $0 \leq \Theta_{\text{signed}}$ immediately implies $q_{\text{signed}} \leq \Theta_{\text{signed}}$, matching the printed reduction to the case $-|\log(\Theta)| < 0$;
- the Step (xi-h) warning: for $N \geq 2$ and $\Theta < 0$, the tensor-power boundary satisfies $N\Theta < \Theta$;
- the Step (xii) shadow: local Frobenioid p_v^N -ambiguity changes log-volume by an integer shift, while global calibration fixes exponent 0;
- the Remark 3.12.1(iii) arithmetic fact $\mathbb{Q}_{>0} \cap \mathbb{Z}^\times = \{1\}$;
- the Remark 3.12.1(ii) generalized-link numerical bound $C_\Theta \geq -\lambda$ for $\lambda \in \mathbb{Q}_{>0}$, with strict strengthening over the $\lambda = 1$ bound when $\lambda < 1$;
- the q^λ -pilot endpoint algebra: from a signed comparison for $-\lambda|\log(q)|$ and the same $C_\Theta|\log(q)|$ upper bound, Lean derives $C_\Theta \geq -\lambda$, and hence $C_\Theta \geq -1$ when $\lambda \leq 1$;
- the Remark 3.12.2 distinct-label implication $q\text{-itw} \Rightarrow q\text{-itw} \wedge \Theta\text{-itw}/\text{indets}$, with q-native and Θ -native labels proved distinct;
- the Remark 3.12.2 toy calculation: forgotten concrete assignments $-h$ and $-2h$ are incompatible for $h > 0$, while $-h \in \mathbb{R}_{\leq -2h+\epsilon}$ implies $h \leq \epsilon$;
- the Fig. 3.9 column table: both $\bullet_0$ and $\circ$ roles are required in both columns, while q-pilot multiradiality is absent;
- the Remark 3.12.2(v) absorption distinction: zero-column unit indeterminacy is absorbed by a hull upper ray, while one-column logarithmic conjugacy is absorbed as log-volume equality;
- the Remark 3.12.3(i) metric analogy: $\text{vol}(S) > 0$ and $\chi_S = -\text{vol}(S)$ imply $\chi_S < 0$;
- the Remark 3.12.4(iii) factor- p analogy: theta-label normalization divides a gap log-volume by ℓ , and multiplication by ℓ recovers it;
- the Remark 3.12.4(v) p-adic analogue: an inclusion, not isomorphism, gives $(1-p)(2g-2) \leq 0$;
- the IUT IV, Theorem 1.10 algebraic handoff: an explicit expression for C_Θ , together with $C_\Theta \geq -1$ and the remaining elementary estimates, implies the displayed $\frac{1}{6}\log(q)$ bound;
- the final elementary coefficient estimates in IUT IV, Theorem 1.10: $(1 - 12/\ell^2)^{-1} \leq 2$ and $(1 - 12/\ell^2)^{-1}\frac{1}{4}(1 + 36d_{\text{mod}}/\ell) \leq 1 + 80d_{\text{mod}}/\ell$;
- the IUT IV, Theorem 1.10 base-change monotonicity: $\log(d_{F_{\text{tpd}}}) + \log(f_{F_{\text{tpd}}}) \leq \log(d_F) + \log(f_F)$ propagates through the final right-hand side;

- the IUT IV, Theorem 1.10 Step (i) identities: $n^{-1} \sum_{m=1}^n m = (n+1)/2$ and $n^{-1} \sum_{m=1}^n m^2 = (2n+1)(n+1)/6$;
- the IUT IV, Theorem 1.10 Step (ii) small-prime error: $\log(2^{11}3^35^2) \leq 21$, represented by the corresponding linear logarithmic bound;
- the IUT IV, Theorem 1.10 Step (i) GL_2 arithmetic constants for $p = 2, 3, 5$, including the product with the quadratic factor from (E5);
- the IUT IV, Theorem 1.10 Step (iii) algebraic estimate combining the Step (ii) K -bound and the displayed $\log(s_Q)$ -bound into $(1 + 4/\ell) \log(d_K, f_K) + (16/\ell) \log(s_Q) \leq (1 + 36d_{\text{mod}}/\ell)S + 2\log(\ell) + 70$;
- the IUT IV, Theorem 1.10 Step (viii) Proposition 1.6 case split: the two cases $e_{\text{mod}}^* \ell \geq \eta_{\text{prm}}$ and $e_{\text{mod}}^* \ell < \eta_{\text{prm}}$ imply the uniform $4/3(e_{\text{mod}}^* \ell + \eta_{\text{prm}})$ bound;
- the IUT IV, Theorem 1.10 Step (viii) final absorption of $2\log(\ell) + 74$ and the prime-product term into $10(e_{\text{mod}}^* \ell + \eta_{\text{prm}})$;
- the IUT IV, Theorem 1.10 final display: the Corollary 3.12 C_Θ handoff, followed by the tripodal-to- F monotonicity, gives the final F -level $\frac{1}{6} \log(q)$ inequality;
- the IUT IV, Theorem A bounded-discrepancy conclusion: boundedness below of $(1 + \epsilon)(\log \text{diff}_X + \log \text{cond}_D) - \text{ht}_{\omega_X(D)}$ implies the corresponding pointwise height inequality on points of bounded degree;
- the IUT IV, Corollary 2.2(i) bounded-discrepancy chain: $\frac{1}{6} \log(q_2) \approx \frac{1}{6} \log(q_V) \approx \frac{1}{6} \text{ht}_\infty \approx \text{ht}_{\omega_X(D)}$ composes to a direct bounded discrepancy relation;
- bounded-discrepancy transfer lemmas: pointwise upper and lower bounds move across an $\approx$ relation with the expected additive constant shift;
- the IUT IV, Corollary 2.2(ii) condition (C1) prime-scale window: $(\log(q_V))^{1/2} \leq \ell \leq 10\delta(\log(q_V))^{1/2} \log(2\delta \log(q_V))$. Lean keeps the square-root and logarithmic factor explicit, proves $\log(q_V) \geq 0$, and derives $1 \leq 10\delta \log(2\delta \log(q_V))$ from the nonempty window;
- the IUT IV, Corollary 2.2(ii) use of (C1) in Theorem 1.10: $\sqrt{h} \leq \ell$ and $80d_{\text{mod}} \leq \delta$ imply $80d_{\text{mod}}/\ell \leq \delta h^{-1/2}$;
- the IUT IV, Corollary 2.2(ii) error-term use of (C1): $d_{\text{mod}}^* \leq \delta$ and $\ell \leq 10\delta h^{1/2} \log(2\delta h)$ imply $20(d_{\text{mod}}^* \ell + \eta_{\text{prm}}) \leq 200\delta^2 h^{1/2} \log(2\delta h) + 20\eta_{\text{prm}}$;
- the IUT IV, Theorem 1.10 to Corollary 2.2(ii) first-bound composition: these two C1 replacements convert the Theorem 1.10 right-hand side into $(1 + \delta h^{-1/2})S + 200\delta^2 h^{1/2} \log(2\delta h) + 20\eta_{\text{prm}}$;
- the IUT IV, Corollary 2.2(ii) q_2/q comparison: the displayed estimate $\frac{1}{6} \log(q_2) - \frac{1}{6} \log(q) \leq \frac{1}{3} h^{1/2} \log(2\delta h)$ follows through the intermediate term $h^{1/2} \log(\ell)$;
- the IUT IV, Corollary 2.2(ii) pre- ϵ_E bound for $h = \log(q_V)$: the Theorem 1.10 bound, the q_2/q estimate, and the bounded discrepancy B_K imply

$$\frac{1}{6} h \leq (1 + \delta h^{-1/2})S + (15\delta)^2 h^{1/2} \log(2\delta h) + \frac{1}{2} C_K,$$

with $C_K = 40\eta_{\text{prm}} + 2B_K$;

- the IUT IV, Corollary 2.2(ii) condition (C2) inequality chain: $\frac{1}{6} \log(q) \leq \frac{1}{6} \log(q_2) \leq \frac{1}{6} \log(q_V) \leq (1 + \epsilon_E)(\log \text{diff}_X + \log \text{cond}_D) + C_K$;
- the IUT IV, Corollary 2.2(ii) definition of ϵ_E : $\epsilon_E = (60\delta)^2 h^{-1/2} \log(2\delta h)$, represented with $h^{1/2}$ and $\log(2\delta h)$ as explicit real parameters; Lean proves positivity and $\epsilon_E \geq 5\delta h^{-1/2}$ from $\delta \geq 2$, $h^{1/2} > 0$, and $\log(2\delta h) \geq 1$;
- the IUT IV, Corollary 2.2(ii) ϵ_E -error conversion: substituting the definition of ϵ_E , Lean proves $(15\delta)^2 h^{1/2} \log(2\delta h) \leq \frac{1}{6} h \cdot \frac{2}{5} \epsilon_E$;
- the IUT IV, Corollary 2.2(ii) bridge to the denominator step: the pre- ϵ_E h -bound, the error conversion, and $\delta h^{-1/2} \leq \epsilon_E/5$ yield the preliminary inequality used to move the $\frac{2}{5} \epsilon_E \frac{1}{6} h$ term to the left;
- the IUT IV, Corollary 2.2(ii) denominator step: Lean moves the term $(2\epsilon_E/5) \frac{1}{6} h$ to the left and obtains the denominator $(1 - 2\epsilon_E/5)^{-1}$;
- the IUT IV, Corollary 2.2(ii) final ϵ_E -absorption: from $0 < \epsilon_E \leq 1$, Lean proves

$$\frac{1 + \epsilon_E/5}{1 - 2\epsilon_E/5} \leq 1 + \epsilon_E, \quad (1 - 2\epsilon_E/5)^{-1} C_K/2 \leq C_K,$$

and hence the displayed C2 bound from the preceding denominator expression;

- the IUT IV, Corollary 2.2(ii) final h -bound: Lean composes the pre- ϵ_E bound, error conversion, move-left step, and absorption to prove $\frac{1}{6} h \leq (1 + \epsilon_E) S + C_K$;
- the IUT IV, Corollary 2.2(ii) final C2 bridge: using $S \leq \log \text{diff}_X(x_E) + \log \text{cond}_D(x_E)$, Lean derives the displayed C2 inequality chain;
- the IUT IV, Corollary 2.2(ii) comparison $\log \text{diff}_X(x_E) = \log(d_{F_{\text{tpd}}})$ and $\log(f_{F_{\text{tpd}}}) \leq \log \text{cond}_D(x_E) \leq \log(f_{F_{\text{tpd}}}) + \log(2\ell)$;
- the IUT IV, Corollary 2.2 to Theorem A passage: bounded-discrepancy equivalence $\frac{1}{6} \log(q_2) \approx \text{ht}_{\omega_X(D)}$, together with the C2 bound for $\frac{1}{6} \log(q_2)$, gives a lower bound for $(1 + \epsilon_E)(\log \text{diff}_X + \log \text{cond}_D) - \text{ht}_{\omega_X(D)}$;
- the IUT IV, Corollary 2.2/2.3 finite-exception passage: a lower bound on the complement of a finite exceptional set and a lower bound on the exceptional set glue to a global lower bound by taking the minimum of the two constants;
- the IUT IV, Corollary 2.3 Diophantine inequality: Lean records the GenEll transfer as an explicit hypothesis and derives the global bounded-discrepancy lower bound for $(1 + \epsilon)(\log \text{diff}_X + \log \text{cond}_D) - \text{ht}_{\omega_X(D)}$;
- comparison levels: pointwise representative, aggregate representative, balanced sign-compatible, and hull/log-volume;
- endpoint audits for the 3.11.5 $\Rightarrow$ 3.12 comparison chart;
- nonarchimedean and archimedean (Ind3) local orientations;
- ordered real-line alignment with cancellation only after scale equality;
- a canonical unit-scale ordered alignment from a nonarchimedean (Ind3) entry;
- a bridge from factored square/full-label SHE obligations to the Hodge-descent (Ind3) route.

5 Current Lean Results

The current first-pass experiment report evaluates to:

route theorem flags	= true,
collapse-obstruction flags	= true,
Gaussian/procession finite-model flags	= true,
balancedLevelRejectedAtFinalRouteTheoremAvailable	= true,
disputeSettledByCurrentStage	= false.

Here the route flags are ordered real-line, nonarchimedean entry, and factored SHE availability. The obstruction flags are raw-cancellation failure, label-independent j^2 failure, and sign-quotient representative descent failure. The Gaussian/procession flags include finite tensor-packet, monoid action, canonical positive square-weight profile, the constant-average check, base-valuation product, coric-transfer, determinant-normalization, upper-ray, q -pilot two-computation, the Step (x) averaged-log-volume indeterminacy corridor, tensor-power warning, and global Frobenioid-calibration shadows, plus the finiteness endpoint for $-|\log(\Theta)| \in \mathbb{R} \cup \{+\infty\}$, the direct Corollary 3.12 statement endpoint with Θ -subject/ q -not-subject pilot bookkeeping, the Step (xi-g)-to-(xi-f) endpoint bridge giving $C_\Theta \geq -1$ from the two-computation q -pilot record, and positive-rational unit rigidity. The generalized-link flag records the rational parameter inequality $C_\Theta \geq -\lambda$, now with the corresponding q^λ -pilot real-valued derivation. The distinct-label flag records the logical intertwining display in Remark 3.12.2 and rejects collapse of the q -native and Θ -native labels. The toy-model flag records the resulting upper-ray arithmetic $h \leq \epsilon$. The column-table flag records the Fig. 3.9 theta/ q similarity-difference split. The absorption flag records the zero-column upper-bound versus one-column equality distinction. The metric flag records the Gauss–Bonnet sign calculation used as analogy for the upper semi-compatibility inequality. The factor- p flag records the F_ℓ -label averaging denominator. The Frobenius-derivative flag records the degree inequality from inclusion. The IUT IV flag records the downstream Theorem 1.10 use of the Corollary 3.12 lower bound on C_Θ , including the final coefficient estimates and the tripodal-to- F log-degree monotonicity. The Step (i) flag records the two elementary sum identities used in the leading-term computation. The Step (ii) small-prime flag records the additive error bound 21. The GL_2 flag records the finite cardinality constants feeding that bound. The Step (iii) flag records the $\log(s_Q)$ coefficient absorption. The Step (viii) flag records the prime-product case split used before the final upper bound for C_Θ , plus the final coarse error absorption. The final display flag records the composed Theorem 1.10 inequality with F -level log-degree terms. The Theorem A flag records the bounded-discrepancy height/log-different/log-conductor inequality. The Corollary 2.2 flag records composition of the displayed bounded-discrepancy chain. The transfer flag records movement of pointwise bounds across bounded-discrepancy equivalence. The Corollary 2.2(C1) flag records the displayed prime-scale window and its elementary consequences. The Corollary 2.2(C2) flag records the displayed three-step logarithmic inequality chain. The C1 coefficient flag records the replacement of $80d_{\text{mod}}/\ell$ by $\delta h^{-1/2}$ in the Theorem 1.10-to-C2 proof. The C1 error-term flag records the corresponding replacement of $20d_{\text{mod}}^*\ell$ by $200\delta^2 h^{1/2} \log(2\delta h)$. The first-bound flag records the composition of those replacements with Theorem 1.10. The ϵ_E -definition flag records the source formula and lower estimate for ϵ_E . The ϵ_E -move-left flag records introduction of the denominator $(1 - 2\epsilon_E/5)^{-1}$. The ϵ_E -absorption flag records the final denominator removal in C2. The ϵ_E -error flag records the conversion of the $(15\delta)^2$ -term to the $\frac{2}{5}\epsilon_E \frac{1}{6}h$ term. The final h -bound flag records the full composition through absorption. The final C2 bridge flag records the last monotonic replacement of the tripodal log-degree sum by the curve different/conductor sum. The h -to-move-left flag records the bridge from the pre- ϵ_E bound to the denominator-stage preliminary inequality. q^2/q flag records the displayed comparison

of $\log(q_2)$ and $\log(q)$. The pre- ϵ_E h -bound flag records the subsequent absorption of the bounded q_V/q_2 discrepancy into C_K . The log-different/log-conductor flag records the bridge from tripodal field terms to curve functions. The Corollary 2.2-to-Theorem A flag records the formal bounded-discrepancy transfer from C2 to the Theorem A inequality. The finite-exception flags record the lower-bound gluing used when finite exceptional sets are enlarged in Corollary 2.2 and then passed toward Corollary 2.3. The Corollary 2.3 flag records the final GenEll-transfer interface for the Diophantine inequality. The theorem-level route is:

$$\begin{array}{ccccc}
\text{nonarch. (Ind3) entry} & \rightarrow & \text{canonical ordered } \mathbb{R}\text{-alignment} & \rightarrow & \text{source/target alignment} \\
& & \downarrow & & \downarrow \\
\text{factored SHE transport} & \rightarrow & \text{Hodge-descent audit} & \rightarrow & q_{\text{signed}} \leq \Theta_{\text{signed}}.
\end{array}$$

The route remains conditional on the displayed hypotheses.

6 Audit Findings

No formal statement currently proves IUT III, Corollary 3.12 from Mochizuki's published hypotheses. The strongest positive result is conditional: if the hull/log-volume upper-ray comparison supplies $-\log(q) \leq -\log(\Theta)$, then Lean checks the formal Step (xi-f) conclusion $C_\Theta \geq -1$. If the nonarchimedean (Ind3) entry, ordered real-line alignment, and Hodge/SHE square-weight transport audit are supplied, Lean checks the final signed q -to- Θ inequality feeding that algebra.

The strongest negative result is also limited: in the current $\mathbb{F}_\ell$ -label model, Lean rejects a single label-independent scale accounting for all representative j^2 factors, and it rejects balanced sign-compatible evidence as the final hull/log-volume comparison level. Lean also rejects direct descent of arbitrary raw representative j^2 theta-degree data to $|F_\ell|$, while accepting the half-range exponent convention stated in IUT II. Lean also checks that an N -th tensor-power boundary is a strict strengthening for $N \geq 2$ and negative pilot log-volume. It also checks the Step (xii) numerical warning: nonzero local Frobenioid shifts change log-volume, whereas global calibration restores the unshifted value. The positive rational unit rigidity fact in Remark 3.12.1(iii) is formalized directly over $\mathbb{Q}$. For generalized links, Lean records that $\lambda < 1$ makes $-\lambda > -1$. Lean also checks that the Remark 3.12.2 simultaneous-intertwining step depends on distinct labels rather than an unlabeled identification. In the accompanying toy model, Lean proves the final upper bound exactly from the upper-ray membership hypothesis. Lean also records that the q -pilot column lacks the theta-side multiradiality used in Theorem 3.11(iii)(c). For Remark 3.12.2(v), Lean separates hull absorption of unit shifts from log-volume equality after logarithmic-conjugate compatibility. For Remark 3.12.3(i), Lean proves the stated negative Euler-characteristic sign from positivity of metric volume. For Remark 3.12.4(iii), Lean checks the ℓ -denominator normalization identity for the theta-label averaging analogy. For Remark 3.12.4(v), Lean proves the displayed nonpositive degree defect from $p \geq 2$ and $g \geq 2$. For IUT IV, Theorem 1.10, Lean proves the algebraic implication from the computed C_Θ expression and $C_\Theta \geq -1$ to the final $\frac{1}{6}\log(q)$ estimate, conditional on the separate elementary estimates. Lean now also proves the two displayed final elementary coefficient estimates from $\ell \geq 5$. It also proves the final monotonic replacement of the tripodal log-degree sum by the larger F -level log-degree sum, conditional on the two component monotonicity inequalities cited from Proposition 1.3(i). Lean also proves the Step (i) sum and square-sum identities over $\mathbb{R}$. For Step (ii), Lean proves the small-prime additive error bound from the displayed logarithmic inequalities. Lean also checks the Step (i) GL_2 constants for $p = 2, 3, 5$ and their product with the quadratic extension factor. For Step (iii), Lean proves the displayed coefficient absorption from the source hypotheses $d_{\text{mod}} \geq 1$, $\ell \geq 5$, $2\log(\ell) \leq \ell$, and nonnegativity of the tripodal log-degree

sum. For Step (viii), Lean checks that the two Proposition 1.6 cases imply the uniform prime-product bound used in the final C_Θ estimate. Lean also checks the subsequent absorption into the coefficient 10 of $e_{\text{mod}}^* \ell + \eta_{\text{prm}}$. Combining the Corollary 3.12 handoff with the tripodal-to- F monotonicity, Lean proves the final displayed F -level inequality of Theorem 1.10 at this real-valued shadow level. For Theorem A, Lean records the exact bounded-discrepancy conclusion and proves the equivalent pointwise height bound from the lower-bound hypothesis. For Corollary 2.2(i), Lean defines equality up to bounded discrepancy, proves reflexivity, symmetry, transitivity, nonnegative scaling, and composes the displayed chain from $\log(q_2)$ to $\text{ht}_{\omega_X(D)}$. Lean also proves that upper and lower pointwise bounds transfer across such a bounded-discrepancy equivalence with the appropriate constant shift. For Corollary 2.2(ii)(C1), Lean records the displayed window for ℓ and derives nonnegativity of $\log(q_V)$, positivity of $(\log(q_V))^{1/2}$, and $1 \leq 10\delta \log(2\delta \log(q_V))$. Lean also verifies the C1 coefficient replacement $80d_{\text{mod}}/\ell \leq \delta h^{-1/2}$ used immediately after applying Theorem 1.10. Lean also verifies the C1 upper-window replacement for the Theorem 1.10 error term $20(d_{\text{mod}}^* \ell + \eta_{\text{prm}})$. Combining these, Lean proves the first displayed post-Theorem-1.10 bound in the proof of Corollary 2.2(ii). Lean also proves the displayed comparison $\frac{1}{6} \log(q_2) - \frac{1}{6} \log(q) \leq \frac{1}{3} h^{1/2} \log(2\delta h)$ from the intermediate $h^{1/2} \log(\ell)$ bounds. Lean then combines this with the bounded q_V/q_2 discrepancy to prove the displayed pre- ϵ_E bound for $\frac{1}{6}h$. For Corollary 2.2(ii), Lean proves the final $\frac{1}{6} \log(q)$ bound from the displayed $\log(q)$, $\log(q_2)$, and $\log(q_V)$ chain. Lean records the proof's definition of ϵ_E and verifies the stated lower estimate $\epsilon_E \geq 5\delta h^{-1/2}$. Lean also verifies the source conversion $(15\delta)^2 h^{1/2} \log(2\delta h) \leq \frac{1}{6}h \cdot \frac{2}{5}\epsilon_E$. Combining this with $\delta h^{-1/2} \leq \epsilon_E/5$, Lean proves the preliminary inequality needed for the move-left step. Lean then verifies the algebraic move-left step that introduces the denominator $(1 - 2\epsilon_E/5)^{-1}$. Lean also proves the final ϵ_E -absorption step used just before C2: the intermediate expression with denominator $1 - 2\epsilon_E/5$ is bounded by $(1 + \epsilon_E)$ times the log-degree sum plus C_K . Lean now composes these steps into the final displayed $h = \log(q_V)$ bound. Lean also derives the displayed C2 chain from this h -bound and the tripodal-to-curve log-degree comparison. It also proves the two sum inequalities relating the tripodal different and conductor terms to $\log \text{diff}_X + \log \text{cond}_D$. For the Corollary 2.2/2.3 exceptional-set passage, Lean proves that a lower bound off a finite exceptional set and a lower bound on the exceptional set imply a global lower bound, and specializes this to the Theorem A discrepancy. For Corollary 2.3, Lean proves the stated bounded-discrepancy conclusion from an explicit GenEll transfer hypothesis; GenEll itself is not formalized here. Finally, Lean combines the Corollary 2.2(i) bounded-discrepancy height comparison with the C2 bound to construct the Theorem A lower-bound shadow.

7 Missing Mathematics

The present code does not yet formalize:

- initial Θ -data as in IUT I;
- actual Hodge theaters, Frobenioids, prime strips, log-shells, or Θ -links;
- the Hodge–Arakelov theta evaluation and Gaussian monoids of IUT II;
- the construction of the actual theta values, Gaussian/splitting monoids, $F_{\text{mod}}^\times$ -actions, Kummer isomorphisms, mono-analyticization, and Frobenioids behind the present shadows;
- the actual construction of log-shells and tensor packets attached to mono-analytic prime-strips in IUT III, Theorem 3.11;

- the genuine holomorphic hull and determinant operations of Step (xi), beyond the present finite determinant/upper-ray/two-computation/tensor-power shadows;
- the global realified Frobenioids of Step (xii), beyond the present integer log-volume shift model;
- the arithmetic-divisor and local log-volume construction behind the IUT IV Theorem 1.10 numerical estimates, and the Diophantine applications;
- the construction of the normalized height, log-different, and log-conductor functions used in IUT IV, Theorem A;
- the construction of the bounded-discrepancy equivalences asserted in IUT IV, Corollary 2.2(i);
- a proof that the current Hodge/SHE transport obligations follow from Mochizuki’s full hypotheses.

8 Next Work

The next significant formal step is not another wrapper. It is to replace current explicit Hodge/SHE transport assumptions by constructions closer to the papers:

$$\begin{aligned}
 & \text{theta values } q^{j^2} \\
 \rightsquigarrow & \text{ Gaussian and splitting monoids} \\
 \rightsquigarrow & \text{ square/full-label preservation} \\
 \rightsquigarrow & \text{ hull/log-volume route.}
 \end{aligned}$$

Only after this bridge is formalized can the first-pass experiment begin to test the actual mathematical load-bearing part of Mochizuki’s response to Scholze–Stix.

References

- [1] S. Mochizuki, *Inter-universal Teichmüller Theory I: Construction of Hodge Theaters*.
- [2] S. Mochizuki, *Inter-universal Teichmüller Theory II: Hodge-Arakelov-theoretic Evaluation*.
- [3] S. Mochizuki, *Inter-universal Teichmüller Theory III: Canonical Splittings of the Log-theta-lattice*.
- [4] S. Mochizuki, *Inter-universal Teichmüller Theory IV: Log-volume Computations and Set-theoretic Foundations*.
- [5] S. Mochizuki, *On the Formalization of IUT: Preliminary Progress Report*, 2026.
- [6] P. Scholze and J. Stix, *Why abc is still a conjecture*.