

CENG 216 – NUMERICAL COMPUTATION

LEAST SQUARES - PART I

Mustafa Özuysal

`mustafaozuysal@iyte.edu.tr`

April 3, 2022

İzmir Institute of Technology

Slides are based on material from the main textbook:

“Numerical Analysis”, The new international edition, 2ed,
by Timothy Sauer

LEAST SQUARES AND THE NORMAL EQUATIONS

INCONSISTENT SYSTEMS OF EQUATIONS

$A\mathbf{x} = \mathbf{b}$ with no solutions

$$x_1 + x_2 = 2$$

$$x_1 - x_2 = 1$$

$$x_1 + x_2 = 3$$

INCONSISTENT SYSTEMS OF EQUATIONS

$A\mathbf{x} = \mathbf{b}$ with no solutions \rightarrow Equations are inconsistent.

$$x_1 + x_2 = 2$$

$$x_1 - x_2 = 1$$

$$x_1 + x_2 = 3$$

INCONSISTENT SYSTEMS OF EQUATIONS

$A\mathbf{x} = \mathbf{b}$ with no solutions \rightarrow Equations are inconsistent.

$$\left. \begin{array}{l} x_1 + x_2 = 2 \\ x_1 - x_2 = 1 \\ x_1 + x_2 = 3 \end{array} \right\} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

INCONSISTENT SYSTEMS OF EQUATIONS

$A\mathbf{x} = \mathbf{b}$ with no solutions \rightarrow Equations are inconsistent.

$$\left. \begin{array}{l} x_1 + x_2 = 2 \\ x_1 - x_2 = 1 \\ x_1 + x_2 = 3 \end{array} \right\} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

$$x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix},$$

and there is no x_1, x_2 pair that satisfies this equality.

INCONSISTENT SYSTEMS OF EQUATIONS: GENERAL CASE

In general,

$$\begin{aligned}\mathbf{A}_{m \times n} \mathbf{x} &= \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= x_1 \mathbf{v}_1 + x_2 \mathbf{v}_2 + \dots + x_n \mathbf{v}_n \\ &= \mathbf{b},\end{aligned}$$

where $\mathbf{v}_i \in \mathbb{R}^m$ and $\mathbf{b} \in \mathbb{R}^m$.

INCONSISTENT SYSTEMS OF EQUATIONS: GENERAL CASE

In general,

$$\begin{aligned}\mathbf{A}_{m \times n} \mathbf{x} &= \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= x_1 \mathbf{v}_1 + x_2 \mathbf{v}_2 + \dots + x_n \mathbf{v}_n \\ &= \mathbf{b},\end{aligned}$$


where $\mathbf{v}_i \in \mathbb{R}^m$ and $\mathbf{b} \in \mathbb{R}^m$. Whenever $m > n$, there is the possibility that the equations will be inconsistent.

INCONSISTENT SYSTEMS OF EQUATIONS: GEOMETRIC VIEW

$$\mathbf{Ax} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \mathbf{b}$$

INCONSISTENT SYSTEMS OF EQUATIONS: GEOMETRIC VIEW


$$\mathbf{Ax} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \mathbf{b}$$




Linear combination of vectors creates another vector in the same plane.

INCONSISTENT SYSTEMS OF EQUATIONS: GEOMETRIC VIEW

$$\mathbf{Ax} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \mathbf{b}$$




Linear combination of vectors creates another vector in the same plane.




Inconsistent System

INCONSISTENT SYSTEMS OF EQUATIONS: GEOMETRIC VIEW

$$\mathbf{Ax} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 = x_1 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \mathbf{b}$$



Linear combination of vectors creates another vector in the same plane.



Inconsistent System

REMINDER: VECTOR AND MATRIX TRANSPOSE

Def: **Transpose** of A is A^T with rows as the columns of A .

REMINDER: VECTOR AND MATRIX TRANSPOSE

Def: **Transpose** of A is A^T with rows as the columns of A .
If A is $m \times n$ then A^T is $n \times m$.

REMINDER: VECTOR AND MATRIX TRANSPOSE

Def: Transpose of A is A^T with rows as the columns of A .
If A is $m \times n$ then A^T is $n \times m$.

$$(A + B)^T = A^T + B^T \text{ and } (AB)^T = B^T A^T.$$

REMINDER: VECTOR AND MATRIX TRANSPOSE

Def: Transpose of \mathbf{A} is \mathbf{A}^\top with rows as the columns of \mathbf{A} .
If \mathbf{A} is $m \times n$ then \mathbf{A}^\top is $n \times m$.

$$(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top \text{ and } (\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top.$$

$$\mathbf{u}^\top \mathbf{v} = \mathbf{v}^\top \mathbf{u} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta_{\mathbf{uv}},$$

where $\theta_{\mathbf{uv}}$ is the angle between \mathbf{u} and \mathbf{v} .

REMINDER: VECTOR AND MATRIX TRANSPOSE

Def: Transpose of \mathbf{A} is \mathbf{A}^\top with rows as the columns of \mathbf{A} .
If \mathbf{A} is $m \times n$ then \mathbf{A}^\top is $n \times m$.

$$(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top \text{ and } (\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top.$$

$$\mathbf{u}^\top \mathbf{v} = \mathbf{v}^\top \mathbf{u} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta_{\mathbf{uv}},$$

where $\theta_{\mathbf{uv}}$ is the angle between \mathbf{u} and \mathbf{v} . If $\mathbf{u} \perp \mathbf{v}$ then
 $\theta_{\mathbf{uv}} = 90^\circ \implies \cos \theta_{\mathbf{uv}} = 0 \implies \mathbf{u}^\top \mathbf{v} = 0$.

REMINDER: VECTOR AND MATRIX TRANSPOSE


Def: Transpose of \mathbf{A} is \mathbf{A}^\top with rows as the columns of \mathbf{A} .
If \mathbf{A} is $m \times n$ then \mathbf{A}^\top is $n \times m$.

$$(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top \text{ and } (\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top.$$

$$\mathbf{u}^\top \mathbf{v} = \mathbf{v}^\top \mathbf{u} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta_{\mathbf{uv}},$$


where $\theta_{\mathbf{uv}}$ is the angle between \mathbf{u} and \mathbf{v} . If $\mathbf{u} \perp \mathbf{v}$ then
 $\theta_{\mathbf{uv}} = 90^\circ \implies \cos \theta_{\mathbf{uv}} = 0 \implies \mathbf{u}^\top \mathbf{v} = 0.$

NORMAL EQUATIONS




$$(\mathbf{b} - \mathbf{Ax}_{\text{LS}}) \perp \{\mathbf{Ax} \mid \mathbf{x} \in \mathbb{R}^n\}$$

NORMAL EQUATIONS




$$(\mathbf{b} - \mathbf{Ax}_{LS}) \perp \{\mathbf{Ax} \mid \mathbf{x} \in \mathbb{R}^n\} \implies (\mathbf{Ax})^\top (\mathbf{b} - \mathbf{Ax}_{LS}) = 0, \forall \mathbf{x} \in \mathbb{R}^n$$

NORMAL EQUATIONS




$$\begin{aligned} (b - Ax_{LS}) \perp \{Ax \mid x \in \mathbb{R}^n\} &\implies (Ax)^T (b - Ax_{LS}) = 0, \forall x \in \mathbb{R}^n \\ &\implies x^T A^T (b - Ax_{LS}) = 0, \forall x \in \mathbb{R}^n \end{aligned}$$

NORMAL EQUATIONS




$$\begin{aligned} (b - Ax_{LS}) \perp \{Ax \mid x \in \mathbb{R}^n\} &\implies (Ax)^T (b - Ax_{LS}) = 0, \forall x \in \mathbb{R}^n \\ &\implies x^T A^T (b - Ax_{LS}) = 0, \forall x \in \mathbb{R}^n \\ &\implies A^T (b - Ax_{LS}) = 0 \end{aligned}$$

NORMAL EQUATIONS



$$\begin{aligned}(\mathbf{b} - \mathbf{Ax}_{\text{LS}}) \perp \{\mathbf{Ax} \mid \mathbf{x} \in \mathbb{R}^n\} &\implies (\mathbf{Ax})^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{x}^\top \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = \mathbf{0} \\&\implies \mathbf{A}^\top \mathbf{Ax}_{\text{LS}} = \mathbf{A}^\top \mathbf{b} \quad \text{(Normal Equations)}\end{aligned}$$


NORMAL EQUATIONS



$$\begin{aligned}(\mathbf{b} - \mathbf{Ax}_{\text{LS}}) \perp \{\mathbf{Ax} \mid \mathbf{x} \in \mathbb{R}^n\} &\implies (\mathbf{Ax})^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{x}^\top \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{\text{LS}}) = \mathbf{0} \\&\implies \mathbf{A}^\top \mathbf{Ax}_{\text{LS}} = \mathbf{A}^\top \mathbf{b} \quad \text{(Normal Equations)}\end{aligned}$$

\mathbf{r} is called the **residual** vector.

NORMAL EQUATIONS



$$\begin{aligned}(\mathbf{b} - \mathbf{Ax}_{LS}) \perp \{\mathbf{Ax} \mid \mathbf{x} \in \mathbb{R}^n\} &\implies (\mathbf{Ax})^\top (\mathbf{b} - \mathbf{Ax}_{LS}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{x}^\top \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{LS}) = 0, \forall \mathbf{x} \in \mathbb{R}^n \\&\implies \mathbf{A}^\top (\mathbf{b} - \mathbf{Ax}_{LS}) = 0 \\&\implies \mathbf{A}^\top \mathbf{Ax}_{LS} = \mathbf{A}^\top \mathbf{b} \quad \text{(Normal Equations)}\end{aligned}$$

\mathbf{r} is called the **residual** vector. The **squared error (SE)** and the **root mean squared error (RMSE)** are defined to be

$$\text{SE} = \|\mathbf{r}\|^2 = r_1^2 + r_2^2 + \dots + r_n^2 \text{ and } \text{RMSE} = \sqrt{\frac{\text{SE}}{m}} = \frac{\|\mathbf{r}\|}{\sqrt{m}}$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\mathbf{A}^T \mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\mathbf{A}^T \mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

$$\mathbf{A}^T \mathbf{b} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\mathbf{A}^T \mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

$$\mathbf{A}^T \mathbf{b} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

Solving the system of normal equations $\mathbf{A}^T \mathbf{A} \mathbf{x}_{\text{LS}} = \mathbf{A}^T \mathbf{b}$

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \implies \mathbf{x}_{\text{LS}} = \begin{bmatrix} \frac{7}{4} \\ \frac{3}{4} \end{bmatrix}.$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \implies \mathbf{x}_{\text{LS}} = \begin{bmatrix} \frac{7}{4} \\ \frac{3}{4} \end{bmatrix}.$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \implies \mathbf{x}_{LS} = \begin{bmatrix} \frac{7}{4} \\ \frac{3}{4} \end{bmatrix}.$$

Then,

$$\mathbf{r} = \mathbf{b} - \mathbf{A}\mathbf{x}_{LS} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} - \begin{bmatrix} 2.5 \\ 1 \\ 2.5 \end{bmatrix} = \begin{bmatrix} -0.5 \\ 0.0 \\ 0.5 \end{bmatrix}$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \implies \mathbf{x}_{LS} = \begin{bmatrix} \frac{7}{4} \\ \frac{3}{4} \end{bmatrix}.$$

Then,

$$\mathbf{r} = \mathbf{b} - \mathbf{A}\mathbf{x}_{LS} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} - \begin{bmatrix} 2.5 \\ 1 \\ 2.5 \end{bmatrix} = \begin{bmatrix} -0.5 \\ 0.0 \\ 0.5 \end{bmatrix}$$

$$SE = (-0.5)^2 + (0)^2 + (0.5)^2 = 0.5$$

EXAMPLE

Find the least squares solution \mathbf{x}_{LS} for $\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$.

$$\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \implies \mathbf{x}_{LS} = \begin{bmatrix} \frac{7}{4} \\ \frac{3}{4} \end{bmatrix}.$$

Then,

$$\mathbf{r} = \mathbf{b} - \mathbf{A}\mathbf{x}_{LS} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} - \begin{bmatrix} 2.5 \\ 1 \\ 2.5 \end{bmatrix} = \begin{bmatrix} -0.5 \\ 0.0 \\ 0.5 \end{bmatrix}$$


$$SE = (-0.5)^2 + (0)^2 + (0.5)^2 = 0.5$$

$$RMSE = \sqrt{\frac{SE}{3}} \approx 0.408$$

MODEL FITTING


FITTING MODELS TO DATA

- Let $(t_1, y_1), (t_2, y_2), \dots, (t_n, y_n)$ be a set of data points.




FITTING MODELS TO DATA

- Let $(t_1, y_1), (t_2, y_2), \dots, (t_n, y_n)$ be a set of **data points**.
- Given a **parametric model**, such as the line $y = c_1 + c_2 t$ with parameters c_1 and c_2 , find the “best” model that “fits” to the data points.



FITTING MODELS TO DATA

- Let $(t_1, y_1), (t_2, y_2), \dots, (t_n, y_n)$ be a set of **data points**.
- Given a **parametric model**, such as the line $y = c_1 + c_2 t$ with parameters c_1 and c_2 , find the “best” model that “fits” to the data points.
- The best parameters minimize the **model error** $\sum_{i=1}^n e_i^2$.




EXAMPLE

Find the line $y = c_1 + c_2 t$ that best fits the three data points $(t, y) = (1, 2)$, $(-1, 1)$, and $(1, 3)$.

$$c_1 + c_2(1) = 2$$

$$c_1 + c_2(-1) = 1$$


$$c_1 + c_2(1) = 3$$



EXAMPLE

Find the line $y = c_1 + c_2 t$ that best fits the three data points $(t, y) = (1, 2)$, $(-1, 1)$, and $(1, 3)$.


$$\left. \begin{array}{l} c_1 + c_2(1) = 2 \\ c_1 + c_2(-1) = 1 \\ c_1 + c_2(1) = 3 \end{array} \right\} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$



EXAMPLE

Find the line $y = c_1 + c_2 t$ that best fits the three data points $(t, y) = (1, 2)$, $(-1, 1)$, and $(1, 3)$.

$$\left. \begin{array}{l} c_1 + c_2(1) = 2 \\ c_1 + c_2(-1) = 1 \\ c_1 + c_2(1) = 3 \end{array} \right\} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} \rightarrow \begin{array}{l} c_1 = \frac{7}{4} \\ c_2 = \frac{3}{4} \end{array}$$




EXAMPLE

Find the line $y = c_1 + c_2 t$ that best fits the three data points $(t, y) = (1, 2), (-1, 1)$, and $(1, 3)$.

$$\left. \begin{array}{l} c_1 + c_2(1) = 2 \\ c_1 + c_2(-1) = 1 \\ c_1 + c_2(1) = 3 \end{array} \right\} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} \rightarrow \begin{array}{l} c_1 = \frac{7}{4} \\ c_2 = \frac{3}{4} \end{array}$$

So the best fitting line is


$$y = \frac{7}{4} + \frac{3}{4}t$$



EXAMPLE

Find the **line** $y = c_1 + c_2t$ and the parabola $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1)$, $(0, 0)$, $(1, 0)$, and $(2, -2)$.


$$\begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix}$$



EXAMPLE

Find the **line** $y = c_1 + c_2t$ and the parabola $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1)$, $(0, 0)$, $(1, 0)$, and $(2, -2)$.

$$\begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \right\} \begin{bmatrix} 4 & 2 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \end{bmatrix}$$




EXAMPLE

Find the **line** $y = c_1 + c_2t$ and the parabola $y = c_1 + c_2t + c_3t^2$ that best fits the four data points

$(t, y) = (-1, 1), (0, 0), (1, 0),$ and $(2, -2)$.

$$\begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \quad \left\} \quad \begin{bmatrix} 4 & 2 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \end{bmatrix} \rightarrow \begin{matrix} c_1 = 0.2 \\ c_2 = -0.9 \end{matrix}$$



EXAMPLE

Find the **line** $y = c_1 + c_2t$ and the parabola $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1), (0, 0), (1, 0),$ and $(2, -2)$.


$$\left\{ \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \right\} \quad \begin{bmatrix} 4 & 2 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \end{bmatrix} \rightarrow \begin{matrix} c_1 = 0.2 \\ c_2 = -0.9 \end{matrix}$$

So the best fitting line is

$$y = 0.2 - 0.9t$$

$$SE = 0.7$$


$$RMSE = 0.418$$



EXAMPLE

Find the line $y = c_1 + c_2t$ and the **parabola** $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1)$, $(0, 0)$, $(1, 0)$, and $(2, -2)$.

$$\begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix}$$




EXAMPLE

Find the line $y = c_1 + c_2t$ and the **parabola** $y = c_1 + c_2t + c_3t^2$ that best fits the four data points

$(t, y) = (-1, 1), (0, 0), (1, 0),$ and $(2, -2)$.


$$\left\{ \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \right\} \quad \begin{bmatrix} 4 & 2 & 6 \\ 2 & 6 & 8 \\ 6 & 8 & 18 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \\ -7 \end{bmatrix}$$



EXAMPLE

Find the line $y = c_1 + c_2t$ and the **parabola** $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1)$, $(0, 0)$, $(1, 0)$, and $(2, -2)$.

$$\left\{ \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \right\} \quad \begin{bmatrix} 4 & 2 & 6 \\ 2 & 6 & 8 \\ 6 & 8 & 18 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \\ -7 \end{bmatrix} \rightarrow \begin{aligned} c_1 &= 0.45 \\ c_2 &= -0.65 \\ c_3 &= -0.25 \end{aligned}$$



EXAMPLE

Find the line $y = c_1 + c_2t$ and the **parabola** $y = c_1 + c_2t + c_3t^2$ that best fits the four data points $(t, y) = (-1, 1), (0, 0), (1, 0),$ and $(2, -2)$.


$$\left\{ \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -2 \end{bmatrix} \right\} \begin{bmatrix} 4 & 2 & 6 \\ 2 & 6 & 8 \\ 6 & 8 & 18 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} -1 \\ -5 \\ -7 \end{bmatrix} \rightarrow \begin{matrix} c_1 = 0.45 \\ c_2 = -0.65 \\ c_3 = -0.25 \end{matrix}$$

So the best fitting parabola is

$$y = 0.45 - 0.65t - 0.25t^2$$

$$SE = 0.45$$

$$RMSE \approx 0.335$$



CONDITIONING OF LEAST SQUARES

So far, to solve the least square problem $\mathbf{Ax} = \mathbf{b}$, we have solved the normal equations

$$\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}.$$

CONDITIONING OF LEAST SQUARES

So far, to solve the least square problem $\mathbf{Ax} = \mathbf{b}$, we have solved the normal equations

$$\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}.$$

However, the numerically this is problematic since

$$\text{cond}(\mathbf{A}^\top \mathbf{A}) = (\text{cond}(\mathbf{A}))^2,$$

so if the condition number of \mathbf{A} is large, the condition number of $\mathbf{A}^\top \mathbf{A}$ is even larger! We need a better approach that does not require explicitly forming $\mathbf{A}^\top \mathbf{A}$.

CONDITIONING OF LEAST SQUARES

So far, to solve the least square problem $\mathbf{Ax} = \mathbf{b}$, we have solved the normal equations

$$\mathbf{A}^\top \mathbf{Ax} = \mathbf{A}^\top \mathbf{b}.$$

However, the numerically this is problematic since

$$\text{cond}(\mathbf{A}^\top \mathbf{A}) = (\text{cond}(\mathbf{A}))^2,$$

so if the condition number of \mathbf{A} is large, the condition number of $\mathbf{A}^\top \mathbf{A}$ is even larger! We need a better approach that does not require explicitly forming $\mathbf{A}^\top \mathbf{A}$. → **QR Factorization**

QR FACTORIZATION

GRAM-SCHMIDT ORTHOGONALIZATION

- Given an input set of m -dimensional vectors, find an **orthogonal** coordinate system for the subspace spanned by the set (all vectors that can be reached by linear combination).

GRAM-SCHMIDT ORTHOGONALIZATION

- Given an input set of m -dimensional vectors, find an **orthogonal** coordinate system for the subspace spanned by the set (all vectors that can be reached by linear combination).
- **Input:** n -linearly independent vectors $\{\mathbf{a}_i\}$.

GRAM-SCHMIDT ORTHOGONALIZATION


- Given an input set of m -dimensional vectors, find an **orthogonal** coordinate system for the subspace spanned by the set (all vectors that can be reached by linear combination).
- **Input:** n -linearly independent vectors $\{\mathbf{a}_i\}$.
- **Output:** n mutually perpendicular unit vectors $\{\mathbf{q}_i\}$ that forms an orthogonal coordinate system for $\text{span}\{\mathbf{a}_i\}$

$$\|\mathbf{q}_i\| = 1 \implies \mathbf{q}_i^\top \mathbf{q}_i = 1, \quad \forall i = 1, \dots, n.$$

$$\mathbf{q}_i \perp \mathbf{q}_j \implies \mathbf{q}_i^\top \mathbf{q}_j = 0, \quad \forall i, j \in \{1, \dots, n\} \text{ and } i \neq j.$$

GRAM-SCHMIDT ORTHOGONALIZATION

Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n \in \mathbb{R}^m$ be linearly independent ($m \geq n$).




GRAM-SCHMIDT ORTHOGONALIZATION


Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n \in \mathbb{R}^m$ be linearly independent ($m \geq n$).

$$\mathbf{y}_1 = \mathbf{a}_1,$$

$$\mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|}$$



GRAM-SCHMIDT ORTHOGONALIZATION



Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n \in \mathbb{R}^m$ be linearly independent ($m \geq n$).

$$\mathbf{y}_1 = \mathbf{a}_1,$$

$$\mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^T \mathbf{a}_2)$$

GRAM-SCHMIDT ORTHOGONALIZATION


Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n \in \mathbb{R}^m$ be linearly independent ($m \geq n$).

$$\mathbf{y}_1 = \mathbf{a}_1,$$


$$\mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2),$$

$$\mathbf{q}_2 = \frac{\mathbf{y}_2}{\|\mathbf{y}_2\|}$$



GRAM-SCHMIDT ORTHOGONALIZATION



Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n \in \mathbb{R}^m$ be linearly independent ($m \geq n$).

$$\begin{aligned} \mathbf{y}_1 &= \mathbf{a}_1, & \mathbf{q}_1 &= \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} \\ \mathbf{y}_2 &= \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2), & \mathbf{q}_2 &= \frac{\mathbf{y}_2}{\|\mathbf{y}_2\|} \\ &\vdots & &\vdots \\ \mathbf{y}_j &= \mathbf{a}_j - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j) - \mathbf{q}_2(\mathbf{q}_2^\top \mathbf{a}_j) & & \\ &\quad - \dots - \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j), & \mathbf{q}_j &= \frac{\mathbf{y}_j}{\|\mathbf{y}_j\|} \end{aligned}$$

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)
- Assume $\mathbf{q}_i \perp \mathbf{q}_k$, $\forall k \leq j-1, k \neq i$,
Prove $\mathbf{q}_i \perp \mathbf{y}_j$:

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)
- Assume $\mathbf{q}_i \perp \mathbf{q}_k$, $\forall k \leq j-1, k \neq i$,
Prove $\mathbf{q}_i \perp \mathbf{y}_j$:

$$\mathbf{q}_i^\top \mathbf{y}_j = \mathbf{q}_i^\top (\mathbf{a}_j - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j) - \mathbf{q}_2(\mathbf{q}_2^\top \mathbf{a}_j) - \dots - \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j))$$

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)
- Assume $\mathbf{q}_i \perp \mathbf{q}_k$, $\forall k \leq j-1, k \neq i$,

Prove $\mathbf{q}_i \perp \mathbf{y}_j$:

$$\begin{aligned}\mathbf{q}_i^\top \mathbf{y}_j &= \mathbf{q}_i^\top (\mathbf{a}_j - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j) - \mathbf{q}_2(\mathbf{q}_2^\top \mathbf{a}_j) - \dots - \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)) \\ &= \mathbf{q}_i^\top \mathbf{a}_j - \underbrace{\mathbf{q}_i^\top \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j)}_0 - \dots - \mathbf{q}_i^\top \mathbf{q}_i(\mathbf{q}_i^\top \mathbf{a}_j) - \dots - \underbrace{\mathbf{q}_i^\top \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)}_0\end{aligned}$$

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)
- Assume $\mathbf{q}_i \perp \mathbf{q}_k$, $\forall k \leq j-1, k \neq i$,

Prove $\mathbf{q}_i \perp \mathbf{y}_j$:

$$\begin{aligned}\mathbf{q}_i^\top \mathbf{y}_j &= \mathbf{q}_i^\top (\mathbf{a}_j - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j) - \mathbf{q}_2(\mathbf{q}_2^\top \mathbf{a}_j) - \dots - \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)) \\ &= \mathbf{q}_i^\top \mathbf{a}_j - \underbrace{\mathbf{q}_i^\top \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j)}_0 - \dots - \mathbf{q}_i^\top \mathbf{q}_i(\mathbf{q}_i^\top \mathbf{a}_j) - \dots - \underbrace{\mathbf{q}_i^\top \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)}_0 \\ &= \mathbf{q}_i^\top \mathbf{a}_j - \underbrace{\mathbf{q}_i^\top \mathbf{q}_i(\mathbf{q}_i^\top \mathbf{a}_j)}_1\end{aligned}$$

GRAM-SCHMIDT ORTHOGONALIZATION

Prove that $\mathbf{q}_i \perp \mathbf{y}_j$, $\forall i < j$: Proof-by-induction

- Prove that $\mathbf{q}_1 \perp \mathbf{y}_2$ hence $\mathbf{q}_1 \perp \mathbf{q}_2$ (Exercise)
- Assume $\mathbf{q}_i \perp \mathbf{q}_k$, $\forall k \leq j-1, k \neq i$,

Prove $\mathbf{q}_i \perp \mathbf{y}_j$:

$$\begin{aligned}\mathbf{q}_i^\top \mathbf{y}_j &= \mathbf{q}_i^\top (\mathbf{a}_j - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j) - \mathbf{q}_2(\mathbf{q}_2^\top \mathbf{a}_j) - \dots - \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)) \\&= \mathbf{q}_i^\top \mathbf{a}_j - \underbrace{\mathbf{q}_i^\top \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_j)}_0 - \dots - \mathbf{q}_i^\top \mathbf{q}_i(\mathbf{q}_i^\top \mathbf{a}_j) - \dots - \underbrace{\mathbf{q}_i^\top \mathbf{q}_{j-1}(\mathbf{q}_{j-1}^\top \mathbf{a}_j)}_0 \\&= \mathbf{q}_i^\top \mathbf{a}_j - \underbrace{\mathbf{q}_i^\top \mathbf{q}_i(\mathbf{q}_i^\top \mathbf{a}_j)}_1 \\&= \mathbf{q}_i^\top \mathbf{a}_j - \mathbf{q}_i^\top \mathbf{a}_j \\&= 0\end{aligned}$$

QR FACTORIZATION BY GRAM-SCHMIDT

Define $r_{jj} = \|\mathbf{y}_j\|$ and $r_{ij} = \mathbf{q}_i^\top \mathbf{a}_j$.

QR FACTORIZATION BY GRAM-SCHMIDT

Define $r_{jj} = \|\mathbf{y}_j\|$ and $r_{ij} = \mathbf{q}_i^\top \mathbf{a}_j$.

$$\mathbf{a}_1 = r_{11}\mathbf{q}_1$$

$$\mathbf{a}_2 = r_{12}\mathbf{q}_1 + r_{22}\mathbf{q}_2$$

$$\vdots$$

$$\mathbf{a}_j = r_{1j}\mathbf{q}_1 + r_{2j}\mathbf{q}_2 + \dots + r_{jj}\mathbf{q}_j$$

QR FACTORIZATION BY GRAM-SCHMIDT

Define $r_{jj} = \|\mathbf{y}_j\|$ and $r_{ij} = \mathbf{q}_i^\top \mathbf{a}_j$.

$$\mathbf{a}_1 = r_{11}\mathbf{q}_1$$

$$\mathbf{a}_2 = r_{12}\mathbf{q}_1 + r_{22}\mathbf{q}_2$$

$$\vdots$$

$$\mathbf{a}_j = r_{1j}\mathbf{q}_1 + r_{2j}\mathbf{q}_2 + \dots + r_{jj}\mathbf{q}_j$$

$$\underbrace{\begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_n \end{bmatrix}}_{\mathbf{A}_{m \times n}} = \underbrace{\begin{bmatrix} \mathbf{q}_1 & \mathbf{q}_2 & \dots & \mathbf{q}_n \end{bmatrix}}_{\mathbf{Q}_{m \times n}} \underbrace{\begin{bmatrix} r_{11} & r_{12} & \dots & r_{1n} \\ 0 & r_{22} & \dots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & r_{nn} \end{bmatrix}}_{\mathbf{R}_{n \times n}}$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} = \frac{\mathbf{y}_1}{3} = \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix}$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} = \frac{\mathbf{y}_1}{3} = \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2)$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} = \frac{\mathbf{y}_1}{3} = \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2) = \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} - \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix} \underbrace{\left(\begin{bmatrix} \frac{1}{3} & \frac{2}{3} & \frac{2}{3} \end{bmatrix} \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} \right)}_2 = \begin{bmatrix} -\frac{14}{3} \\ \frac{5}{3} \\ \frac{2}{3} \end{bmatrix},$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} = \frac{\mathbf{y}_1}{3} = \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2) = \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} - \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix} \underbrace{\left(\begin{bmatrix} \frac{1}{3} & \frac{2}{3} & \frac{2}{3} \end{bmatrix} \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} \right)}_2 = \begin{bmatrix} -\frac{14}{3} \\ \frac{5}{3} \\ \frac{2}{3} \end{bmatrix},$$

$$\mathbf{q}_2 = \frac{\mathbf{y}_2}{\|\mathbf{y}_2\|} = \frac{\mathbf{y}_2}{5} = \begin{bmatrix} -\frac{14}{15} \\ \frac{1}{3} \\ \frac{2}{15} \end{bmatrix}$$

EXAMPLE

Find the reduced QR factorization of $\mathbf{A} = \begin{bmatrix} 1 & -4 \\ 2 & 3 \\ 2 & 2 \end{bmatrix}$.

$$\mathbf{y}_1 = \mathbf{a}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{q}_1 = \frac{\mathbf{y}_1}{\|\mathbf{y}_1\|} = \frac{\mathbf{y}_1}{3} = \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix}$$

$$\mathbf{y}_2 = \mathbf{a}_2 - \mathbf{q}_1(\mathbf{q}_1^\top \mathbf{a}_2) = \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} - \begin{bmatrix} \frac{1}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{bmatrix} \underbrace{\left(\begin{bmatrix} \frac{1}{3} & \frac{2}{3} & \frac{2}{3} \end{bmatrix} \begin{bmatrix} -4 \\ 3 \\ 2 \end{bmatrix} \right)}_2 = \begin{bmatrix} -\frac{14}{3} \\ \frac{5}{3} \\ \frac{2}{3} \end{bmatrix},$$

$$\mathbf{q}_2 = \frac{\mathbf{y}_2}{\|\mathbf{y}_2\|} = \frac{\mathbf{y}_2}{5} = \begin{bmatrix} -\frac{14}{15} \\ \frac{1}{3} \\ \frac{2}{15} \end{bmatrix} \rightarrow \mathbf{A} = \begin{bmatrix} \frac{1}{3} & -\frac{14}{15} \\ \frac{2}{3} & \frac{1}{3} \\ \frac{2}{3} & \frac{2}{15} \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 0 & 5 \end{bmatrix}$$

SUMMARY OF REDUCED QR FACTORIZATION WITH GRAM-SCHMIDT

Input: $\mathbf{A}_{m \times n} = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \cdots \quad \mathbf{a}_n]$ with linearly independent columns.

```
for  $j \leftarrow 1$  to  $n$  do
   $\mathbf{y} = \mathbf{a}_j$ 
  for  $i \leftarrow 1$  to
     $j - 1$  do
     $r_{ij} = \mathbf{q}_i^\top \mathbf{a}_j$ 
     $\mathbf{y} =$ 
     $\mathbf{y} - r_{ij} \mathbf{q}_i$ 
  end
   $r_{jj} = \|\mathbf{y}\|$ 
   $\mathbf{q}_j = \frac{\mathbf{y}}{r_{jj}}$ 
end
```

Output: $\mathbf{Q}_{m \times n}$ and $\mathbf{R}_{n \times n}$ such that

$$\mathbf{A}_{m \times n} = [\mathbf{q}_1 \quad \mathbf{q}_2 \quad \cdots \quad \mathbf{q}_n] \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{nn} \end{bmatrix}$$
$$= \mathbf{Q}_{m \times n} \mathbf{R}_{n \times n}$$