

CENG 216 – NUMERICAL COMPUTATION

SOLVING EQUATIONS

Mustafa Özuysal

`mustafaozuysal@iyte.edu.tr`

March 24, 2022

İzmir Institute of Technology

Slides are based on material from the main textbook:

“Numerical Analysis”, The new international edition, 2ed,
by Timothy Sauer

INTRODUCTION

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

$$\cdot f(x) = g(x) \rightarrow f(x) - g(x) = 0$$

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$
 - Initialization: How to pick x_0 ?

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$
 - Initialization: How to pick x_0 ?
 - Convergence: Will x_N be a solution for large enough N ?

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$
 - Initialization: How to pick x_0 ?
 - Convergence: Will x_N be a solution for large enough N ?
 - Accuracy: How to decide if x_N is a good solution?

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$
 - Initialization: How to pick x_0 ?
 - Convergence: Will x_N be a solution for large enough N ?
 - Accuracy: How to decide if x_N is a good solution?
 - Convergence Speed: How large should N be?

ITERATIVE NUMERICAL SOLUTION OF $f(x) = 0$

- $f(x) = g(x) \rightarrow f(x) - g(x) = 0$
- Numerical
 - As opposed to symbolic (algebraic) computation
 - As opposed to analytic (closed-form) solution
 - Computer Algebra Systems (CAS)
- Iterative
 - $x_0 \rightarrow x_1 \rightarrow \cdots \rightarrow x_i \rightarrow \cdots \rightarrow x_N$
 - Initialization: How to pick x_0 ?
 - Convergence: Will x_N be a solution for large enough N ?
 - Accuracy: How to decide if x_N is a good solution?
 - Convergence Speed: How large should N be?
 - Complexity: Evaluation of $f(x_i)$ and possibly $f'(x_i), f''(x_i), \dots$

THE BISECTION METHOD

DEFINITION AND EXISTENCE OF ROOTS

- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.

DEFINITION AND EXISTENCE OF ROOTS


- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.
- How to check existence of a root in a given interval $[a, b]$?

DEFINITION AND EXISTENCE OF ROOTS

- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.
- How to check existence of a root in a given interval $[a, b]$?
- **Thm:** Let $f(x)$ be a continuous function on $[a, b]$ such that $f(a)f(b) < 0$ then $f(x)$ has a root in $[a, b]$.


DEFINITION AND EXISTENCE OF ROOTS

- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.
- How to check existence of a root in a given interval $[a, b]$?
- **Thm:** Let $f(x)$ be a continuous function on $[a, b]$ such that $f(a)f(b) < 0$ then $f(x)$ has a root in $[a, b]$.




DEFINITION AND EXISTENCE OF ROOTS

- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.
- How to check existence of a root in a given interval $[a, b]$?
- **Thm:** Let $f(x)$ be a continuous function on $[a, b]$ such that $f(a)f(b) < 0$ then $f(x)$ has a root in $[a, b]$.



DEFINITION AND EXISTENCE OF ROOTS

- **Def:** The function $f(x)$ has a **root** at $x = r$ if $f(r) = 0$.
- How to check existence of a root in a given interval $[a, b]$?
- **Thm:** Let $f(x)$ be a continuous function on $[a, b]$ such that $f(a)f(b) < 0$ then $f(x)$ has a root in $[a, b]$.



THE BISECTION IDEA

- Search in an address book


THE BISECTION IDEA

- Search in an address book → Binary Search

THE BISECTION IDEA

- Search in an address book \rightarrow Binary Search
- Bisection: Given $[a, b]$ such that $f(a)f(b) < 0$

while $\frac{(b-a)}{2} > TOL$ **do**



end

THE BISECTION IDEA

- Search in an address book \rightarrow Binary Search
- Bisection: Given $[a, b]$ such that $f(a)f(b) < 0$

while $\frac{(b-a)}{2} > TOL$ **do**

$$c = \frac{(a+b)}{2}$$

end

THE BISECTION IDEA

- Search in an address book \rightarrow Binary Search
- Bisection: Given $[a, b]$ such that $f(a)f(b) < 0$

```
while  $\frac{(b-a)}{2} > TOL$  do
```

```
   $c = \frac{(a+b)}{2}$ 
```

```
  if  $f(c) = 0$  then
```

```
    | stop
```

```
  end
```

```
end
```


THE BISECTION IDEA

- Search in an address book \rightarrow Binary Search
- Bisection: Given $[a, b]$ such that $f(a)f(b) < 0$

```
while  $\frac{(b-a)}{2} > TOL$  do  
   $c = \frac{(a+b)}{2}$   
  if  $f(c) = 0$  then  
    | stop  
  end  
  if  $f(a)f(c) < 0$  then  
    |  $b = c$   
  else  
    |  $a = c$   
  end  
end
```

THE BISECTION IDEA

- Search in an address book \rightarrow Binary Search
- Bisection: Given $[a, b]$ such that $f(a)f(b) < 0$

```
while  $\frac{(b-a)}{2} > TOL$  do
   $c = \frac{(a+b)}{2}$ 
  if  $f(c) = 0$  then
    | stop
  end
  if  $f(a)f(c) < 0$  then
    |  $b = c$ 
  else
    |  $a = c$ 
  end
end
```

- The final interval $[a, b]$ contains the root, which is approximately at $\frac{a+b}{2}$.

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
-----	-----	--------	-----	--------	-----	--------	-------------

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125
5	0.65625	-0.06113	0.68750	0.01245	0.67187	-0.02483	0.01562

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125
5	0.65625	-0.06113	0.68750	0.01245	0.67187	-0.02483	0.01562
6	0.67187	-0.02483	0.68750	0.01245	0.67969	-0.00631	0.00781

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125
5	0.65625	-0.06113	0.68750	0.01245	0.67187	-0.02483	0.01562
6	0.67187	-0.02483	0.68750	0.01245	0.67969	-0.00631	0.00781
7	0.67969	-0.00631	0.68750	0.01245	0.68359	0.00304	0.00391

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125
5	0.65625	-0.06113	0.68750	0.01245	0.67187	-0.02483	0.01562
6	0.67187	-0.02483	0.68750	0.01245	0.67969	-0.00631	0.00781
7	0.67969	-0.00631	0.68750	0.01245	0.68359	0.00304	0.00391
8	0.67969	-0.00631	0.68359	0.00304	0.68164	-0.00165	0.00195

EXAMPLE

Find the root of the function $f(x) = x^3 + x - 1$ in $[0, 1]$.

i	a	$f(a)$	b	$f(b)$	c	$f(c)$	$(b - a)/2$
0	0.00000	-1.00000	1.00000	1.00000	0.50000	-0.37500	0.50000
1	0.50000	-0.37500	1.00000	1.00000	0.75000	0.17187	0.25000
2	0.50000	-0.37500	0.75000	0.17187	0.62500	-0.13086	0.12500
3	0.62500	-0.13086	0.75000	0.17187	0.68750	0.01245	0.06250
4	0.62500	-0.13086	0.68750	0.01245	0.65625	-0.06113	0.03125
5	0.65625	-0.06113	0.68750	0.01245	0.67187	-0.02483	0.01562
6	0.67187	-0.02483	0.68750	0.01245	0.67969	-0.00631	0.00781
7	0.67969	-0.00631	0.68750	0.01245	0.68359	0.00304	0.00391
8	0.67969	-0.00631	0.68359	0.00304	0.68164	-0.00165	0.00195

$$r_c = 0.682617, \quad \max. \text{ error} = \pm 0.000977$$

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$:

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$:

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$
- ...
- At $i = n$:

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** $= |x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$
- ...
- At $i = n$: Absolute Error $< \frac{b-a}{2^{n+1}}$

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$
- ...
- At $i = n$: Absolute Error $< \frac{b-a}{2^{n+1}}$
- Number of required function evaluations is

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$
- ...
- At $i = n$: Absolute Error $< \frac{b-a}{2^{n+1}}$
- Number of required function evaluations is $n + 2$.

ACCURACY AND CONVERGENCE SPEED

- Reminder: **Absolute Error** = $|x_c - r|$, where x_c is the computed root.
- At $i = 0$: Absolute Error $< \frac{b-a}{2}$
- At $i = 1$: Absolute Error $< \frac{b-a}{2^2}$
- ...
- At $i = n$: Absolute Error $< \frac{b-a}{2^{n+1}}$
- Number of required function evaluations is $n + 2$.
- **Def:** A solution is correct to p decimal places if the error is less than $0.5 \cdot 10^{-p}$.

EXAMPLE

Find the number of iterations of the bisection method required to compute a root of the function $f(x) = \cos x - x$ in $[0, 1]$ to six correct decimal places.

EXAMPLE

Find the number of iterations of the bisection method required to compute a root of the function $f(x) = \cos x - x$ in $[0, 1]$ to six correct decimal places.

$$\frac{1}{2^{n+1}} < 0.5 \cdot 10^{-6}$$

EXAMPLE

Find the number of iterations of the bisection method required to compute a root of the function $f(x) = \cos x - x$ in $[0, 1]$ to six correct decimal places.

$$\frac{1}{2^{n+1}} < 0.5 \cdot 10^{-6} \rightarrow n > \frac{6}{\log_{10} 2}$$

EXAMPLE

Find the number of iterations of the bisection method required to compute a root of the function $f(x) = \cos x - x$ in $[0, 1]$ to six correct decimal places.

$$\frac{1}{2^{n+1}} < 0.5 \cdot 10^{-6} \rightarrow n > \frac{6}{\log_{10} 2} \approx \frac{6}{0.301} \approx 19.9$$

EXAMPLE

Find the number of iterations of the bisection method required to compute a root of the function $f(x) = \cos x - x$ in $[0, 1]$ to six correct decimal places.

$$\frac{1}{2^{n+1}} < 0.5 \cdot 10^{-6} \rightarrow n > \frac{6}{\log_{10} 2} \approx \frac{6}{0.301} \approx 19.9$$

Take $n = 20$.

FIXED POINT ITERATIONS

FIXED POINTS

- **Def:** A real number r is a fixed point of $f(x)$ if $f(r) = r$.

FIXED POINTS

- **Def:** A real number r is a fixed point of $f(x)$ if $f(r) = r$.
- Fixed point iterations:

$$x_0 = \text{"initial guess"}$$

$$x_{i+1} = f(x_i), \quad \forall i = 0, 1, 2, \dots$$

FIXED POINTS

- **Def:** A real number r is a fixed point of $f(x)$ if $f(r) = r$.
- Fixed point iterations:

$$x_0 = \text{"initial guess"}$$

$$x_{i+1} = f(x_i), \quad \forall i = 0, 1, 2, \dots$$

- $f(x) = \cos(x)$, $x_0 = 0.0$:

$$x_1 = \cos(0.0) = 1.0$$

FIXED POINTS

- **Def:** A real number r is a fixed point of $f(x)$ if $f(r) = r$.
- Fixed point iterations:

$$x_0 = \text{"initial guess"}$$

$$x_{i+1} = f(x_i), \quad \forall i = 0, 1, 2, \dots$$

- $f(x) = \cos(x)$, $x_0 = 0.0$:

$$x_1 = \cos(0.0) = 1.0$$

$$x_2 = \cos(1.0) = 0.5403023058681398$$

FIXED POINTS

- **Def:** A real number r is a fixed point of $f(x)$ if $f(r) = r$.
- Fixed point iterations:

$$x_0 = \text{"initial guess"}$$

$$x_{i+1} = f(x_i), \quad \forall i = 0, 1, 2, \dots$$

- $f(x) = \cos(x)$, $x_0 = 0.0$:

$$x_1 = \cos(0.0) = 1.0$$

$$x_2 = \cos(1.0) = 0.5403023058681398$$

$$\vdots$$

$$x_k = \cos(0.739085133215) = 0.739085133215$$

ROOT FINDING EXAMPLE

Find a root of the function $f(x) = x^3 + x - 1.0$.

$$0 = x^3 + x - 1.0$$

$$x = 1.0 - x^3 = p(x)$$

ROOT FINDING EXAMPLE

Find a root of the function $f(x) = x^3 + x - 1.0$.

$$0 = x^3 + x - 1.0$$

$$x = 1.0 - x^3 = p(x)$$

$$0 = y^3 + y - 1.0$$

$$y^3 = 1.0 - y$$

$$y = \sqrt[3]{1.0 - y} = q(y)$$

ROOT FINDING EXAMPLE

Find a root of the function $f(x) = x^3 + x - 1.0$.

$$0 = x^3 + x - 1.0$$

$$x = 1.0 - x^3 = p(x)$$

$$0 = y^3 + y - 1.0$$

$$y^3 = 1.0 - y$$

$$y = \sqrt[3]{1.0 - y} = q(y)$$

$$0 = z^3 + z - 1.0$$

$$z = 1 - z^3$$

$$3z^3 + z = 1 + 2z^3$$

$$(3z^2 + 1)z = 1 + 2z^3$$

$$z = \frac{1 + 2z^3}{3z^2 + 1} = r(z)$$

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780
6	0.00337606	0.65900615	0.68232780

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780
6	0.00337606	0.65900615	0.68232780
7	0.99999996	0.69863261	0.68232780

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780
6	0.00337606	0.65900615	0.68232780
7	0.99999996	0.69863261	0.68232780
8	0.00000012	0.67044850	0.68232780

ROOT FINDING EXAMPLE


i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780
6	0.00337606	0.65900615	0.68232780
7	0.99999996	0.69863261	0.68232780
8	0.00000012	0.67044850	0.68232780
9	1.00000000	0.69072912	0.68232780
10	0.00000000	0.67625892	0.68232780
11	1.00000000	0.68664554	0.68232780
12	0.00000000	0.67922234	0.68232780

ROOT FINDING EXAMPLE

i	$x_i = p(x_{i-1})$	$y_i = q(y_{i-1})$	$z_i = r(z_{i-1})$
0	0.50000000	0.50000000	0.50000000
1	0.87500000	0.79370053	0.71428571
2	0.33007813	0.59088011	0.68317972
3	0.96403747	0.74236393	0.68232842
4	0.10405419	0.63631020	0.68232780
5	0.99887338	0.71380081	0.68232780
6	0.00337606	0.65900615	0.68232780
7	0.99999996	0.69863261	0.68232780
8	0.00000012	0.67044850	0.68232780
9	1.00000000	0.69072912	0.68232780
10	0.00000000	0.67625892	0.68232780
11	1.00000000	0.68664554	0.68232780
12	0.00000000	0.67922234	0.68232780
\vdots	\vdots	\vdots	\vdots
24	0.00000000	0.68227157	0.68232780
25	1.00000000	0.68236807	0.68232780


THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams




THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams




THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams




THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams




THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams





THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams





THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams





THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams





THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams






THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams






THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams






THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams






THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams



THE GEOMETRY OF FIXED POINT ITERATION

Cobweb Diagrams



LINEAR CONVERGENCE OF FIXED POINT ITERATIONS

- **Def:** Let $e_i = |x_i - r|$ be the error at step i . If

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i} = s < 1$$

then the method is linearly convergent with rate s .

LINEAR CONVERGENCE OF FIXED POINT ITERATIONS

- **Def:** Let $e_i = |x_i - r|$ be the error at step i . If

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i} = s < 1$$

then the method is linearly convergent with rate s .

- **Thm:** Assume $f(x)$ is continuously differentiable,

$$f(r) = r \text{ and } s = |f'(r)| < 1$$

then fixed point iterations converges linearly with rate s to the fixed point r for initial guesses sufficiently close to r .

LINEAR CONVERGENCE OF FIXED POINT ITERATIONS

- **Def:** Let $e_i = |x_i - r|$ be the error at step i . If

$$\lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i} = s < 1$$

then the method is linearly convergent with rate s .

- **Thm:** Assume $f(x)$ is continuously differentiable,

$$f(r) = r \text{ and } s = |f'(r)| < 1$$

then fixed point iterations converges linearly with rate s to the fixed point r for initial guesses sufficiently close to r .


- $f(x) = \cos x, f'(x) = -\sin x, f'(0.739085133215) \approx -0.67$.

EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.


EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.




EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.




EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



$$x_0 = 0.1000$$

$$x_1 = 0.2700$$

$$x_2 = 0.6831$$


$$x_3 = 1.4461$$

$$x_4 = 1.9579$$

$$x_5 = 1.6487$$

EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



$$x_0 = 0.1000$$

$$x_1 = 0.2700$$

$$x_2 = 0.6831$$

$$x_3 = 1.4461$$

$$x_4 = 1.9579$$

$$x_5 = 1.6487$$


$$\vdots$$

$$x_{11} = 1.7660$$

$$x_{12} = 1.8260$$

EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



$$x_0 = 0.1000$$

$$x_1 = 0.2700$$

$$x_2 = 0.6831$$

$$x_3 = 1.4461$$

$$x_4 = 1.9579$$

$$x_5 = 1.6487$$

$$\vdots$$


$$x_{11} = 1.7660$$

$$x_{12} = 1.8260$$

$$f'(x) = 2.8 - 2x$$

EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



$$x_0 = 0.1000$$

$$x_1 = 0.2700$$

$$x_2 = 0.6831$$

$$x_3 = 1.4461$$

$$x_4 = 1.9579$$

$$x_5 = 1.6487$$

$$\vdots$$

$$x_{11} = 1.7660$$


$$x_{12} = 1.8260$$

$$f'(x) = 2.8 - 2x$$

$$f'(1.8) = -0.8$$

EXAMPLE

Find the fixed points of $2.8x - x^2 = f(x)$.



$$x_0 = 0.1000$$

$$x_1 = 0.2700$$

$$x_2 = 0.6831$$

$$x_3 = 1.4461$$

$$x_4 = 1.9579$$

$$x_5 = 1.6487$$

$$\vdots$$

$$x_{11} = 1.7660$$

$$x_{12} = 1.8260$$

$$f'(x) = 2.8 - 2x$$

$$f'(1.8) = -0.8$$

$$f'(0.0) = 2.8$$

STOPPING CRITERIA

$$|x_{i+1} - x_i| < TOL$$

STOPPING CRITERIA

$$|x_{i+1} - x_i| < TOL$$

$$\frac{|x_{i+1} - x_i|}{|x_{i+1}|} < TOL$$

STOPPING CRITERIA


$$|x_{i+1} - x_i| < TOL$$

$$\frac{|x_{i+1} - x_i|}{|x_{i+1}|} < TOL$$


$$\frac{|x_{i+1} - x_i|}{\max(|x_{i+1}|, \theta)} < TOL, \text{ with } \theta > 0.0$$

NEWTON'S METHOD

NEWTON-RAPHSON APPROACH




NEWTON-RAPHSON APPROACH



$$y - f(x_i) = f'(x_i)(x - x_i)$$


NEWTON-RAPHSON APPROACH



$$y - f(x_i) = f'(x_i)(x - x_i)$$

$$y_1 = 0 \implies x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$

NEWTON-RAPHSON APPROACH



$$y - f(x_i) = f'(x_i)(x - x_i)$$

$$y_1 = 0 \implies x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$

Alternatively,

$$0 = f(\overbrace{x_i + \Delta x_i}^{x_{i+1}})$$

$$0 = f(x_i) + \Delta x_i f'(x_i) + H.O.T$$

$$\Delta x_i \approx -\frac{f(x_i)}{f'(x_i)}$$

EXAMPLE

Find a root of the equation $x^3 + x - 1$.

$$f'(x) = 3x^2 + 1$$

EXAMPLE

Find a root of the equation $x^3 + x - 1$.

$$f'(x) = 3x^2 + 1$$

$$x_{i+1} = x_i - \frac{x_i^3 + x_i - 1}{3x_i^2 + 1}$$

$$= \frac{2x_i^3 + 1}{3x_i^2 + 1}$$

EXAMPLE

Find a root of the equation $x^3 + x - 1$.

$$f'(x) = 3x^2 + 1$$

$$x_{i+1} = x_i - \frac{x_i^3 + x_i - 1}{3x_i^2 + 1}$$

$$= \frac{2x_i^3 + 1}{3x_i^2 + 1}$$

$$x_0 = -0.7$$

$$x_1 = 0.12712551$$

$$\vdots$$

$$x_6 = 0.68232780$$

$$x_7 = 0.68232780$$

QUADRATIC CONVERGENCE OF NEWTON'S METHOD

- **Def:** Iterations are quadratically convergent if

$$M = \lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} < \infty$$

QUADRATIC CONVERGENCE OF NEWTON'S METHOD

- **Def:** Iterations are quadratically convergent if

$$M = \lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} < \infty$$

- **Thm:** Let f be twice continuously differentiable and $f(r) = 0$. If $f'(r) \neq 0$ then Newton's Method is quadratically convergent to r with $M = \frac{f''(r)}{2f'(r)}$.

QUADRATIC CONVERGENCE OF NEWTON'S METHOD

- **Def:** Iterations are quadratically convergent if

$$M = \lim_{i \rightarrow \infty} \frac{e_{i+1}}{e_i^2} < \infty$$

- **Thm:** Let f be twice continuously differentiable and $f(r) = 0$. If $f'(r) \neq 0$ then Newton's Method is quadratically convergent to r with $M = \frac{f''(r)}{2f'(r)}$.
- If $f'(r) = 0$, which happens in case of double roots, convergence is linear. See Modified Newton's Method in the textbook.

EXAMPLE OF NON-CONVERGENCE

Apply Newton's Method to $f(x) = 4x^4 - 6x^2 - \frac{11}{4}$ starting at $x_0 = \frac{1}{2}$

EXAMPLE OF NON-CONVERGENCE

Apply Newton's Method to $f(x) = 4x^4 - 6x^2 - \frac{11}{4}$ starting at $x_0 = \frac{1}{2}$

$$x_{i+1} = x_i - \frac{4x_i^4 - 6x_i^2 - \frac{11}{4}}{16x_i^3 - 12x_i}$$

EXAMPLE OF NON-CONVERGENCE

Apply Newton's Method to $f(x) = 4x^4 - 6x^2 - \frac{11}{4}$ starting at $x_0 = \frac{1}{2}$

$$x_{i+1} = x_i - \frac{4x_i^4 - 6x_i^2 - \frac{11}{4}}{16x_i^3 - 12x_i}$$

$$x_1 = -\frac{1}{2} \text{ and } x_2 = \frac{1}{2}$$

LIMITS OF ACCURACY

FORWARD AND BACKWARD ERROR

Use Bisection Method to find the root of

$$f(x) = x^3 - 2x^2 + \frac{4}{3} - \frac{8}{27} \text{ to within six correct digits.}$$

FORWARD AND BACKWARD ERROR

Use Bisection Method to find the root of

$f(x) = x^3 - 2x^2 + \frac{4}{3} - \frac{8}{27}$ to within six correct digits.

$$\left. \begin{array}{l} f(0) = -\frac{8}{27} \\ f(1) = \frac{1}{27} \end{array} \right\}$$

FORWARD AND BACKWARD ERROR

Use Bisection Method to find the root of
 $f(x) = x^3 - 2x^2 + \frac{4}{3} - \frac{8}{27}$ to within six correct digits.

$$\left. \begin{array}{l} f(0) = -\frac{8}{27} \\ f(1) = \frac{1}{27} \end{array} \right\} \text{Solution in } [0, 1]$$

FORWARD AND BACKWARD ERROR

Use Bisection Method to find the root of

$f(x) = x^3 - 2x^2 + \frac{4}{3} - \frac{8}{27}$ to within six correct digits.

$$\left. \begin{array}{l} f(0) = -\frac{8}{27} \\ f(1) = \frac{1}{27} \end{array} \right\} \text{Solution in } [0, 1] \rightarrow f\left(\frac{2}{3}\right) = f(0.6\bar{6}) = 0$$

FORWARD AND BACKWARD ERROR

Use Bisection Method to find the root of

$f(x) = x^3 - 2x^2 + \frac{4}{3} - \frac{8}{27}$ to within six correct digits.

$$\left. \begin{array}{l} f(0) = -\frac{8}{27} \\ f(1) = \frac{1}{27} \end{array} \right\} \text{Solution in } [0, 1] \rightarrow f\left(\frac{2}{3}\right) = f(0.\overline{66}) = 0$$

Due to rounding errors $f(0.6666641) = 0$ so Bisection terminates with $r = 0.6666641$ which is close in y but not in x .

FORWARD AND BACKWARD ERROR

Def: Assume f is a function and r is a root such that $f(r) = 0$.
Assume x_a is an approximation of r .

FORWARD AND BACKWARD ERROR

Def: Assume f is a function and r is a root such that $f(r) = 0$. Assume x_a is an approximation of r . For the root finding problem:

- backward error = $|f(x_a)|$

FORWARD AND BACKWARD ERROR

Def: Assume f is a function and r is a root such that $f(r) = 0$. Assume x_a is an approximation of r . For the root finding problem:

- backward error = $|f(x_a)|$
- forward error = $|r - x_a|$

FORWARD AND BACKWARD ERROR

Def: Assume f is a function and r is a root such that $f(r) = 0$. Assume x_a is an approximation of r . For the root finding problem:

- backward error = $|f(x_a)|$
- forward error = $|r - x_a|$

$x_a = 0.6666641 \rightarrow$ backward error $\approx \epsilon_{\text{mach}}$, forward error $\approx 10^{-5}$.

EXAMPLE: WILKINSON POLYNOMIAL

$$\begin{aligned} W(x) &= (x - 1)(x - 2) \cdots (x - 20) \\ &= x^{20} + ax^{19} + bx^{18} + \dots \end{aligned}$$

EXAMPLE: WILKINSON POLYNOMIAL

$$\begin{aligned}W(x) &= (x - 1)(x - 2) \cdots (x - 20) \\&= x^{20} + ax^{19} + bx^{18} + \dots\end{aligned}$$

$$x_0 = 16.0$$

$$x_a = 16.01468 \dots$$

SENSITIVITY OF ROOT FINDING

Def: A problem is called **sensitive** if small error in the input lead to large error in the solution.

SENSITIVITY OF ROOT FINDING

Def: A problem is called **sensitive** if small error in the input lead to large error in the solution.

$$r \rightarrow f(x) = 0$$

$$r + \Delta r \rightarrow f(x) + \epsilon g(x) = 0$$

SENSITIVITY OF ROOT FINDING

Def: A problem is called **sensitive** if small error in the input lead to large error in the solution.

$$r \rightarrow f(x) = 0$$

$$r + \Delta r \rightarrow f(x) + \epsilon g(x) = 0$$

$$f(r + \Delta r) + \epsilon g(r + \Delta r) = 0$$

$$f(r) + \Delta r f'(r) + \epsilon g(r) + \epsilon \Delta r g'(r) + \mathcal{O}(\Delta r^2) = 0$$

SENSITIVITY OF ROOT FINDING

Def: A problem is called **sensitive** if small error in the input lead to large error in the solution.

$$r \rightarrow f(x) = 0$$

$$r + \Delta r \rightarrow f(x) + \epsilon g(x) = 0$$

$$f(r + \Delta r) + \epsilon g(r + \Delta r) = 0$$

$$f(r) + \Delta r f'(r) + \epsilon g(r) + \epsilon \Delta r g'(r) + \mathcal{O}(\Delta r^2) = 0$$

$$\Delta r (f'(r) + \epsilon g'(r)) \approx -f(r) - \epsilon g(r) = -\epsilon g(r)$$

SENSITIVITY OF ROOT FINDING

Def: A problem is called **sensitive** if small error in the input lead to large error in the solution.

$$r \rightarrow f(x) = 0$$

$$r + \Delta r \rightarrow f(x) + \epsilon g(x) = 0$$

$$f(r + \Delta r) + \epsilon g(r + \Delta r) = 0$$

$$f(r) + \Delta r f'(r) + \epsilon g(r) + \epsilon \Delta r g'(r) + \mathcal{O}(\Delta r^2) = 0$$

$$\Delta r (f'(r) + \epsilon g'(r)) \approx -f(r) - \epsilon g(r) = -\epsilon g(r)$$

$$\Delta r \approx \frac{-\epsilon g(r)}{f'(r) + \epsilon g'(r)} \approx -\epsilon \frac{g(r)}{f'(r)} \quad \epsilon \ll f'(r), f'(r) \neq 0$$

EXAMPLE

Estimate the largest root of

$$P(x) = (x - 1)(x - 2)(x - 3)(x - 4)(x - 5)(x - 6) - 10^{-6}x^7.$$

EXAMPLE

Estimate the largest root of

$$P(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6) - 10^{-6}x^7.$$

$$f(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6), g(x) = x^7, \epsilon = 10^{-6}.$$

EXAMPLE

Estimate the largest root of

$$P(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6) - 10^{-6}x^7.$$

$$f(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6), g(x) = x^7, \epsilon = 10^{-6}.$$

Largest root of $f(x) = 6$.

EXAMPLE

Estimate the largest root of

$$P(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6) - 10^{-6}x^7.$$

$$f(x) = (x-1)(x-2)(x-3)(x-4)(x-5)(x-6), g(x) = x^7, \epsilon = 10^{-6}.$$

Largest root of $f(x) = 6$.

$$\Delta r \approx -\frac{\epsilon 6^7}{5!} = -2332.6\epsilon \implies r + \Delta r = 6.0023328$$

ERROR MAGNIFICATION FACTOR FOR ROOT FINDING

- Def:

$$\text{error magnification factor} = \frac{\text{relative forward error}}{\text{relative backward error}}$$

ERROR MAGNIFICATION FACTOR FOR ROOT FINDING

- Def:

$$\begin{aligned}\text{error magnification factor} &= \frac{\text{relative forward error}}{\text{relative backward error}} \\ &= \left| \frac{\Delta r / r}{\epsilon g(r) / g(r)} \right| = \frac{|g(r)|}{|r f'(r)|}\end{aligned}$$

- Def: **Condition number** is the maximum error magnification factor over all input changes.

ERROR MAGNIFICATION FACTOR FOR ROOT FINDING

- Def:

$$\begin{aligned}\text{error magnification factor} &= \frac{\text{relative forward error}}{\text{relative backward error}} \\ &= \left| \frac{\Delta r / r}{\epsilon g(r) / g(r)} \right| = \frac{|g(r)|}{|r f'(r)|}\end{aligned}$$

- Def: **Condition number** is the maximum error magnification factor over all input changes.
- High condition number \rightarrow **ill-conditioned** problem.

ERROR MAGNIFICATION FACTOR FOR ROOT FINDING

- Def:

$$\begin{aligned}\text{error magnification factor} &= \frac{\text{relative forward error}}{\text{relative backward error}} \\ &= \left| \frac{\Delta r / r}{\epsilon g(r) / g(r)} \right| = \frac{|g(r)|}{|r f'(r)|}\end{aligned}$$

- Def: **Condition number** is the maximum error magnification factor over all input changes.
- High condition number \rightarrow **ill-conditioned** problem.
- Condition number $\approx 1 \rightarrow$ **well-conditioned** problem.

ERROR MAGNIFICATION FACTOR FOR ROOT FINDING

- Def:

$$\begin{aligned}\text{error magnification factor} &= \frac{\text{relative forward error}}{\text{relative backward error}} \\ &= \left| \frac{\Delta r / r}{\epsilon g(r) / g(r)} \right| = \frac{|g(r)|}{|r f'(r)|}\end{aligned}$$

- Def: **Condition number** is the maximum error magnification factor over all input changes.
- High condition number \rightarrow **ill-conditioned** problem.
- Condition number $\approx 1 \rightarrow$ **well-conditioned** problem.
- We will revisit conditioning for different numerical problems throughout the course.

EXERCISE: BALL THROWING ROBOT ARM

- A ball throwing robot arm throws a ball with initial velocity V_0 at an angle θ from the ground. Assuming that the robot arm is at $x = 0$, the ball will hit the ground at $x_f = \frac{V_0^2 \sin 2\theta}{g}$, where $g = 9.8 \text{ m/s}^2$ is the gravitational constant.

EXERCISE: BALL THROWING ROBOT ARM

- A ball throwing robot arm throws a ball with initial velocity V_0 at an angle θ from the ground. Assuming that the robot arm is at $x = 0$, the ball will hit the ground at $x_f = \frac{V_0^2 \sin 2\theta}{g}$, where $g = 9.8 \text{ m/s}^2$ is the gravitational constant.
- We want to ensure that the ball hits the ground at $x_f = 0.17$ meters.

EXERCISE: BALL THROWING ROBOT ARM

- A ball throwing robot arm throws a ball with initial velocity V_0 at an angle θ from the ground. Assuming that the robot arm is at $x = 0$, the ball will hit the ground at $x_f = \frac{V_0^2 \sin 2\theta}{g}$, where $g = 9.8 \text{ m/s}^2$ is the gravitational constant.
- We want to ensure that the ball hits the ground at $x_f = 0.17$ meters.
- The robot design constrains the speed to be $V_0 = k(1 + \cos \theta)$, where $k = 0.75$ is a design parameter.

EXERCISE: BALL THROWING ROBOT ARM

- A ball throwing robot arm throws a ball with initial velocity V_0 at an angle θ from the ground. Assuming that the robot arm is at $x = 0$, the ball will hit the ground at $x_f = \frac{V_0^2 \sin 2\theta}{g}$, where $g = 9.8 \text{ m/s}^2$ is the gravitational constant.
- We want to ensure that the ball hits the ground at $x_f = 0.17$ meters.
- The robot design constrains the speed to be $V_0 = k(1 + \cos \theta)$, where $k = 0.75$ is a design parameter.
- Solve for θ in the interval $[0^\circ, 40^\circ]$ and calculate the required V_0 .

SOLUTION: BALL THROWING ROBOT ARM

- Writing $V_0 = k(1 + \cos \theta)$ into the equation for x_f , we have

$$x_f = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta.$$

SOLUTION: BALL THROWING ROBOT ARM

- Writing $V_0 = k(1 + \cos \theta)$ into the equation for x_f , we have

$$x_f = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta.$$

- This can solve this as a root finding problem with

$$f(\theta) = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta - x_f = 0$$

SOLUTION: BALL THROWING ROBOT ARM

- Writing $V_0 = k(1 + \cos \theta)$ into the equation for x_f , we have

$$x_f = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta.$$

- This can solve this as a root finding problem with

$$f(\theta) = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta - x_f = 0$$

- $f(0^\circ) = -0.17 < 0$ and $f(40^\circ) \approx 0.0063 > 0$

SOLUTION: BALL THROWING ROBOT ARM

- Writing $V_0 = k(1 + \cos \theta)$ into the equation for x_f , we have

$$x_f = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta.$$

- This can solve this as a root finding problem with

$$f(\theta) = \frac{k^2}{g} (1 + \cos \theta)^2 \sin 2\theta - x_f = 0$$

- $f(0^\circ) = -0.17 < 0$ and $f(40^\circ) \approx 0.0063 > 0$
- After 20 iterations $\theta \approx 28.53^\circ$, $f(\theta) \approx 6.76 \cdot 10^{-8}$, and $V_0 \approx 1.41$ m/s.