

CENG 216 – NUMERICAL COMPUTATION

NUMERICAL DIFFERENTIATION AND INTEGRATION

Mustafa Özuysal

`mustafaozuysal@iyte.edu.tr`

May 22, 2022

İzmir Institute of Technology

Slides are based on material from the main textbook:

“Numerical Analysis”, The new international edition, 2ed,
by Timothy Sauer

NUMERICAL DIFFERENTIATION

THE NEED TO COMPUTE NUMERICAL DERIVATIVES

- When solving problems with methods that require derivatives such as Newton's Method, an explicit analytical formula for $f'(x)$ may not be always available.

THE NEED TO COMPUTE NUMERICAL DERIVATIVES

- When solving problems with methods that require derivatives such as Newton's Method, an explicit analytical formula for $f'(x)$ may not be always available.
- Real-world data is mostly discrete and tabulated so it is not possible to directly compute derivatives using a continuous formula: Sensor data acquired with a fixed or varying time interval.

THE NEED TO COMPUTE NUMERICAL DERIVATIVES

- When solving problems with methods that require derivatives such as Newton's Method, an explicit analytical formula for $f'(x)$ may not be always available.
- Real-world data is mostly discrete and tabulated so it is not possible to directly compute derivatives using a continuous formula: Sensor data acquired with a fixed or varying time interval.
- For some complex functions, analytical derivatives are not available or difficult to compute.

THE NEED TO COMPUTE NUMERICAL DERIVATIVES

- When solving problems with methods that require derivatives such as Newton's Method, an explicit analytical formula for $f'(x)$ may not be always available.
- Real-world data is mostly discrete and tabulated so it is not possible to directly compute derivatives using a continuous formula: Sensor data acquired with a fixed or varying time interval.
- For some complex functions, analytical derivatives are not available or difficult to compute.
- Numerical derivatives are easy to compute, so we can check analytical derivatives by comparing their results to numerical ones.

FINITE DIFFERENCE FORMULAS

Def: Forward Difference Formula

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

FINITE DIFFERENCE FORMULAS

Def: Forward Difference Formula

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Derivation:

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(c) \quad x \leq c \leq x+h, \text{ (Taylor Expansion)}$$

FINITE DIFFERENCE FORMULAS

Def: Forward Difference Formula

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Derivation:

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(c) \quad x \leq c \leq x+h, \text{ (Taylor Expansion)}$$

$$f'(x) = \frac{f(x+h) - f(x)}{h} - \underbrace{\frac{h}{2}f''(c)}_{\mathcal{O}(h)},$$

FINITE DIFFERENCE FORMULAS

Def: Forward Difference Formula

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Derivation:

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(c) \quad x \leq c \leq x+h, \text{ (Taylor Expansion)}$$

$$f'(x) = \frac{f(x+h) - f(x)}{h} - \underbrace{\frac{h}{2}f''(c)}_{\mathcal{O}(h)},$$

$$\approx \frac{f(x+h) - f(x)}{h}$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(x) = \frac{f(x + 0.1) - f(x)}{0.1}$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(x) = \frac{f(x + 0.1) - f(x)}{0.1}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{2}}{0.1}$$

$$\approx -0.2381 \quad (\text{Forward difference result})$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(x) = \frac{f(x + 0.1) - f(x)}{0.1}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{2}}{0.1}$$

$$\approx -0.2381 \quad (\text{Forward difference result})$$

$$f'(x) = -x^{-2} \rightarrow f'(2) = -0.25 \quad (\text{Analytical formula})$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(x) = \frac{f(x + 0.1) - f(x)}{0.1}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{2}}{0.1}$$

$$\approx -0.2381 \quad (\text{Forward difference result})$$

$$f'(x) = -x^{-2} \rightarrow f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward error})$$

BACKWARD AND CENTERED DIFFERENCES

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(c) \text{ (Forward difference)}$$

BACKWARD AND CENTERED DIFFERENCES

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(c) \text{ (Forward difference)}$$

$$f(x-h) = f(x) - hf'(x) + \frac{h^2}{2}f''(x) - \frac{h^3}{6}f'''(c) \text{ (Backward difference)}$$

BACKWARD AND CENTERED DIFFERENCES

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(c) \text{ (Forward difference)}$$

$$f(x-h) = f(x) - hf'(x) + \frac{h^2}{2}f''(x) - \frac{h^3}{6}f'''(c) \text{ (Backward difference)}$$

\downarrow

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} - \underbrace{\frac{h^2}{12}f'''(c)}_{\mathcal{O}(h^2)}$$

BACKWARD AND CENTERED DIFFERENCES

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(c) \text{ (Forward difference)}$$

$$f(x-h) = f(x) - hf'(x) + \frac{h^2}{2}f''(x) - \frac{h^3}{6}f'''(c) \text{ (Backward difference)}$$

↓

$$\begin{aligned} f'(x) &= \frac{f(x+h) - f(x-h)}{2h} - \underbrace{\frac{h^2}{12}f'''(c)}_{\mathcal{O}(h^2)} \\ &\approx \frac{f(x+h) - f(x-h)}{2h}, \quad \text{(Centered difference formula)} \end{aligned}$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(2) = -0.2381 \quad (\text{Forward difference result})$$

$$f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward difference error})$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(2) = -0.2381 \quad (\text{Forward difference result})$$

$$f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward difference error})$$

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(2) = -0.2381 \quad (\text{Forward difference result})$$

$$f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward difference error})$$

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{1.9}}{0.2}$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(2) = -0.2381 \quad (\text{Forward difference result})$$

$$f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward difference error})$$

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{1.9}}{0.2}$$

$$\approx -0.2506$$

EXAMPLE

Calculate the approximate derivative of $f(x) = \frac{1}{x}$ at $x = 2$ with step size $h = 0.1$.

$$f'(2) = -0.2381 \quad (\text{Forward difference result})$$

$$f'(2) = -0.25 \quad (\text{Analytical formula})$$

$$\| -0.2381 - (-0.25) \| = 0.0119 \quad (\text{Forward difference error})$$

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

$$f'(2) = \frac{\frac{1}{2.1} - \frac{1}{1.9}}{0.2}$$

$$\approx -0.2506$$

$$\| -0.2506 - (-0.25) \| = 0.0006 \quad (\text{Centered difference error})$$

SECOND DERIVATIVE FORMULAS

$$\begin{aligned} f''(x) &= \frac{f(x-h) - 2f(x) + f(x+h)}{h^2} - \underbrace{\frac{h^2}{12} f^{(iv)}(c)}_{\mathcal{O}(h^2)} \\ &\approx \frac{f(x-h) - 2f(x) + f(x+h)}{h^2} \end{aligned}$$

ROUNDING ERRORS IN FINITE DIFFERENCES

Ex: Approximate the derivative of $f(x) = e^x$ at $x = 0$.

$$f'(x) \approx \frac{e^{x+h} - e^x}{h}, \quad (\star)$$

$$f'(x) \approx \frac{e^{x+h} - e^{x-h}}{2h}, \quad (\star\star)$$

ROUNDING ERRORS IN FINITE DIFFERENCES

Ex: Approximate the derivative of $f(x) = e^x$ at $x = 0$.

$$f'(x) \approx \frac{e^{x+h} - e^x}{h}, \quad (\star)$$

$$f'(x) \approx \frac{e^{x+h} - e^{x-h}}{2h}, \quad (\star\star)$$

h	\star	Error in \star	$\star\star$	Error in $\star\star$
10^{-5}	1.00000500000696	-0.00000500000696	1.00000000001210	-0.00000000001210
10^{-6}	1.00000049996218	-0.00000049996218	0.99999999997324	0.00000000002676
10^{-7}	1.00000004943368	-0.00000004943368	0.99999999947364	0.00000000052636
10^{-8}	0.9999999392253	0.0000000607747	0.9999999392253	0.0000000607747
10^{-9}	1.00000008274037	-0.00000008274037	1.00000002722922	-0.00000002722922
10^{-10}	1.00000008274037	-0.00000008274037	1.00000008274037	-0.00000008274037
10^{-11}	1.00000008274037	-0.00000008274037	1.00000008274037	-0.00000008274037

ROUNDING ERRORS IN FINITE DIFFERENCES

Ex: Approximate the derivative of $f(x) = e^x$ at $x = 0$.

$$f'(x) \approx \frac{e^{x+h} - e^x}{h}, \quad (\star)$$

$$f'(x) \approx \frac{e^{x+h} - e^{x-h}}{2h}, \quad (\star\star)$$

Error drops as h gets smaller but then it starts to increase. Why?

h	\star	Error in \star	$\star\star$	Error in $\star\star$
10^{-5}	1.00000500000696	-0.00000500000696	1.00000000001210	-0.00000000001210
10^{-6}	1.00000049996218	-0.00000049996218	0.99999999997324	0.00000000002676
10^{-7}	1.00000004943368	-0.00000004943368	0.99999999947364	0.00000000052636
10^{-8}	0.99999999392253	0.00000000607747	0.99999999392253	0.00000000607747
10^{-9}	1.00000008274037	-0.00000008274037	1.00000002722922	-0.00000002722922
10^{-10}	1.00000008274037	-0.00000008274037	1.00000008274037	-0.00000008274037
10^{-11}	1.00000008274037	-0.00000008274037	1.00000008274037	-0.00000008274037

ROUNDING ERRORS IN FINITE DIFFERENCES

$$\left. \begin{aligned} \hat{f}(x+h) &= f(x+h) + \epsilon_1 \\ \hat{f}(x-h) &= f(x-h) + \epsilon_2 \end{aligned} \right\} |\epsilon_1|, |\epsilon_2| \approx \epsilon_{\text{mach}}, \text{ rounding errors}$$

ROUNDING ERRORS IN FINITE DIFFERENCES

$$\left. \begin{aligned} \hat{f}(x+h) &= f(x+h) + \epsilon_1 \\ \hat{f}(x-h) &= f(x-h) + \epsilon_2 \end{aligned} \right\} |\epsilon_1|, |\epsilon_2| \approx \epsilon_{\text{mach}}, \text{ rounding errors}$$

$$f'_{\text{correct}}(x) - f'_{\text{machine}}(x) = f'(x) - \frac{\hat{f}(x+h) - \hat{f}(x-h)}{2h}$$

ROUNDING ERRORS IN FINITE DIFFERENCES

$$\left. \begin{aligned} \hat{f}(x+h) &= f(x+h) + \epsilon_1 \\ \hat{f}(x-h) &= f(x-h) + \epsilon_2 \end{aligned} \right\} |\epsilon_1|, |\epsilon_2| \approx \epsilon_{\text{mach}}, \text{ rounding errors}$$

$$\begin{aligned} f'_{\text{correct}}(x) - f'_{\text{machine}}(x) &= f'(x) - \frac{\hat{f}(x+h) - \hat{f}(x-h)}{2h} \\ &= \underbrace{f'(x)}_{f'_{\text{correct}}(x)} - \underbrace{\frac{f(x+h) - f(x-h)}{2h}}_{f'_{\text{formula}}(x)} + \underbrace{\frac{\epsilon_2 - \epsilon_1}{2h}}_{\text{rounding error}} \end{aligned}$$

ROUNDING ERRORS IN FINITE DIFFERENCES

$$\left. \begin{aligned} \hat{f}(x+h) &= f(x+h) + \epsilon_1 \\ \hat{f}(x-h) &= f(x-h) + \epsilon_2 \end{aligned} \right\} |\epsilon_1|, |\epsilon_2| \approx \epsilon_{\text{mach}}, \text{ rounding errors}$$

$$\begin{aligned} f'_{\text{correct}}(x) - f'_{\text{machine}}(x) &= f'(x) - \frac{\hat{f}(x+h) - \hat{f}(x-h)}{2h} \\ &= \underbrace{f'(x)}_{f'_{\text{correct}}(x)} - \underbrace{\frac{f(x+h) - f(x-h)}{2h}}_{f'_{\text{formula}}(x)} + \underbrace{\frac{\epsilon_2 - \epsilon_1}{2h}}_{\text{rounding error}} \end{aligned}$$

As h gets smaller, error in the finite difference formulas becomes smaller, but the rounding error grows. There is an optimal h where the sum of the two error is at a minimum.

NUMERICAL INTEGRATION

NEWTON-COTES FORMULAS FOR NUMERICAL INTEGRATION

- Given a complex function $f(x)$, sample the function at n points: $(x_1, y_1), \dots, (x_n, y_n)$.

NEWTON-COTES FORMULAS FOR NUMERICAL INTEGRATION

- Given a complex function $f(x)$, sample the function at n points: $(x_1, y_1), \dots, (x_n, y_n)$.
- Fit a low degree polynomial to data points to get $P(x)$.


NEWTON-COTES FORMULAS FOR NUMERICAL INTEGRATION

- Given a complex function $f(x)$, sample the function at n points: $(x_1, y_1), \dots, (x_n, y_n)$.
- Fit a low degree polynomial to data points to get $P(x)$.
- Approximate the true integral $\int_a^b f(x) dx$ by $\int_a^b P(x) dx$.

NEWTON-COTES FORMULAS FOR NUMERICAL INTEGRATION


- Given a complex function $f(x)$, sample the function at n points: $(x_1, y_1), \dots, (x_n, y_n)$.
- Fit a low degree polynomial to data points to get $P(x)$.
- Approximate the true integral $\int_a^b f(x) dx$ by $\int_a^b P(x) dx$.
- Compute $\int_a^b P(x) dx$ by the true analytical expression since it is easy to integrate polynomials.

TRAPEZOID RULE ($n = 2$)



$$f(x) \approx y_0 \frac{x - x_1}{x_0 - x_1} + y_1 \frac{x - x_0}{x_1 - x_0}$$


TRAPEZOID RULE ($n = 2$)



$$f(x) \approx y_0 \frac{x - x_1}{x_0 - x_1} + y_1 \frac{x - x_0}{x_1 - x_0}$$

$$\int_{x_0}^{x_1} f(x) dx \approx y_0 \int_{x_0}^{x_1} \frac{x - x_1}{x_0 - x_1} dx + y_1 \int_{x_0}^{x_1} \frac{x - x_0}{x_1 - x_0} dx$$


TRAPEZOID RULE ($n = 2$)



$$f(x) \approx y_0 \frac{x - x_1}{x_0 - x_1} + y_1 \frac{x - x_0}{x_1 - x_0}$$

$$\begin{aligned} \int_{x_0}^{x_1} f(x) dx &\approx y_0 \int_{x_0}^{x_1} \frac{x - x_1}{x_0 - x_1} dx + y_1 \int_{x_0}^{x_1} \frac{x - x_0}{x_1 - x_0} dx \\ &\approx y_0 \int_0^{x_1 - x_0} \frac{u + x_0 - x_1}{x_0 - x_1} du + y_1 \int_{x_0 - x_1}^0 \frac{v + x_1 - x_0}{x_1 - x_0} dv \quad \begin{pmatrix} u = x - x_0, \\ v = x - x_1 \end{pmatrix} \end{aligned}$$


TRAPEZOID RULE ($n = 2$)



$$f(x) \approx y_0 \frac{x - x_1}{x_0 - x_1} + y_1 \frac{x - x_0}{x_1 - x_0}$$

$$\begin{aligned} \int_{x_0}^{x_1} f(x) dx &\approx y_0 \int_{x_0}^{x_1} \frac{x - x_1}{x_0 - x_1} dx + y_1 \int_{x_0}^{x_1} \frac{x - x_0}{x_1 - x_0} dx \\ &\approx y_0 \int_0^{x_1 - x_0} \frac{u + x_0 - x_1}{x_0 - x_1} du + y_1 \int_{x_0 - x_1}^0 \frac{v + x_1 - x_0}{x_1 - x_0} dv \quad \begin{pmatrix} u = x - x_0, \\ v = x - x_1 \end{pmatrix} \\ &\approx y_0 \int_0^h \frac{u - h}{-h} du + y_1 \int_{-h}^0 \frac{v + h}{h} dv \quad (h = x_1 - x_0) \end{aligned}$$


TRAPEZOID RULE ($n = 2$)



$$f(x) \approx y_0 \frac{x - x_1}{x_0 - x_1} + y_1 \frac{x - x_0}{x_1 - x_0}$$


$$\begin{aligned} \int_{x_0}^{x_1} f(x) dx &\approx y_0 \int_{x_0}^{x_1} \frac{x - x_1}{x_0 - x_1} dx + y_1 \int_{x_0}^{x_1} \frac{x - x_0}{x_1 - x_0} dx \\ &\approx y_0 \int_0^{x_1 - x_0} \frac{u + x_0 - x_1}{x_0 - x_1} du + y_1 \int_{x_0 - x_1}^0 \frac{v + x_1 - x_0}{x_1 - x_0} dv \quad \begin{pmatrix} u = x - x_0, \\ v = x - x_1 \end{pmatrix} \\ &\approx y_0 \int_0^h \frac{u - h}{-h} du + y_1 \int_{-h}^0 \frac{v + h}{h} dv \quad (h = x_1 - x_0) \\ &\approx -y_0 \left. \frac{\frac{u^2}{2} - hu}{h} \right|_0^h + y_1 \left. \frac{\frac{v^2}{2} + hv}{h} \right|_{-h}^0 = y_0 \frac{h}{2} + y_1 \frac{h}{2} = h \frac{(y_0 + y_1)}{2}. \end{aligned}$$

SIMPSON'S RULE ($n = 3$)



$$f(x) \approx y_0 \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}$$


SIMPSON'S RULE ($n = 3$)



$$f(x) \approx y_0 \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}$$

$$\int_{x_0}^{x_2} f(x) dx \approx y_0 \int_{x_0}^{x_2} \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} dx + y_1 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} dx + y_2 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} dx$$


SIMPSON'S RULE ($n = 3$)



$$f(x) \approx y_0 \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}$$

$$\begin{aligned} \int_{x_0}^{x_2} f(x) dx &\approx y_0 \int_{x_0}^{x_2} \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} dx + y_1 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} dx \\ &\quad + y_2 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} dx \\ &= y_0 \frac{h}{3} + y_1 \frac{4h}{3} + y_2 \frac{h}{3} \quad (h = x_2 - x_1 = x_1 - x_0) \end{aligned}$$

SIMPSON'S RULE ($n = 3$)



$$f(x) \approx y_0 \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}$$

$$\begin{aligned} \int_{x_0}^{x_2} f(x) dx &\approx y_0 \int_{x_0}^{x_2} \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} dx + y_1 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} dx \\ &\quad + y_2 \int_{x_0}^{x_2} \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} dx \\ &= y_0 \frac{h}{3} + y_1 \frac{4h}{3} + y_2 \frac{h}{3} \quad (h = x_2 - x_1 = x_1 - x_0) \\ &= \frac{h}{3} (y_0 + 4y_1 + y_2). \end{aligned}$$

EXAMPLE

Apply Trapezoid and Simpson's rules to $\int_1^2 \ln x \, dx$.

EXAMPLE

Apply Trapezoid and Simpson's rules to $\int_1^2 \ln x \, dx$.

Exact value:

$$\int_1^2 \ln x \, dx = x \ln x \Big|_1^2 - \int_1^2 dx = 2 \ln 2 - \ln 1 - 1 \approx 0.3863$$

EXAMPLE

Apply Trapezoid and Simpson's rules to $\int_1^2 \ln x \, dx$.

Exact value:

$$\int_1^2 \ln x \, dx = x \ln x \Big|_1^2 - \int_1^2 dx = 2 \ln 2 - \ln 1 - 1 \approx 0.3863$$

Trapezoid R.:

$$\int_1^2 \ln x \, dx \approx \frac{h}{2}(y_0 + y_1) = \frac{1}{2}(\ln 1 + \ln 2) = \frac{\ln 2}{2} \approx 0.3466$$

EXAMPLE

Apply Trapezoid and Simpson's rules to $\int_1^2 \ln x \, dx$.

Exact value:

$$\int_1^2 \ln x \, dx = x \ln x \Big|_1^2 - \int_1^2 dx = 2 \ln 2 - \ln 1 = 2 \ln 2 - 0 \approx 0.3863$$

Trapezoid R.:


$$\int_1^2 \ln x \, dx \approx \frac{h}{2}(y_0 + y_1) = \frac{1}{2}(\ln 1 + \ln 2) = \frac{\ln 2}{2} \approx 0.3466$$

Simpson's R.:

$$\int_1^2 \ln x \, dx \approx \frac{h}{3}(y_0 + 4y_1 + y_2) = \frac{0.5}{3}(\ln 1 + 4 \ln \frac{3}{2} + \ln 2) \approx 0.3858$$

COMPOSITE NEWTON-COTES FORMULAS

Composite Trapezoid Rule




$$\int_a^b f(x) dx \approx \frac{h}{2} \left[f(a) + f(b) + 2 \sum_{i=1}^{m-1} f(x_i) \right],$$

$$m = 4, a = x_0, b = x_4, h = \frac{b - a}{m}.$$

COMPOSITE NEWTON-COTES FORMULAS


Composite Trapezoid Rule



$$\int_a^b f(x) dx \approx \frac{h}{2} \left[f(a) + f(b) + 2 \sum_{i=1}^{m-1} f(x_i) \right],$$

$$m = 4, a = x_0, b = x_4, h = \frac{b - a}{m}.$$

Composite Simpson's Rule



$$\int_a^b f(x) dx \approx \frac{h}{3} \left[f(a) + f(b) + 4 \sum_{i=1}^m f(x_{2i-1}) + 2 \sum_{i=1}^{m-1} f(x_{2i}) \right],$$

$$m = 2, a = x_0, b = x_4, h = \frac{b - a}{2m}.$$