

CENG 216 – NUMERICAL COMPUTATION

INTERPOLATION - PART I

Mustafa Özuysal
mustafaozuysal@iyte.edu.tr

May 9, 2022

İzmir Institute of Technology


Slides are based on material from the main textbook:
“Numerical Analysis”, The new international edition, 2ed,
by Timothy Sauer

DATA AND INTERPOLATING FUNCTIONS

INTERPOLATION

A function is said to interpolate a set of data points if it passes through those points.


Def: The function $y = P(x)$ interpolates the data points $(x_1, y_1), \dots, (x_n, y_n)$ if $P(x_i) = y_i \ \forall i = 1, \dots, n$.



INTERPOLATION

A function is said to interpolate a set of data points if it passes through those points.

Def: The function $y = P(x)$ interpolates the data points $(x_1, y_1), \dots, (x_n, y_n)$ if $P(x_i) = y_i \ \forall i = 1, \dots, n$.



LAGRANGE INTERPOLATION

To interpolate the n points $(x_1, y_1), \dots, (x_n, y_n)$, we define the $n - 1$ degree polynomials $L_k(x) \forall k, k = 1, \dots, n$:

$$L_k(x) = \frac{(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}.$$

LAGRANGE INTERPOLATION

To interpolate the n points $(x_1, y_1), \dots, (x_n, y_n)$, we define the $n - 1$ degree polynomials $L_k(x) \forall k, k = 1, \dots, n$:

$$L_k(x) = \frac{(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}.$$

Note that

$$L_k(x_k) = 1 \quad \text{and} \quad L_k(x_j) = 0 \quad \text{for } j \neq k.$$

LAGRANGE INTERPOLATION

To interpolate the n points $(x_1, y_1), \dots, (x_n, y_n)$, we define the $n - 1$ degree polynomials $L_k(x) \forall k, k = 1, \dots, n$:

$$L_k(x) = \frac{(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}.$$

Note that

$$L_k(x_k) = 1 \text{ and } L_k(x_j) = 0 \text{ for } j \neq k.$$

LAGRANGE INTERPOLATION

To interpolate the n points $(x_1, y_1), \dots, (x_n, y_n)$, we define the $n - 1$ degree polynomials $L_k(x) \forall k, k = 1, \dots, n$:

$$L_k(x) = \frac{(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}.$$

Note that

$$L_k(x_k) = 1 \text{ and } L_k(x_j) = 0 \text{ for } j \neq k.$$

Then the **interpolating polynomial** of degree $n - 1$ for these data points is

$$P_{n-1}(x) = y_1 L_1(x) + y_2 L_2(x) + \dots + y_n L_n(x),$$

LAGRANGE INTERPOLATION

To interpolate the n points $(x_1, y_1), \dots, (x_n, y_n)$, we define the $n - 1$ degree polynomials $L_k(x) \forall k, k = 1, \dots, n$:

$$L_k(x) = \frac{(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_1) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}.$$

Note that

$$L_k(x_k) = 1 \text{ and } L_k(x_j) = 0 \text{ for } j \neq k.$$

Then the **interpolating polynomial** of degree $n - 1$ for these data points is

$$P_{n-1}(x) = y_1 L_1(x) + y_2 L_2(x) + \dots + y_n L_n(x),$$

since

$$P_{n-1}(x_k) = y_1 \cdot 0 + y_2 \cdot 0 + \dots + y_k \cdot 1 + \dots + y_n \cdot 0 = y_k \quad \forall k = 1, \dots, n.$$

EXAMPLE

Interpolate points $(0, 1)$, $(2, 2)$, and $(3, 4)$.

EXAMPLE

Interpolate points $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$P_2(x) = 1 \frac{(x-2)(x-3)}{(0-2)(0-3)} + 2 \frac{(x-0)(x-3)}{(2-0)(2-3)} + 4 \frac{(x-0)(x-2)}{(3-0)(3-2)}$$

EXAMPLE

Interpolate points $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{aligned} P_2(x) &= 1 \frac{(x-2)(x-3)}{(0-2)(0-3)} + 2 \frac{(x-0)(x-3)}{(2-0)(2-3)} + 4 \frac{(x-0)(x-2)}{(3-0)(3-2)} \\ &= \frac{1}{6}(x-2)(x-3) - x(x-3) + \frac{4}{3}x(x-2) \end{aligned}$$

EXAMPLE

Interpolate points $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{aligned}P_2(x) &= 1 \frac{(x-2)(x-3)}{(0-2)(0-3)} + 2 \frac{(x-0)(x-3)}{(2-0)(2-3)} + 4 \frac{(x-0)(x-2)}{(3-0)(3-2)} \\&= \frac{1}{6}(x-2)(x-3) - x(x-3) + \frac{4}{3}x(x-2) \\&= \frac{1}{2}x^2 - \frac{1}{2}x + 1\end{aligned}$$

EXAMPLE

Interpolate points $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{aligned}P_2(x) &= 1 \frac{(x-2)(x-3)}{(0-2)(0-3)} + 2 \frac{(x-0)(x-3)}{(2-0)(2-3)} + 4 \frac{(x-0)(x-2)}{(3-0)(3-2)} \\&= \frac{1}{6}(x-2)(x-3) - x(x-3) + \frac{4}{3}x(x-2) \\&= \frac{1}{2}x^2 - \frac{1}{2}x + 1\end{aligned}$$

We can check the interpolation property:

$$P_2(0) = 1, P_2(2) = 2, \text{ and } P_2(3) = 4.$$

EXISTANCE AND UNIQUENESS

Thm: Main Theorem of Polynomial Interpolation

Let $(x_1, y_1), \dots, (x_n, y_n)$ be n points in the plane with distinct x_i . Then there exists **one and only one** polynomial of degree $n - 1$ or less that satisfies $P(x_i) = y_i \ \forall i = 1, \dots, n$.

EXISTANCE AND UNIQUENESS

Thm: Main Theorem of Polynomial Interpolation

Let $(x_1, y_1), \dots, (x_n, y_n)$ be n points in the plane with distinct x_i . Then there exists **one and only one** polynomial of degree $n - 1$ or less that satisfies $P(x_i) = y_i \ \forall i = 1, \dots, n$.

Existence Proof: Lagrange interpolation formula.

EXISTANCE AND UNIQUENESS

Thm: Main Theorem of Polynomial Interpolation

Let $(x_1, y_1), \dots, (x_n, y_n)$ be n points in the plane with distinct x_i . Then there exists **one and only one** polynomial of degree $n - 1$ or less that satisfies $P(x_i) = y_i \ \forall i = 1, \dots, n$.

Existence Proof: Lagrange interpolation formula.

Uniqueness Proof: Let $P(x)$ and $Q(x)$ be distinct $n - 1$ degree polynomials that interpolate the n points.

EXISTANCE AND UNIQUENESS

Thm: Main Theorem of Polynomial Interpolation

Let $(x_1, y_1), \dots, (x_n, y_n)$ be n points in the plane with distinct x_i . Then there exists **one and only one** polynomial of degree $n - 1$ or less that satisfies $P(x_i) = y_i \ \forall i = 1, \dots, n$.

Existence Proof: Lagrange interpolation formula.

Uniqueness Proof: Let $P(x)$ and $Q(x)$ be distinct $n - 1$ degree polynomials that interpolate the n points. $H(x) = P(x) - Q(x)$ is of degree $n - 1$ and $H(x_i) = 0 \ \forall i = 1, \dots, n \rightarrow H(x)$ has n distinct zeros.

EXISTANCE AND UNIQUENESS

Thm: Main Theorem of Polynomial Interpolation

Let $(x_1, y_1), \dots, (x_n, y_n)$ be n points in the plane with distinct x_i . Then there exists **one and only one** polynomial of degree $n - 1$ or less that satisfies $P(x_i) = y_i \ \forall i = 1, \dots, n$.

Existence Proof: Lagrange interpolation formula.

Uniqueness Proof: Let $P(x)$ and $Q(x)$ be distinct $n - 1$ degree polynomials that interpolate the n points. $H(x) = P(x) - Q(x)$ is of degree $n - 1$ and $H(x_i) = 0 \ \forall i = 1, \dots, n \rightarrow H(x)$ has n distinct zeros. **Fundamental Theorem of Algebra** states that polynomial of degree d has at most d zeros or its is the zero polynomial $\rightarrow H(x) = 0 = P(x) - Q(x) \rightarrow P(x) = Q(x)$ ✖

EXAMPLE

Find polynomial of degree three or less interpolating $(0, 2)$, $(1, 1)$, $(2, 0)$ and $(3, -1)$.

$$\begin{aligned} P(x) = & 2 \frac{(x-1)(x-2)(x-3)}{(0-1)(0-2)(0-3)} + 1 \frac{x(x-2)(x-3)}{(1-0)(1-2)(1-3)} \\ & + 0 \frac{x(x-1)(x-3)}{(2-0)(2-1)(2-3)} - 1 \frac{x(x-1)(x-2)}{(3-0)(3-1)(3-2)} \end{aligned}$$

EXAMPLE

Find polynomial of degree three or less interpolating $(0, 2)$, $(1, 1)$, $(2, 0)$ and $(3, -1)$.

$$\begin{aligned}P(x) &= 2 \frac{(x-1)(x-2)(x-3)}{(0-1)(0-2)(0-3)} + 1 \frac{x(x-2)(x-3)}{(1-0)(1-2)(1-3)} \\&\quad + 0 \frac{x(x-1)(x-3)}{(2-0)(2-1)(2-3)} - 1 \frac{x(x-1)(x-2)}{(3-0)(3-1)(3-2)} \\&= -x + 2,\end{aligned}$$

EXAMPLE

Find polynomial of degree three or less interpolating $(0, 2)$, $(1, 1)$, $(2, 0)$ and $(3, -1)$.

$$\begin{aligned} P(x) &= 2 \frac{(x-1)(x-2)(x-3)}{(0-1)(0-2)(0-3)} + 1 \frac{x(x-2)(x-3)}{(1-0)(1-2)(1-3)} \\ &\quad + 0 \frac{x(x-1)(x-3)}{(2-0)(2-1)(2-3)} - 1 \frac{x(x-1)(x-2)}{(3-0)(3-1)(3-2)} \\ &= -x + 2, \end{aligned}$$

which is a line and the points are colinear. There is no parabola (degree=2) and no cubic (degree=3) passing through the same points.

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots \\ f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots \\ f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots$$
$$f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$
$$f[x_k] = f(x_k)$$

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots \\ f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$

$$f[x_k] = f(x_k)$$

$$f[x_k x_{k+1}] = \frac{f[x_{k+1}] - f[x_k]}{x_{k+1} - x_k}$$

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots \\ f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$

$$f[x_k] = f(x_k)$$

$$f[x_k x_{k+1}] = \frac{f[x_{k+1}] - f[x_k]}{x_{k+1} - x_k}$$

$$f[x_k x_{k+1} x_{k+2}] = \frac{f[x_{k+1} x_{k+2}] - f[x_k x_{k+1}]}{x_{k+2} - x_k}$$

NEWTON'S DIVIDED DIFFERENCES

Simpler way of writing the interpolating polynomial.

Def: Denote by $f[x_1 \dots x_n]$, the coefficient of the x^{n-1} term in the unique polynomial that interpolates $(x_1, f(x_1)), \dots, (x_n, f(x_n))$.

$$P(x) = f[x_1] + f[x_1 x_2](x - x_1) + f[x_1 x_2 x_3](x - x_1)(x - x_2) + \dots \\ f[x_1 \dots x_n](x - x_1) \dots (x - x_{n-1})$$

$$f[x_k] = f(x_k)$$

$$f[x_k x_{k+1}] = \frac{f[x_{k+1}] - f[x_k]}{x_{k+1} - x_k}$$

$$f[x_k x_{k+1} x_{k+2}] = \frac{f[x_{k+1} x_{k+2}] - f[x_k x_{k+1}]}{x_{k+2} - x_k}$$

$$f[x_k x_{k+1} x_{k+2} x_{k+3}] = \frac{f[x_{k+1} x_{k+2} x_{k+3}] - f[x_k x_{k+1} x_{k+2}]}{x_{k+3} - x_k}$$

$$\dots = \dots$$

NEWTON'S DIVIDED DIFFERENCES: THE NOTATION

Interpolate $(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3))$.

$$\begin{array}{l|l} x_1 & f[x_1] = f(x_1) \\ x_2 & f[x_2] = f(x_2) \\ x_3 & f[x_3] = f(x_3) \end{array}$$

NEWTON'S DIVIDED DIFFERENCES: THE NOTATION

Interpolate $(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3))$.

$$\begin{array}{l|l} x_1 & f[x_1] = f(x_1) \\ & f[x_1, x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1} \\ x_2 & f[x_2] = f(x_2) \\ & \\ x_3 & f[x_3] = f(x_3) \end{array}$$

NEWTON'S DIVIDED DIFFERENCES: THE NOTATION

Interpolate $(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3))$.

$$\begin{array}{l|l} x_1 & f[x_1] = f(x_1) \\ & f[x_1 \ x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1} \\ x_2 & f[x_2] = f(x_2) \\ & f[x_2 \ x_3] = \frac{f[x_3] - f[x_2]}{x_3 - x_2} \\ x_3 & f[x_3] = f(x_3) \end{array}$$

NEWTON'S DIVIDED DIFFERENCES: THE NOTATION

Interpolate $(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3))$.

$$\begin{array}{l|l} x_1 & f[x_1] = f(x_1) \\ x_2 & f[x_2] = f(x_2) \\ x_3 & f[x_3] = f(x_3) \end{array} \quad \begin{array}{l} f[x_1 \ x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1} \\ f[x_2 \ x_3] = \frac{f[x_3] - f[x_2]}{x_3 - x_2} \end{array} \quad f[x_1 \ x_2 \ x_3] = \frac{f[x_2 \ x_3] - f[x_1 \ x_2]}{x_3 - x_1}$$

NEWTON'S DIVIDED DIFFERENCES: THE NOTATION

Interpolate $(x_1, f(x_1)), (x_2, f(x_2)), (x_3, f(x_3))$.

$$\begin{array}{l|l} x_1 & f[x_1] = f(x_1) \\ & f[x_1 \ x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1} \\ x_2 & f[x_2] = f(x_2) \\ & f[x_2 \ x_3] = \frac{f[x_3] - f[x_2]}{x_3 - x_2} \\ x_3 & f[x_3] = f(x_3) \end{array} \quad f[x_1 \ x_2 \ x_3] = \frac{f[x_2 \ x_3] - f[x_1 \ x_2]}{x_3 - x_1}$$

And the interpolating polynomial is

$$P_2(x) = f[x_1] + f[x_1 \ x_2](x - x_1) + f[x_1 \ x_2 \ x_3](x - x_1)(x - x_2)$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

0	1
2	2
3	4

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{2-1}{2-0} = \frac{1}{2} \\ 2 & 2 \\ & \\ 3 & 4 \end{array}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{2-1}{2-0} = \frac{1}{2} \\ 2 & 2 \\ & \frac{4-2}{3-2} = \frac{2}{1} = 2 \\ 3 & 4 \end{array}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{2-1}{2-0} = \frac{1}{2} \\ 2 & 2 \\ & \frac{4-2}{3-2} = \frac{2}{1} = 2 \\ 3 & 4 \end{array} \quad \frac{2-\frac{1}{2}}{3-0} = \frac{1}{2}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{2-1}{2-0} = \frac{1}{2} \\ 2 & 2 \\ & \frac{4-2}{3-2} = \frac{2}{1} = 2 \\ 3 & 4 \end{array} \quad \frac{2-\frac{1}{2}}{3-0} = \frac{1}{2}$$

And the interpolating polynomial is

$$P_2(x) = 1 + \frac{1}{2}(x - 0) + \frac{1}{2}(x - 0)(x - 2)$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, and $(3, 4)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{2-1}{2-0} = \frac{1}{2} \\ 2 & 2 \\ & \frac{4-2}{3-2} = \frac{2}{1} = 2 \\ 3 & 4 \end{array} \qquad \frac{2-\frac{1}{2}}{3-0} = \frac{1}{2}$$

And the interpolating polynomial is

$$\begin{aligned} P_2(x) &= 1 + \frac{1}{2}(x-0) + \frac{1}{2}(x-0)(x-2) \\ &= 1 + \frac{1}{2}x^2 - \frac{2}{2}x \\ &= \frac{1}{2}x^2 - \frac{1}{2}x + 1 \end{aligned}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

0	1
2	2
3	4
1	0

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

0	1		
		$\frac{1}{2}$	
2	2		$\frac{1}{2}$
		2	
3	4		
1	0		

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

0	1		
		$\frac{1}{2}$	
2	2		$\frac{1}{2}$
		2	
3	4		
		$\frac{0-4}{1-3} = 2$	
1	0		

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

$$\begin{array}{r|rrrr} 0 & 1 & & & \\ & & \frac{1}{2} & & \\ 2 & 2 & & \frac{1}{2} & \\ & & 2 & & \\ 3 & 4 & & \frac{2-2}{1-2} = 0 & \\ & & \frac{0-4}{1-3} = 2 & & \\ 1 & 0 & & & \end{array}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

$$\begin{array}{c|c} 0 & 1 \\ & \frac{1}{2} \\ 2 & 2 \\ & 2 \\ & \frac{1}{2} \\ 3 & 4 \\ & \frac{2-2}{1-2} = 0 \\ & \frac{0-4}{1-3} = 2 \\ 1 & 0 \\ & \frac{0-\frac{1}{2}}{1-0} = -\frac{1}{2} \end{array}$$

EXAMPLE

Use divided differences to find the interpolating polynomial passing through $(0, 1)$, $(2, 2)$, $(3, 4)$, and $(1, 0)$.

0	1				
		$\frac{1}{2}$			
2	2		$\frac{1}{2}$		
		2		$\frac{0-\frac{1}{2}}{1-0} = -\frac{1}{2}$	
3	4		$\frac{2-2}{1-2} = 0$		
		$\frac{0-4}{1-3} = 2$			
1	0				

And the interpolating polynomial is

$$P_2(x) = 1 + \underbrace{\frac{1}{2}(x-0) + \frac{1}{2}(x-0)(x-2) - \frac{1}{2}(x-0)(x-2)(x-3)}_{\text{same as before}}$$

APPROXIMATING FUNCTIONS BY POLYNOMIALS

Find an approximating polynomial to $f(x) = \sin(x)$ by interpolating at four equally spaced points on $[0, \frac{\pi}{2}]$.

APPROXIMATING FUNCTIONS BY POLYNOMIALS

Find an approximating polynomial to $f(x) = \sin(x)$ by interpolating at four equally spaced points on $[0, \frac{\pi}{2}]$.

0	0.0000
$\frac{\pi}{6}$	0.5000
$\frac{2\pi}{6}$	0.8660
$\frac{3\pi}{6}$	1.0000

APPROXIMATING FUNCTIONS BY POLYNOMIALS

Find an approximating polynomial to $f(x) = \sin(x)$ by interpolating at four equally spaced points on $[0, \frac{\pi}{2}]$.

0	0.0000			
		0.9549		
$\frac{\pi}{6}$	0.5000		− 0.2443	
		0.6990		− 0.1139
$\frac{2\pi}{6}$	0.8660		− 0.4232	
		0.2559		
$\frac{3\pi}{6}$	1.0000			

APPROXIMATING FUNCTIONS BY POLYNOMIALS

Find an approximating polynomial to $f(x) = \sin(x)$ by interpolating at four equally spaced points on $[0, \frac{\pi}{2}]$.

0	0.0000			
		0.9549		
$\frac{\pi}{6}$	0.5000		- 0.2443	
		0.6990		- 0.1139
$\frac{2\pi}{6}$	0.8660		- 0.4232	
		0.2559		
$\frac{3\pi}{6}$	1.0000			

$$P(x) = 0.0000 + 0.9545x - 0.2443x(x - \frac{\pi}{6}) - 0.1139x(x - \frac{\pi}{6})(x - \frac{2\pi}{6})$$

APPROXIMATING FUNCTIONS BY POLYNOMIALS

$$P(x) = 0.0000 + 0.9545x - 0.2443x\left(x - \frac{\pi}{6}\right) - 0.1139x\left(x - \frac{\pi}{6}\right)\left(x - \frac{2\pi}{6}\right) \\ \approx \sin(x)$$

APPROXIMATING FUNCTIONS BY POLYNOMIALS

$$P(x) = 0.0000 + 0.9545x - 0.2443x\left(x - \frac{\pi}{6}\right) - 0.1139x\left(x - \frac{\pi}{6}\right)\left(x - \frac{2\pi}{6}\right) \\ \approx \sin(x)$$

$[0, \frac{\pi}{2}]$ is called the **fundamental domain**. It is sufficient for approximating $\sin(x)$ since if $x \in [\frac{\pi}{2}, \pi] \rightarrow \sin(x) = \sin(\pi - x)$ and if $x \in [\pi, 2\pi] \rightarrow \sin(x) = -\sin(2\pi - x)$.

APPROXIMATING FUNCTIONS BY POLYNOMIALS

$$P(x) = 0.0000 + 0.9545x - 0.2443x\left(x - \frac{\pi}{6}\right) - 0.1139x\left(x - \frac{\pi}{6}\right)\left(x - \frac{2\pi}{6}\right) \\ \approx \sin(x)$$

$[0, \frac{\pi}{2}]$ is called the **fundamental domain**. It is sufficient for approximating $\sin(x)$ since if $x \in [\frac{\pi}{2}, \pi] \rightarrow \sin(x) = \sin(\pi - x)$ and if $x \in [\pi, 2\pi] \rightarrow \sin(x) = -\sin(2\pi - x)$.

x	$\sin(x)$	$P(x)$	$ \sin(x) - P(x) $
1	0.8415	0.8411	0.0004
2	0.9093	0.9192	0.0009
3	0.1411	0.1428	0.0017
4	-0.7568	-0.7557	0.0011
14	0.9906	0.9928	0.0022
1000	0.8269	0.8263	0.0006

INTERPOLATION ERROR

BOUNDING THE INTERPOLATION ERROR

Thm: Assume that $P(x)$ is the (degree $n - 1$ or less) interpolating polynomial fitting the n points $(x_1, y_1), \dots, (x_n, y_n)$ sampled from a function $f(x)$.

BOUNDING THE INTERPOLATION ERROR

Thm: Assume that $P(x)$ is the (degree $n - 1$ or less) interpolating polynomial fitting the n points $(x_1, y_1), \dots, (x_n, y_n)$ sampled from a function $f(x)$. The interpolation error is

$$f(x) - P(x) = \frac{(x - x_1)(x - x_2) \dots (x - x_n)}{n!} f^{(n)}(c),$$

where $f^{(n)}(x)$ is the n^{th} derivative of $f(x)$ and c lies between the largest and the smallest of x_1, x_2, \dots, x_n .

EXAMPLE

$$\begin{aligned}\sin(x) - P(x) &= \frac{(x - 0)(x - \frac{\pi}{6})(x - \frac{\pi}{3})(x - \frac{\pi}{2})}{4!} f^{(4)}(c) \\ &= \frac{(x - 0)(x - \frac{\pi}{6})(x - \frac{\pi}{3})(x - \frac{\pi}{2})}{4!} \sin(c)\end{aligned}$$

EXAMPLE

$$\begin{aligned}\sin(x) - P(x) &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} f^{(4)}(c) \\ &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} \sin(c) \\ |\sin(x) - P(x)| &\leq \frac{|(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})|}{24} |1|\end{aligned}$$

EXAMPLE

$$\begin{aligned}\sin(x) - P(x) &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} f^{(4)}(c) \\ &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} \sin(c) \\ |\sin(x) - P(x)| &\leq \frac{|(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})|}{24} |1|\end{aligned}$$

- At $x = 1$, the worst case error is ≈ 0.0005348 (> 0.0004 from the table of actual errors that we obtained).

EXAMPLE

$$\begin{aligned}\sin(x) - P(x) &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} f^{(4)}(c) \\ &= \frac{(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})}{4!} \sin(c) \\ |\sin(x) - P(x)| &\leq \frac{|(x-0)(x-\frac{\pi}{6})(x-\frac{\pi}{3})(x-\frac{\pi}{2})|}{24} |1|\end{aligned}$$

- At $x = 1$, the worst case error is ≈ 0.0005348 (> 0.0004 from the table of actual errors that we obtained).
- Error will be smaller in the middle of the interval due to smaller terms in the numerator.

RUNGE PHENOMENON

Try to interpolate with data points

$$x \in \{-3, -2.5, -2.0, \dots, 2.0, 2.5, 3.0\}$$


with y equal to zeros except at $x = 0, y = 1$.

RUNGE PHENOMENON

Try to interpolate with data points

$$x \in \{-3, -2.5, -2.0, \dots, 2.0, 2.5, 3.0\}$$

with y equal to zeros except at $x = 0, y = 1$.




RUNGE PHENOMENON

Try to interpolate with data points

$$x \in \{-3, -2.5, -2.0, \dots, 2.0, 2.5, 3.0\}$$

with y equal to zeros except at $x = 0, y = 1$.




Very large errors near
the ends of the interval!

RUNGE PHENOMENON

Try to interpolate with data points

$$x \in \{-3, -2.5, -2.0, \dots, 2.0, 2.5, 3.0\}$$

with y equal to zeros except at $x = 0, y = 1$.



Very large errors near the ends of the interval!

→ Place more points closer to the end points instead of uniformly sampling x → **Chebyshev Interpolation** (which we will skip for the course. See §3.3 for more.)