# CENG 216 - NUMERICAL COMPUTATION

#### Introduction

Mustafa Özuysal mustafaozuysal@iyte.edu.tr February 16, 2022

İzmir Institute of Technology

**COURSE INFORMATION** 

#### **PEOPLE**

Assoc. Prof. Mustafa Özuysal Furkan Eren Uzyıldırım Ersin Çine mustafaozuysal@iyte.edu.tr furkanuzyildirim@iyte.edu.tr ersincine@iyte.edu.tr

Time Monday, 09:45–12:30

Teams Code e65du56

1

# GRADING

Assignment	Grade Percentage
Homeworks	40%
Midterm Exam	30%
Final Exam	30%

#### **TEXTBOOKS**

#### Main Textbook:

Numerical Analysis, The new international edition, 2ed, Timothy Sauer

## **Secondary Textbooks:**

Numerical Algorithms: Methods for Computer Vision, Machine Learning, and Graphics, J. Solomon

#### https


//people.csail.mit.edu/jsolomon/share/book/numerical\_book.pdf Matrix Computations (4th Ed.), G. H. Golub and C. F. Van Loan

#### COURSE CONTENTS


This course covers the fundamentals of numerical computation starting with finite representation of real numbers and investigation of errors resulting from the discrete and approximate nature of the computation using such a representation.

The course includes topics from numerical linear algebra, interpolation, numerical differentiation and integration, numerical solution of ordinary differential equations, basics of numerical optimization, and generation of random numbers with their application to numerical problems.


# **NUMERICAL COMPUTATION**



# **NUMERICAL COMPUTATION**



# NUMERICAL COMPUTATION



Introduction

- Introduction
- · Fundamentals (Chapter o)

- Introduction
- · Fundamentals (Chapter o)
- · Solving Equations (Chapter 1)

- Introduction
- · Fundamentals (Chapter o)
- · Solving Equations (Chapter 1)
- Systems of Equations (Chapter 2)

- Introduction
- · Fundamentals (Chapter o)
- · Solving Equations (Chapter 1)
- Systems of Equations (Chapter 2)
- Interpolation (Chapter 3)

- Introduction
- · Fundamentals (Chapter o)
- Solving Equations (Chapter 1)
- Systems of Equations (Chapter 2)
- Interpolation (Chapter 3)
- · Least Squares (Chapter 4)

- Introduction
- Fundamentals (Chapter o)
- Solving Equations (Chapter 1)
- Systems of Equations (Chapter 2)
- Interpolation (Chapter 3)
- Least Squares (Chapter 4)
- · Numerical Differentiation and Integration (Chapter 5)

- Introduction
- · Fundamentals (Chapter o)
- · Solving Equations (Chapter 1)
- Systems of Equations (Chapter 2)
- Interpolation (Chapter 3)
- Least Squares (Chapter 4)
- Numerical Differentiation and Integration (Chapter 5)
- Ordinary Differential Equations (Chapter 6)

#### LEARNING OUTCOMES I

- To be able to explain the effects of the finite representation of real numbers on the implementation of a given algorithm.
- To be able to derive the numerical error in computations and compare the numerical error of different algorithms for the same problem.
- To be able to solve numerical problems requiring differentiation, integration, interpolation and/or optimization.


#### **LEARNING OUTCOMES II**


- To be able to apply iterative solutions to numerical problems.
- To be able to derive linear/nonlinear systems for a given problem description.
- To be able to select and apply a numerical algorithm to a given linear/nonlinear system.

# Introduction









 $\cdot$  Integers in X bits

- Integers in X bits
  - · Exact representation, finite range

- Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 =

- Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components

- Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - 10/3 = 3.3333...

- $\cdot$  Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - $\cdot 10/3 = 3.3333...$
  - · Keep as fraction, encode integer components:  $\frac{10}{3}$

- $\cdot$  Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - $\cdot$  10/3 = 3.3333...
  - Keep as fraction, encode integer components:  $\frac{10}{3}$
  - Or approximate with finite number of digits
- · Real numbers

- $\cdot$  Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - $\cdot$  10/3 = 3.3333...
  - Keep as fraction, encode integer components:  $\frac{10}{3}$
  - Or approximate with finite number of digits
- · Real numbers
  - · When do we start to have a *real* problem?

- $\cdot$  Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - $\cdot$  10/3 = 3.3333...
  - · Keep as fraction, encode integer components:  $\frac{10}{3}$
  - Or approximate with finite number of digits
- · Real numbers
  - · When do we start to have a real problem?
  - ·  $\pi$


- $\cdot$  Integers in X bits
  - · Exact representation, finite range
  - Arithmetic is modular: Using unsigned 8 bits 255 + 1 = 0
- Fractions with integer components
  - $\cdot$  10/3 = 3.3333...
  - · Keep as fraction, encode integer components:  $\frac{10}{3}$
  - Or approximate with finite number of digits
- · Real numbers
  - · When do we start to have a real problem?
  - · π
  - $\cdot \sqrt{2}$

#### AN OLD COMPUTATIONAL EXAMPLE

Around the year 1000, Persian scientist Abu Rayhan al-Biruni calculated the Earth's radius as 6339.9 km, which is 6356.75 km.


#### AN OLD COMPUTATIONAL EXAMPLE

Around the year 1000, Persian scientist Abu Rayhan al-Biruni calculated the Earth's radius as 6339.9 km, which is 6356.75 km.



# AN OLD COMPUTATIONAL EXAMPLE

Around the year 1000, Persian scientist Abu Rayhan al-Biruni calculated the Earth's radius as 6339.9 km, which is 6356.75 km.



 $R=rac{h\cos heta}{1-\cos heta}$  ightarrowHow do you compute  $\cos heta$  in the year 1000?

# COMPUTING TRIGONOMETRIC FUNCTIONS WITHOUT A CALCULATOR

How do you compute  $\cos\theta$  in the year 1000?

# COMPUTING TRIGONOMETRIC FUNCTIONS WITHOUT A CALCULATOR

How do you compute  $\cos\theta$  in the year 1000? You look at a table of pre-calculated values.

# COMPUTING TRIGONOMETRIC FUNCTIONS WITHOUT A CALCULATOR

How do you compute  $\cos\theta$  in the year 1000? You look at a table of pre-calculated values.

Alexandrian scientist Ptolemy already tabulated some trigonometric quantities as early as the  $2^{nd}$  century. He did his calculations based on the Pythagorean theorem and some trigonometric identities such as

$$\sin^2\theta + \sin^2(90 - \theta) = 1,$$

and

$$\sin\frac{\theta}{2} = \sqrt{\frac{1 - \cos\theta}{2}}.$$

With these mathematical tools, it is possible to compute  $\sin 45^{\circ}$ ,  $\sin 30^{\circ}$ ,  $\sin 75^{\circ}$ ,  $\sin 36^{\circ}$ ,  $\sin 72^{\circ}$ , and  $\sin 3^{\circ}$  accurately, which means we can compute sine for degree that is a multiple of three<sup>1</sup>.

<sup>&</sup>lt;sup>1</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

With these mathematical tools, it is possible to compute  $\sin 45^{\circ}$ ,  $\sin 30^{\circ}$ ,  $\sin 75^{\circ}$ ,  $\sin 36^{\circ}$ ,  $\sin 72^{\circ}$ , and  $\sin 3^{\circ}$  accurately, which means we can compute sine for degree that is a multiple of three<sup>1</sup>.

However, it is not possible to compute  $\sin 1^\circ$ . Ptolemy approximated it by using the following theorem

### **Theorem**

If 
$$\beta < \alpha$$
, then  $\frac{\alpha}{\beta} > \frac{\sin \alpha}{\sin \beta}$ .

By substituting  $\alpha=1, \beta=\frac{3}{4}$  and then  $\alpha=\frac{3}{2}, \beta=1$  we get

<sup>&</sup>lt;sup>1</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

With these mathematical tools, it is possible to compute  $\sin 45^{\circ}, \sin 30^{\circ}, \sin 75^{\circ}, \sin 36^{\circ}, \sin 72^{\circ}$ , and  $\sin 3^{\circ}$  accurately, which means we can compute sine for degree that is a multiple of three<sup>1</sup>.

However, it is not possible to compute  $\sin 1^{\circ}$ . Ptolemy approximated it by using the following theorem

#### **Theorem**

If 
$$\beta < \alpha$$
, then  $\frac{\alpha}{\beta} > \frac{\sin \alpha}{\sin \beta}$ .

By substituting  $\alpha=1, \beta=\frac{3}{4}$  and then  $\alpha=\frac{3}{2}, \beta=1$  we get

$$\frac{2\sin\frac{3}{2}^{\circ}}{3} < \sin 1^{\circ} < \frac{4\sin\frac{3}{4}^{\circ}}{3}$$

$$0.01745130 < \sin 1^{\circ} < 0.01745279$$

<sup>&</sup>lt;sup>1</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

With these mathematical tools, it is possible to compute  $\sin 45^{\circ}$ ,  $\sin 30^{\circ}$ ,  $\sin 75^{\circ}$ ,  $\sin 36^{\circ}$ ,  $\sin 72^{\circ}$ , and  $\sin 3^{\circ}$  accurately, which means we can compute sine for degree that is a multiple of three<sup>1</sup>.

However, it is not possible to compute  $\sin 1^{\circ}$ . Ptolemy approximated it by using the following theorem

#### **Theorem**

If 
$$\beta < \alpha$$
, then  $\frac{\alpha}{\beta} > \frac{\sin \alpha}{\sin \beta}$ .

By substituting  $\alpha=1, \beta=\frac{3}{4}$  and then  $\alpha=\frac{3}{2}, \beta=1$  we get

$$\frac{2\sin\frac{3}{2}^{\circ}}{3} < \sin 1^{\circ} < \frac{4\sin\frac{3}{4}^{\circ}}{3}$$

 $0.01745130 < \sin 1^{\circ} < 0.01745279$ 

<sup>&</sup>lt;sup>1</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

### A More Accurate Solution for $\sin 1^{\circ}$

Persian astronomer Al-Kāshī went further by exploiting a cubic equation<sup>2</sup>

$$\sin 3\theta = 3\sin \theta - 4\sin^3 \theta$$
$$\sin 3^\circ = 3\sin 1^\circ - 4\sin^3 1^\circ$$

<sup>&</sup>lt;sup>2</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

# A More Accurate Solution for $\sin 1^\circ$

Persian astronomer Al-Kāshī went further by exploiting a cubic equation<sup>2</sup>

$$\sin 3\theta = 3\sin \theta - 4\sin^3 \theta$$
$$\sin 3^\circ = 3\sin 1^\circ - 4\sin^3 1^\circ$$

Writing  $x=\sin 1^\circ \to x=\frac{\sin 3^\circ + 4x^3}{3}$ , we can find  $\sin 1^\circ$  as the intersection point of the two simple curves y=x and  $y=\frac{4}{3}x^3+\frac{\sin 3^\circ}{3}$ .

<sup>&</sup>lt;sup>2</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

### A More Accurate Solution for $\sin 1^\circ$

Persian astronomer Al-Kāshī went further by exploiting a cubic equation<sup>2</sup>

$$\sin 3\theta = 3\sin \theta - 4\sin^3 \theta$$
$$\sin 3^\circ = 3\sin 1^\circ - 4\sin^3 1^\circ$$

Writing  $x=\sin 1^\circ \to x=\frac{\sin 3^\circ + 4x^3}{3}$ , we can find  $\sin 1^\circ$  as the intersection point of the two simple curves y=x and  $y=\frac{4}{3}x^3+\frac{\sin 3^\circ}{3}$ .

**Note** We could solve the cubic equation analytically (instead of numerically as follows) but the solution was not known at the time of Al-Kāshī.

<sup>&</sup>lt;sup>2</sup>See "Heavenly Mathematics: The Forgotten Art of Spherical Trigonometry"

#### Fixed Point Iterations for $\sin 1^{\circ}$

$$x_0 = 0.017450000000000$$

$$x_1 = 0.017452403505815$$

#### FIXED POINT ITERATIONS FOR $\sin 1^\circ$

$$x_0 = 0.017450000000000$$

$$x_1 = 0.017452403505815$$

$$x_2 = 0.017452406433712$$

Let's start from the previous solution  $x_0=0.01745$  and iterate with  $x_{k+1}=\frac{4}{3}x_k^3+\frac{\sin 3^\circ}{3}$  keeping 16 digits:

We have converged to x=0.017452406437284. You can verify with your favorite calculator that this is the correct value for  $\sin 1^\circ$  with 16 digits.

# **NUMERICAL COMPUTATION OVERVIEW**

· Numbers on a digital computers have finite-precision.

- · Numbers on a digital computers have finite-precision.
- · Not all problems have closed-form analytic solutions.

- · Numbers on a digital computers have finite-precision.
- · Not all problems have closed-form analytic solutions.
- Sometimes the numerical solution is faster than the analytic one.

- Numbers on a digital computers have finite-precision.
- Not all problems have closed-form analytic solutions.
- Sometimes the numerical solution is faster than the analytic one.
- Numerical solutions are more generic so in some cases implementation is simpler.

- · Numbers on a digital computers have finite-precision.
- · Not all problems have closed-form analytic solutions.
- Sometimes the numerical solution is faster than the analytic one.
- Numerical solutions are more generic so in some cases implementation is simpler.
- Sometimes the numerical algorithm may not converge or may not be numerically stable.

### **NUMERICAL COMPUTATION OVERVIEW**

- · Numbers on a digital computers have finite-precision.
- · Not all problems have closed-form analytic solutions.
- Sometimes the numerical solution is faster than the analytic one.
- Numerical solutions are more generic so in some cases implementation is simpler.
- Sometimes the numerical algorithm may not converge or may not be numerically stable.

The course will cover all these points for numerical solutions of a variety of problems faced by computer engineers and scientists.