

CENG 216 – NUMERICAL COMPUTATION

SYSTEMS OF EQUATIONS - PART I

Mustafa Özuysal

`mustafaozuysal@iyte.edu.tr`

March 13, 2022

İzmir Institute of Technology

Slides are based on material from the main textbook:

“Numerical Analysis”, The new international edition, 2ed,
by Timothy Sauer

SYSTEMS OF EQUATIONS

- Previous weeks: Find the root of a single equation

INTRODUCTION

- Previous weeks: Find the root of a single equation
- Next two weeks: Find a common solution to sets of equations

INTRODUCTION

- Previous weeks: Find the root of a single equation
- Next two weeks: Find a common solution to sets of equations
- This week (Part I): Sections 2.1 and 2.2 from the main textbook


A SYSTEM OF EQUATIONS

Given the system

$$x + y = 3$$

$$3x - 4y = 2,$$

our aim is to find the common solution x^* to both equations.



GAUSSIAN ELIMINATION

NAIVE GAUSSIAN ELIMINATION

1. Swap one equation for another
2. Add or subtract one equation from another
3. Multiply an equation by a non-zero number

NAIVE GAUSSIAN ELIMINATION

1. Swap one equation for another
2. Add or subtract one equation from another
3. Multiply an equation by a non-zero number

$$x + y = 3$$

$$3x - 4y = 2$$

NAIVE GAUSSIAN ELIMINATION

1. Swap one equation for another
2. Add or subtract one equation from another
3. Multiply an equation by a non-zero number

$$\begin{array}{ccc} x + y = 3 & \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} & x + y = 3 \\ 3x - 4y = 2 & & -7y = -7 \end{array}$$

NAIVE GAUSSIAN ELIMINATION

1. Swap one equation for another
2. Add or subtract one equation from another
3. Multiply an equation by a non-zero number

$$\begin{array}{lcl} x + y = 3 & \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} & x + y = 3 \\ 3x - 4y = 2 & & -7y = -7 \end{array} \rightarrow y = 1, x = 2$$

NAIVE GAUSSIAN ELIMINATION

1. Swap one equation for another
2. Add or subtract one equation from another
3. Multiply an equation by a non-zero number

$$\begin{array}{rcl} x + y = 3 & \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} & x + y = 3 \\ 3x - 4y = 2 & & -7y = -7 \end{array} \rightarrow y = 1, x = 2$$

In matrix notation:

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 3 & -4 & 2 \end{array} \right] \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \left[\begin{array}{cc|c} 1 & 1 & 3 \\ 0 & -7 & -7 \end{array} \right] \rightarrow y = 1, x = 2$$

EXAMPLE

$$x + 2y - z = 3$$

$$2x + y - 2z = 3$$

$$-3x + y + z = -6$$

EXAMPLE

$$x + 2y - z = 3$$

$$2x + y - 2z = 3$$

$$-3x + y + z = -6$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right]$$

EXAMPLE

$$x + 2y - z = 3$$

$$2x + y - 2z = 3$$

$$-3x + y + z = -6$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ -3 & 1 & 1 & -6 \end{array} \right]$$

EXAMPLE

$$x + 2y - z = 3$$

$$2x + y - 2z = 3$$

$$-3x + y + z = -6$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ -3 & 1 & 1 & -6 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 7 & -2 & 3 \end{array} \right]$$

EXAMPLE

$$x + 2y - z = 3$$

$$2x + y - 2z = 3$$

$$-3x + y + z = -6$$

$$\left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ -3 & 1 & 1 & -6 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 7 & -2 & 3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 0 & -2 & -4 \end{array} \right]$$

EXAMPLE

$$\begin{array}{rcl} x + 2y - z = 3 & & x + 2y - z = 3 \\ 2x + y - 2z = 3 & \rightarrow & -3y = -3 \\ -3x + y + z = -6 & & -2z = -4 \end{array}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ -3 & 1 & 1 & -6 \end{array} \right] \\ & \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 7 & -2 & 3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 0 & -2 & -4 \end{array} \right] \end{aligned}$$

EXAMPLE

$$\begin{array}{rcl}
 x + 2y - z = 3 & x + 2y - z = 3 & z = 2 \\
 2x + y - 2z = 3 & \rightarrow \quad -3y = -3 & \xrightarrow{\text{backsubstitution}} y = 1 \\
 -3x + y + z = -6 & -2z = -4 & x = 3
 \end{array}$$

$$\begin{aligned}
 & \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 2 & 1 & -2 & 3 \\ -3 & 1 & 1 & -6 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ -3 & 1 & 1 & -6 \end{array} \right] \\
 & \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 7 & -2 & 3 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 2 & -1 & 3 \\ 0 & -3 & 0 & -3 \\ 0 & 0 & -2 & -4 \end{array} \right]
 \end{aligned}$$

OPERATION COUNTS

For an $n \times n$ square matrix:

OPERATION COUNTS

For an $n \times n$ square matrix:

- Elimination takes $\frac{2}{3}n^3 + \frac{1}{2}n^2 - \frac{7}{6}n \approx \frac{2}{3}n^3$ operations.

OPERATION COUNTS

For an $n \times n$ square matrix:

- Elimination takes $\frac{2}{3}n^3 + \frac{1}{2}n^2 - \frac{7}{6}n \approx \frac{2}{3}n^3$ operations.
- Backsubstitution takes n^2 operations.
- See main textbook for the derivation of these.

OPERATION COUNTS

For an $n \times n$ square matrix:

- Elimination takes $\frac{2}{3}n^3 + \frac{1}{2}n^2 - \frac{7}{6}n \approx \frac{2}{3}n^3$ operations.
- Backsubstitution takes n^2 operations.
- See main textbook for the derivation of these.

Ex: $n = 500$, if elimination takes ~ 1 second, then what is the time cost of backsubstitution?

OPERATION COUNTS

For an $n \times n$ square matrix:

- Elimination takes $\frac{2}{3}n^3 + \frac{1}{2}n^2 - \frac{7}{6}n \approx \frac{2}{3}n^3$ operations.
- Backsubstitution takes n^2 operations.
- See main textbook for the derivation of these.

Ex: $n = 500$, if elimination takes ~ 1 second, then what is the time cost of backsubstitution?

$$\frac{2}{3}500^3 t_{op} = 1 \text{ sec.}$$

$$500^2 t_{op} = \frac{\frac{3}{2}}{500} \text{ sec.}$$
$$\approx 0.003 \text{ sec.}$$

THE LU FACTORIZATION

MATRIX FORM OF GAUSSIAN ELIMINATION

$$x + y = 3$$

$$3x - 4y = 2$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

Def:

- $m \times n$ matrix \mathbf{L} is **lower triangular** if $l_{ij} = 0$, for $i < j$.

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

Def:

- $m \times n$ matrix \mathbf{L} is **lower triangular** if $l_{ij} = 0$, for $i < j$.
- $m \times n$ matrix \mathbf{U} is **upper triangular** if $u_{ij} = 0$, for $i > j$.

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

Def:

- $m \times n$ matrix \mathbf{L} is **lower triangular** if $l_{ij} = 0$, for $i < j$.
- $m \times n$ matrix \mathbf{U} is **upper triangular** if $u_{ij} = 0$, for $i > j$.

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

Def:

- $m \times n$ matrix \mathbf{L} is **lower triangular** if $l_{ij} = 0$, for $i < j$.
- $m \times n$ matrix \mathbf{U} is **upper triangular** if $u_{ij} = 0$, for $i > j$.

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{array}{rcl} x + y = 3 \\ 3x - 4y = 2 \end{array} \rightarrow \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix} \rightarrow \mathbf{Ax} = \mathbf{b}$$

Def:

- $m \times n$ matrix \mathbf{L} is **lower triangular** if $l_{ij} = 0$, for $i < j$.
- $m \times n$ matrix \mathbf{U} is **upper triangular** if $u_{ij} = 0$, for $i > j$.

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = \mathbf{U}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

L: 2×2 matrix with 1's on the diagonal and 3 in l_{21} .

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

L: 2×2 matrix with 1's on the diagonal and 3 in l_{21} .

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

L: 2×2 matrix with 1's on the diagonal and 3 in l_{21} .

$$L = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

L: 2×2 matrix with 1's on the diagonal and 3 in l_{21} .

$$L = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$$

$$\rightarrow LU = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} .$$

MATRIX FORM OF GAUSSIAN ELIMINATION

$$\begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 3 \times \text{row 1}} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = U$$

L: 2×2 matrix with 1's on the diagonal and 3 in l_{21} .

$$L = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$$

$$\rightarrow LU = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -7 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 3 & -4 \end{bmatrix} = A.$$

EXAMPLE

Find the LU factorization of $A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}$.

EXAMPLE

Find the LU factorization of $A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}$.

$$\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix}$$

EXAMPLE

Find the LU factorization of $A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}$.

$$\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} \xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix}$$
$$\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix}$$

EXAMPLE

Find the LU factorization of $A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}$.

$$\begin{aligned} \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U. \end{aligned}$$

EXAMPLE

$$\begin{aligned} A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U. \end{aligned}$$

EXAMPLE

$$\begin{aligned} A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U. \end{aligned}$$

$$L = \begin{bmatrix} 1 & & \\ & 0 & 0 \\ & 1 & 0 \\ & & 1 \end{bmatrix}$$

EXAMPLE

$$\begin{aligned} A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U. \end{aligned}$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ & & 1 \end{bmatrix}$$

EXAMPLE

$$\begin{aligned} A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\ &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U. \end{aligned}$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & & 1 \end{bmatrix}$$

EXAMPLE

$$\begin{aligned}
 A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\
 &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\
 &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U.
 \end{aligned}$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix}$$

EXAMPLE

$$\begin{aligned}
 A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} &\xrightarrow[\text{from row 2}]{\text{Subtract } 2 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ -3 & 1 & 1 \end{bmatrix} \\
 &\xrightarrow[\text{from row 3}]{\text{Subtract } -3 \times \text{row 1}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 7 & -2 \end{bmatrix} \\
 &\xrightarrow[\text{from row 3}]{\text{Subtract } -\frac{7}{3} \times \text{row 2}} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = U.
 \end{aligned}$$

$$LU = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix} = A.$$

$$A\mathbf{x} = \mathbf{b} \rightarrow LU\mathbf{x} = \mathbf{b}$$

BACK SUBSTITUTION WITH LU

$$A\mathbf{x} = \mathbf{b} \rightarrow LU\mathbf{x} = \mathbf{b}$$

1. Define $\mathbf{c} = U\mathbf{x}$ and solve $L\mathbf{c} = \mathbf{b}$ for \mathbf{c} .

BACK SUBSTITUTION WITH LU

$$A\mathbf{x} = \mathbf{b} \rightarrow L\mathbf{U}\mathbf{x} = \mathbf{b}$$

1. Define $\mathbf{c} = \mathbf{U}\mathbf{x}$ and solve $L\mathbf{c} = \mathbf{b}$ for \mathbf{c} .
2. Solve $\mathbf{U}\mathbf{x} = \mathbf{c}$ for \mathbf{x} .

EXAMPLE

$$\overbrace{\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}}^{\mathbf{A}} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \overbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}^{\mathbf{b}}$$

EXAMPLE

$$\overbrace{\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}}^{\mathbf{A}} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \overbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}^{\mathbf{b}} \rightarrow \overbrace{\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix}}^{\mathbf{L}} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \overbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}^{\mathbf{b}}$$

EXAMPLE

$$\overbrace{\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}}^{\mathbf{A}} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \overbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}^{\mathbf{b}} \rightarrow \overbrace{\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix}}^{\mathbf{L}} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \overbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}^{\mathbf{b}}$$

$$c_1 = 3$$

$$\rightarrow c_2 = -3$$

$$c_3 = -6 + 9 - 7 = -4$$

EXAMPLE

$$\underbrace{\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}_b \rightarrow \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix}}_L \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}_b$$

$$c_1 = 3$$

$$\rightarrow c_2 = -3$$

$$c_3 = -6 + 9 - 7 = -4$$

$$\underbrace{\begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix}}_U \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ -3 \\ -4 \end{bmatrix}}_c$$

EXAMPLE

$$\underbrace{\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & -2 \\ -3 & 1 & 1 \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}_b \rightarrow \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & -\frac{7}{3} & 1 \end{bmatrix}}_L \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ -6 \end{bmatrix}}_b$$

$$c_1 = 3$$

$$\rightarrow c_2 = -3$$

$$c_3 = -6 + 9 - 7 = -4$$

$$\underbrace{\begin{bmatrix} 1 & 2 & -1 \\ 0 & -3 & 0 \\ 0 & 0 & -2 \end{bmatrix}}_U \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ -3 \\ -4 \end{bmatrix}}_c \rightarrow \begin{aligned} x_3 &= 2 \\ x_2 &= 1 \\ x_1 &= 3 + 2 - 2 = 3. \end{aligned}$$

COMPLEXITY OF LU DECOMPOSITION

Solving multiple equations with the same A but different \mathbf{b} .

$$A\mathbf{x} = \mathbf{b}_1$$

$$A\mathbf{x} = \mathbf{b}_2$$

$$\vdots$$

$$A\mathbf{x} = \mathbf{b}_k$$

COMPLEXITY OF LU DECOMPOSITION

Solving multiple equations with the same \mathbf{A} but different \mathbf{b} .

$$\left. \begin{array}{l} \mathbf{Ax} = \mathbf{b}_1 \\ \mathbf{Ax} = \mathbf{b}_2 \\ \vdots \\ \mathbf{Ax} = \mathbf{b}_k \end{array} \right\} \begin{array}{l} \text{cost of naive} \\ \text{Gaussian elimination} \end{array} \rightarrow \frac{2}{3}kn^3$$

COMPLEXITY OF LU DECOMPOSITION

Solving multiple equations with the same \mathbf{A} but different \mathbf{b} .

$$\left. \begin{array}{l} \mathbf{Ax} = \mathbf{b}_1 \\ \mathbf{Ax} = \mathbf{b}_2 \\ \vdots \\ \mathbf{Ax} = \mathbf{b}_k \end{array} \right\} \begin{array}{l} \text{cost of naive} \\ \text{Gaussian elimination} \end{array} \rightarrow \frac{2}{3}kn^3$$
$$\left. \begin{array}{l} \mathbf{Ax} = \mathbf{b}_1 \\ \mathbf{Ax} = \mathbf{b}_2 \\ \vdots \\ \mathbf{Ax} = \mathbf{b}_k \end{array} \right\} \begin{array}{l} \text{LU decomposition} \\ \text{and backsubstitution} \end{array} \rightarrow \frac{2}{3}n^3 + 2kn^2$$

COMPLEXITY OF LU DECOMPOSITION

Solving multiple equations with the same \mathbf{A} but different \mathbf{b} .

$$\left. \begin{array}{l} \mathbf{Ax} = \mathbf{b}_1 \\ \mathbf{Ax} = \mathbf{b}_2 \\ \vdots \\ \mathbf{Ax} = \mathbf{b}_k \end{array} \right\} \begin{array}{l} \text{cost of naive} \\ \text{Gaussian elimination} \rightarrow \frac{2}{3}kn^3 \\ \\ \text{LU decomposition} \\ \text{and backsubstitution} \rightarrow \frac{2}{3}n^3 + 2kn^2 \end{array}$$

Solving k equations with LU factorization means that we need to run the costly elimination step ($\frac{2}{3}n^3$ operations) only once. For some applications this can be done in an offline preprocessing step and the solutions can be performed online as data becomes available.

LU FACTORIZATION FOR GENERAL INVERTIBLE MATRICES

Ex: Prove that $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ does not have an LU factorization although it is invertible.

LU FACTORIZATION FOR GENERAL INVERTIBLE MATRICES

Ex: Prove that $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ does not have an LU factorization although it is invertible.

$$\text{Assume } \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} b & c \\ 0 & d \end{bmatrix} = \begin{bmatrix} b & c \\ ab & ac + d \end{bmatrix}$$

LU FACTORIZATION FOR GENERAL INVERTIBLE MATRICES

Ex: Prove that $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ does not have an LU factorization although it is invertible.

$$\text{Assume } \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} b & c \\ 0 & d \end{bmatrix} = \begin{bmatrix} b & c \\ ab & ac + d \end{bmatrix}$$

$$\rightarrow b = 0 \text{ and } ab = 1 \text{ (Contradiction ✖)}$$

LU FACTORIZATION FOR GENERAL INVERTIBLE MATRICES

Ex: Prove that $A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ does not have an LU factorization although it is invertible.

$$\text{Assume } \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix} \begin{bmatrix} b & c \\ 0 & d \end{bmatrix} = \begin{bmatrix} b & c \\ ab & ac + d \end{bmatrix}$$

$$\rightarrow b = 0 \text{ and } ab = 1 \text{ (Contradiction ✖)}$$

Next week, we will generalize LU factorization to handle these kinds of matrices.