Truscan Blogs November

Blog 1

Diabetes: Foods to Eat and Foods to Avoid – A Practical Guide for an Indian Diet

A diabetes diagnosis often creates one immediate question: “What can I eat now?”

Many people assume that managing diabetes means completely avoiding rice, fruits, traditional Indian meals, or anything that tastes sweet. In reality, diabetes nutrition is more nuanced. The goal is not necessarily to eliminate every carbohydrate or favourite food. Instead, a sustainable diabetes diet focuses on what you eat, how much you eat, how often you eat it, and what other foods you combine with it.

For people living with diabetes, food choices play an important role in maintaining blood glucose within an appropriate target range. A balanced eating pattern can also support healthy body weight, cardiovascular health, and overall well-being.

Health organizations like the Centers for Disease Control and Prevention (CDC) recommend considering nutrition needs, lifestyle, preferences, and medications when

developing a diabetes meal plan. They also highlight approaches such as carbohydrate awareness and the Plate Method. For Indian families, diabetes meal planning can be especially practical when traditional foods are modified rather than completely eliminated.

What Is a Diabetes-Friendly Diet?

A diabetes-friendly diet is an eating pattern designed to provide adequate nutrition while helping manage blood glucose.

There is no universal diabetes diet that works exactly the same way for everyone. A meal plan depends on several individual factors:

- Type of diabetes
- Age and body weight
- Physical activity level
- Medications and insulin use
- Other medical conditions
- Personal food preferences and cultural eating patterns
- Individual blood glucose response

Therefore, the question should not simply be *"Which foods are good for diabetes?"*

A better question is: “How can I build balanced meals that fit my health needs?”

A practical starting point is to include plenty of non-starchy vegetables, an appropriate protein source, and a controlled portion of carbohydrate-containing foods.

The Diabetes Plate Method

One of the simplest ways to think about a diabetes-friendly meal is the Plate Method. The CDC recommends filling approximately:

- ½ of the plate with non-starchy vegetables
- ¼ of the plate with lean protein
- ¼ of the plate with carbohydrate-containing foods
- Beverage: Water or a low-calorie drink

This approach is particularly useful for Indian meals because it focuses on proportions rather than requiring people to weigh every ingredient. For example, instead of eating a large serving of rice with a small amount of vegetable curry, the meal can be structured around vegetables first, followed by protein, and a controlled rice portion.

Foods to Eat With Diabetes

1. Non-Starchy Vegetables

Vegetables should form a significant part of diabetes-friendly meals. They provide fiber, vitamins, and

minerals while contributing fewer carbohydrates than rice, bread, or potatoes.

Good options include:

- Leafy Greens: Spinach (*palak*), fenugreek (*methi*), green leafy vegetables
- Gourds: Bitter gourd (*karela*), bottle gourd (*lauki*), ridge gourd (*turai*)
- Cruciferous & Others: Cabbage, cauliflower, broccoli, beans, cucumber, tomato, capsicum, brinjal, radish

Tip: Include vegetables at lunch and dinner rather than treating them as an optional side dish.

2. Protein-Rich Foods

Protein helps make meals more satisfying and can slow down digestion when paired with carbohydrates.

Suitable protein sources include:

- Vegetarian: Dal, chickpeas, green gram, sprouts, beans, paneer, tofu, unsweetened curd
- Non-Vegetarian: Eggs, fish, chicken

3. Whole Fruits

A common misconception is that people with diabetes should never eat fruit. Whole fruit can be part of a

balanced diet when portion size and total carbohydrate intake are managed.

Good options include:

- Guava, apple, orange, pear, papaya, berries, pomegranate, and melon

Note: Whole fruit is preferable to fruit juice because juice provides quick carbohydrates without the intact fiber found in whole fruit. Fruit juice can raise blood glucose much faster than whole fruit.

4. Whole Grains and High-Fibre Carbohydrates

Carbohydrates are not automatically unhealthy. High-fiber options can be included in appropriate portions:

- Whole wheat, oats, barley, brown rice, selected millets, and whole-grain preparations

Caution: A food being labeled “whole grain” or "healthy" does not mean unlimited quantities are appropriate for blood glucose management.

Common Questions About Indian Carbohydrates

Can People With Diabetes Eat Rice?

Yes. Rice can be included in a diabetes meal plan depending on overall dietary requirements and blood

glucose response. The quantity eaten and the foods eaten alongside it matter.

- Less balanced meal: Large serving of white rice + potato curry + fried side dish
- More balanced approach: Controlled rice portion + large serving of vegetables + dal/protein + salad

What About Ragi, Millets, and Other “Healthy” Grains?

Millets (*ragi, jowar, bajra*) and traditional grains are nutritious additions to a varied diet. However, assuming that a food labeled “diabetic-friendly,” “millet-based,” or “sugar-free” can be consumed without portion control is a common mistake. Millets still contain carbohydrates, so overall portion size remains important.

Foods to Limit or Avoid With Diabetes

1. Sugary Drinks

Sugary beverages deliver a large amount of sugar quickly without offering fullness.

- Soft drinks, sweetened packaged juices, sweetened tea/coffee, energy drinks, and sugar-heavy milkshakes

2. Sweets and Desserts

Indian sweets contain high amounts of sugar and refined carbohydrates.

- *Gulab jamun, jalebi, mysore pak, laddu, rasgulla*, cakes, pastries, and sweet biscuits

3. Refined Carbohydrates

Highly refined foods are easily overconsumed and contain less dietary fiber.

- *Maida*-based bakery products, white bread, sweet biscuits, cakes, pastries, refined noodles, and packaged snacks

4. Deep-Fried Foods

- *Pakoda, bhajji, samosa*, and deep-fried fast foods add substantial calories and fat, which can make weight management difficult and affect cardiovascular health.

Is “Sugar-Free” Food Safe for Diabetes?

Not necessarily. “Sugar-free” does not automatically mean “low carbohydrate” or “low calorie.” Some packaged foods use alternative sweeteners but still contain substantial amounts of carbohydrates or fats. Always check the nutritional facts label.

Diabetes-Friendly Meal Ideas

Breakfast Ideas

- Vegetable omelette with a whole-grain accompaniment
- Vegetable oats or vegetable upma (in a controlled portion)
- Idli with sambar and vegetables
- Vegetable dosa with a protein-rich side
- Moong dal chilla
- Unsweetened curd with suitable seeds/additions

Lunch & Dinner Ideas

- Vegetables: Large serving of cooked vegetable curry or salad
- Protein: Dal, sprouts, fish, chicken, paneer, or tofu
- Carbohydrates: Controlled portion of rice or whole-wheat/millet roti
- Dairy: Unsweetened curd where appropriate

Common Diabetes Diet Mistakes

1. Eliminating All Carbohydrates: Carbohydrates are present in many nutritious foods; complete removal is rarely necessary.
2. Drinking Fruit Juice Instead of Eating Fruit: Juice lacks fiber and raises blood sugar faster.

3. Overeating “Healthy” Foods: Nuts, seeds, millets, and whole grains still contain calories and carbohydrates that require portion control.
4. Skipping Meals: Skipping meals can cause blood sugar fluctuations, especially for individuals taking certain medications.
5. Changing Medication Based on Diet Alone: Medication adjustments should only be made under direct medical guidance.

Why Blood Glucose Monitoring and Diagnostic Testing Matter

Food is only one component of diabetes management. Depending on your health profile, healthcare professionals may recommend routine lab testing:

- Blood Glucose Monitoring & HbA1c Testing
- Lipid Profile (Heart Health)
- Kidney & Liver Function Assessments
- Blood Pressure Checks

Testing helps evaluate how well your diet, lifestyle, and treatment plan are working together. Diagnostic centers like TruScan Diagnostics offer services such as pathology testing and the TruScan Basic Health Package to support routine laboratory evaluation when clinically recommended. Laboratory results should always be

interpreted by a qualified doctor in the context of your medical history.

Frequently Asked Questions (FAQs)

Q1. What are the best foods to eat for diabetes?

Ans: A diabetes-friendly diet includes plenty of non-starchy vegetables (like spinach, methi, cauliflower, cucumber, and bitter gourd), protein sources (dal, sprouts, paneer, tofu, eggs, and fish), whole fruits in controlled portions, and fibre-rich whole grains (oats, brown rice, whole wheat, and millets).

Q2. What foods should people with diabetes avoid or limit?

Ans: It is best to limit or avoid sugary beverages (soft drinks, packaged juices, sweetened tea/coffee), refined carbohydrates (maida products, white bread, bakery biscuits), traditional sweets (*gulab jamun, jalebi, laddu*), and deep-fried foods (*samosa, pakoda, bhajji*).

Q3. Can people with diabetes eat rice?

Ans: Yes. Rice can be part of a diabetes diet. The key is portion control and meal balance. Instead of eating a large plate of rice alone, keep the rice portion to about ¼ of your

plate and combine it with plenty of non-starchy vegetables and a good protein source like dal, sprouts, or chicken.

Q4. Is brown rice better than white rice for diabetes?

Ans: Brown rice generally contains more dietary fibre and nutrients than white rice, which helps slow down digestion. However, brown rice still contains carbohydrates, so portion control is equally important for both white and brown rice.

Q5. Can diabetics eat fruits?

Ans: Yes, whole fruits can be eaten. Fruits like guava, apple, orange, papaya, pear, and berries offer valuable vitamins and fibre. However, eat whole fruit instead of drinking fruit juice, as juices lack intact fibre and can raise blood sugar levels quickly.

Q6. Can people with diabetes eat ragi, jowar, or other millets without limits?

Ans: No. While millets like ragi, jowar, and bajra are rich in fibre and minerals, they still contain carbohydrates. Eating large portions of millet-based foods (like ragi dosa or millet roti) will still impact blood sugar, so portion control remains necessary.

Q7. Are "sugar-free" packaged foods completely safe for diabetes?

Ans: Not necessarily. A label that says "sugar-free" does not mean it is free from carbohydrates, calories, or fat. Always read the Nutritional Information panel to check the total carbohydrate content rather than relying only on front-of-pack claims.

Q8. Should people with diabetes get regular blood tests?

Ans: Yes. Nutrition is only one part of diabetes management. Routine diagnostic checks—such as HbA1c, Fasting Blood Glucose, Postprandial Blood Glucose, Lipid Profile, and Kidney Function Tests—help your doctor evaluate how well your diet, lifestyle, and medications are working.

Q9. Where can I get diabetes-related lab tests in Bengaluru?

Ans: Patients in Bengaluru can access diagnostic facilities like TruScan Diagnostics, which offer relevant pathology services and routine health checkup packages (such as the TruScan Basic Health Package) as recommended by their physician.

Meta Title: Diabetes Diet Plan: Foods to Eat & Avoid in an Indian Diet

Meta Description: Discover what foods to eat and avoid with diabetes in an Indian diet. Learn how to balance rice, rotis, fruits & millets using the CDC Plate Method.

H1 Tag: `<h1>Diabetes Diet Plan: Foods to Eat and Foods to Avoid (An Indian Guide)</h1>`

Blog 2

Home Blood Sample Collection: How It Works and What Tests Can Be Done

Getting a blood test no longer necessarily means spending time travelling to a diagnostic centre, waiting for your turn and rearranging your entire day. Home blood sample collection has become an increasingly useful option for people who want diagnostic testing to fit more conveniently into their routine.

For working professionals, older adults, parents, people with mobility challenges and individuals managing regular health monitoring, home sample collection can make the testing process more convenient.

But how does blood sample collection at home actually work? What tests can be collected? Do you need to fast? How should you prepare? And how is the sample transported to a laboratory?

This guide explains the process and highlights how TruScan Diagnostics can fit into a convenient diagnostic testing journey in Bengaluru.

What Is Home Blood Sample Collection?

Home blood sample collection is a diagnostic service in which a trained healthcare professional visits the patient's selected location to collect a blood specimen.

The sample is then handled according to appropriate laboratory procedures and transported for testing.

The patient does not necessarily need to visit the laboratory for the collection itself.

This can be particularly useful when a person requires routine laboratory testing but has difficulty travelling.

How Does Home Sample Collection Work?

Although the exact process may vary depending on the diagnostic provider and test, the general process involves several stages.

Step 1: Select the Required Tests

The first step is determining which laboratory tests are required.

Some tests may be recommended by a doctor, while others may be part of preventive health screening.

Examples can include:

- **Blood glucose testing**
- **Lipid testing**
- **Complete blood count**
- **Liver-related investigations**
- **Kidney-related investigations**
- **Thyroid testing**
- **Vitamin-related testing**

- **Hormonal testing**
- **Fertility-related investigations**
- **Other pathology tests**

Not every test is suitable for home collection, so test-specific requirements should always be confirmed beforehand.

Step 2: Check Preparation Requirements

Some blood tests require fasting.

Others do not.

Some tests may have additional preparation requirements involving food, medication, exercise or timing.

It is important not to assume that every blood test requires fasting.

Before the appointment, confirm the preparation instructions for the exact tests being performed.

If you take regular medication, do not stop or change it unless your doctor specifically instructs you to do so.

Step 3: Schedule the Collection

A suitable time and location are selected for sample collection.

For people living or working in Bengaluru, local home collection can be particularly convenient when regular testing is required.

Step 4: Sample Collection

A trained professional collects the required specimen using appropriate collection equipment.

The professional generally confirms the patient's details and the tests required before proceeding.

Step 5: Sample Handling and Transportation

After collection, specimens must be correctly identified, handled and transported according to the requirements of the tests being performed.

This stage is important because laboratory accuracy depends not only on the analytical process but also on proper pre-analytical handling.

Step 6: Laboratory Testing

Once the specimen reaches the appropriate laboratory facility, it is processed according to the relevant testing procedures.

Step 7: Report Review

The final laboratory report provides test results that can be discussed with the patient's doctor.

A laboratory report should not be interpreted in isolation. A doctor may consider symptoms, medical history, medications, physical examination and other investigations before reaching a clinical conclusion.

What Blood Tests Can Be Done at Home?

The answer depends on the diagnostic provider and the specific test.

Common blood investigations may include:

Complete Blood Count

A CBC can provide information about different blood cell types and may be used in evaluating a variety of health conditions.

Blood Glucose Tests

Blood glucose testing can be used in diabetes screening and monitoring.

Lipid Profile

A lipid profile evaluates measures related to blood fats, such as cholesterol and triglycerides.

Thyroid Testing

Thyroid-related blood tests can help evaluate thyroid function when clinically indicated.

Liver Function-Related Tests

Blood investigations can provide information about certain aspects of liver function.

Kidney Function-Related Tests

Certain blood tests can provide information relevant to kidney function.

Vitamin and Nutritional Testing

Specific vitamin or nutritional markers may be tested when medically appropriate.

Hormonal Testing

Certain hormone tests may require specific timing or preparation, so instructions should be confirmed before collection.

Fertility-Related Testing

Some fertility investigations involve blood testing, while others require different specimen types and specialised collection procedures.

Does Every Test Require Fasting?

No.

Fasting requirements depend on the test.

Some investigations are commonly performed after fasting, while others can be conducted without fasting.

Therefore, do not fast simply because you are getting "a blood test."

Instead, confirm the requirements for the specific tests included in your prescription or health package.

How Should You Prepare for a Home Blood Test?

Follow the Test Instructions

Preparation should be based on the tests being performed.

Stay Hydrated Unless Instructed Otherwise

Adequate hydration may make blood collection easier for some people, but follow any specific instructions given for your test.

Wear Comfortable Clothing

Choose clothing that allows easy access to the arm.

Keep Your Prescription or Test List Ready

Having the doctor's prescription or requested test list available can reduce confusion.

Tell the Collector About Relevant Information

If there are specific instructions regarding your medications, fasting status or test timing, communicate them clearly.

Who Can Benefit From Home Sample Collection?

Home collection can be useful for:

- **Older adults**
- **Working professionals**
- **People with mobility limitations**
- **Individuals undergoing regular monitoring**
- **People who find laboratory visits difficult**
- **Families requiring multiple tests**
- **People seeking greater convenience**

It can also be useful for preventive health screening when several members of a family need testing.

Home Blood Sample Collection in Bengaluru

Bengaluru is a large and busy city where commuting can consume significant time.

For residents in areas such as JP Nagar and surrounding parts of Bengaluru, diagnostic home collection can make routine testing easier to incorporate into a busy schedule.

However, convenience should never come at the expense of proper specimen handling, patient identification or appropriate laboratory procedures.

TruScan Diagnostics and Home Sample Collection

TruScan Diagnostics is a relevant diagnostic brand for people looking for pathology and laboratory testing services in Bengaluru.

The TruScan SEO service structure specifically identifies home sample collection, Bengaluru and JP Nagar as strong local search opportunities, with Pathology and TruScan Services as important internal destinations.

For patients considering home blood testing, TruScan can be positioned as part of a convenient diagnostic journey:

Book the required test → Prepare according to instructions → Complete home sample collection → Laboratory processing → Receive the report → Discuss the results with your doctor.

The exact availability of individual tests, collection locations and preparation requirements should always be confirmed with TruScan before scheduling.

Home Collection vs Visiting a Diagnostic Centre

Both options can be appropriate.

Home Collection

Advantages can include:

- **Convenience**
- **Less travel**
- **Useful for elderly patients**
- **Better suited to busy schedules**
- **Convenient for repeat testing**

Diagnostic Centre Visit

A centre visit may be preferable when:

- **A specialised procedure is required**
- **The test requires equipment unavailable at home**
- **Multiple services need to be completed at the centre**

- **The healthcare provider specifically recommends an in-centre procedure**

The best option depends on the test and the patient's circumstances.

How Important Is Sample Handling?

Very important.

Laboratory testing involves several stages, and the period between collecting the sample and analysing it is known as the pre-analytical phase.

Correct identification, collection, storage and transportation are therefore important components of diagnostic quality.

This is one reason patients should use established diagnostic services and follow collection instructions carefully.

Common Questions About Home Blood Tests

Can I get a blood test done at home?

Many routine blood tests can be collected at home, depending on the diagnostic provider and the test.

How does home blood sample collection work?

A trained collection professional visits the requested location, collects the required specimen and ensures it is appropriately handled for laboratory testing.

What tests can be collected at home?

Many routine pathology blood tests may be suitable for home collection, but specialised tests may have different requirements.

Can I get a blood test at home in Bengaluru?

Home blood collection may be available in selected parts of Bengaluru depending on the diagnostic provider and service coverage.

Do I need to fast before a home blood test?

Only when required for the particular test. Always confirm the instructions before the appointment.

Can children have blood samples collected at home?

Depending on the test and provider, paediatric home collection may be possible. Confirm availability and any special requirements beforehand.

Choosing a Diagnostic Provider

When selecting a diagnostic service, consider:

1. **Whether the required test is available.**
2. **Whether home collection is available in your location.**
3. **Whether trained personnel perform collection.**
4. **Whether appropriate sample handling procedures are followed.**
5. **Whether clear preparation instructions are provided.**
6. **Whether reports are accessible conveniently.**
7. **Whether the laboratory has relevant pathology capabilities.**

Final Takeaway

Home blood sample collection can make routine diagnostic testing considerably more convenient, especially for people with demanding schedules, mobility challenges or regular monitoring requirements.

However, convenience should always be combined with proper preparation, trained collection, appropriate specimen handling and reliable laboratory processing.

For patients searching for home blood sample collection in Bengaluru or JP Nagar, TruScan Diagnostics can be considered as a local diagnostic option. Patients should confirm the availability of the specific test and collection service before booking.

Most importantly, use laboratory testing as part of a broader healthcare process. Discuss which investigations are appropriate with your doctor, follow test-specific preparation instructions and review your results with a qualified healthcare professional.

Meta Title: Home Blood Sample Collection in Bengaluru | How It Works & Tests

Meta Description: Book home blood sample collection in Bengaluru & JP Nagar. Learn how it works, test preparation, fasting rules, and available pathology tests.

H1 Tag: `<h1>Home Blood Sample Collection: How It Works and What Tests Can Be Done</h1>`

Blog 3

How Age Affects Male Fertility and Sperm Analysis Test Results

When people discuss fertility and age, the focus is often on women. However, male age can also influence reproductive health, and changes in sperm production, semen parameters and general health can become more relevant as men get older.

This does not mean that men suddenly become infertile after reaching a particular age. Men can father children well into their 40s, 50s and beyond. However, increasing age may be associated with changes in certain sperm parameters and with a gradual increase in some reproductive and pregnancy-related risks.

This is why questions such as “Does male fertility decrease with age?”, “Does age affect sperm count?”, “Does sperm quality decrease after 40?”, “What does a semen analysis test show?” and “When should men get a semen analysis?” are important questions for couples planning a pregnancy.

The most important principle is that age alone cannot determine whether a man is fertile or infertile.

Male fertility depends on multiple factors, including sperm production, sperm movement, sperm morphology, reproductive anatomy, hormones, general health, lifestyle, medications and the reproductive health of the female partner.

The World Health Organization's current infertility guideline recognises infertility as a condition requiring appropriate evaluation and treatment, with male factors representing an important part of fertility care.

For men concerned about fertility, a semen analysis or sperm analysis test is one of the key investigations used to assess semen and sperm characteristics. The WHO's laboratory manual provides standardised procedures for

semen examination and emphasises the importance of laboratory quality and consistent testing.

Does Male Fertility Decrease With Age?

The short answer is yes, some aspects of male reproductive function can change with age—but male fertility does not suddenly end at a specific age.

Men continue producing sperm throughout adult life, although the quantity and characteristics of sperm can change over time.

The American Society for Reproductive Medicine notes that semen parameters decline detectably after approximately age 35, while also explaining that male fertility does not appear to be substantially affected before around age 50.

This illustrates why it is difficult to define one specific age at which a man's fertility becomes "low."

For example:

- A 42-year-old man may have normal semen parameters.
- A 32-year-old man may have a significantly abnormal semen analysis.
- A 48-year-old man may conceive naturally.
- A younger man may experience infertility because of a medical condition.

Therefore, male fertility after 40 should be evaluated according to the individual rather than judged purely by age.

How Does Age Affect Male Fertility?

Ageing can be associated with changes in several aspects of male reproductive health.

Potential changes include:

- Sperm concentration
- Total sperm count
- Sperm motility
- Sperm morphology
- Semen volume
- Sperm DNA integrity
- Hormonal function
- Testicular function

However, these changes do not occur at exactly the same rate in every man.

A man's overall health can significantly influence reproductive function.

For example, an older man who maintains healthy metabolic function, avoids smoking and manages chronic diseases may have very different reproductive parameters from another man of the same age who has obesity, uncontrolled diabetes, heavy alcohol use or other medical conditions.

This is why fertility assessment should combine age + medical history + lifestyle + semen analysis + clinical examination + other appropriate tests.

Male Fertility After 30, 40 and 50

Male Fertility in the 30s

Many men in their 30s have normal reproductive function.

However, semen parameters may already begin to change gradually with age in some men.

A man who is planning to delay fatherhood may therefore benefit from understanding that male fertility is not completely unaffected by age.

This does not mean that every man in his 30s needs fertility testing.

Testing is generally guided by reproductive plans, symptoms, medical history and clinical circumstances.

Male Fertility After 40

Men can absolutely father children after 40.

However, paternal age becomes increasingly relevant when evaluating reproductive health.

The AUA/ASRM guideline advises clinicians to counsel couples with advanced paternal age, defined in that guideline as 40 years or older, about increased risks of adverse health outcomes for offspring.

This does not mean that pregnancy after 40 is unsafe or impossible.

It means that age is one factor worth discussing with a healthcare professional when planning a pregnancy.

Male Fertility After 50

Men can also father children after 50.

However, increasing paternal age can be associated with changes in sperm characteristics and certain reproductive risks.

For men considering fatherhood later in life, discussing reproductive health with a doctor can help identify potentially modifiable risk factors and determine whether fertility testing is appropriate.

What Is Sperm Quality?

The term sperm quality refers to several characteristics rather than one measurement.

A semen analysis can provide information about several of these characteristics.

The major parameters include:

Sperm Concentration

Sperm concentration refers to the number of sperm present per unit volume of semen.

A low concentration may indicate reduced sperm production, although the interpretation depends on the complete report.

Total Sperm Number

This considers the total number of sperm present in the entire ejaculate.

Total sperm number can provide information that complements sperm concentration.

Sperm Motility

Motility describes sperm movement.

For natural conception, sperm need to move through the reproductive tract and reach the egg.

Therefore, sperm movement is an important component of semen analysis.

Sperm Morphology

Morphology refers to the shape and structure of sperm.

Laboratories evaluate sperm according to established criteria.

Morphology can contribute to fertility assessment, but it should not be interpreted in isolation.

Semen Volume

Semen volume measures the amount of semen produced during ejaculation.

Changes in semen volume can have several causes and need to be interpreted alongside the other findings.

The WHO's sixth-edition semen manual provides standardised laboratory methods for examining these and other semen characteristics.

Does Age Affect Sperm Count?

Age can affect sperm count and other semen parameters, but the effect varies from person to person.

As men age, changes in sperm production may occur.

However, there is no single age at which every man's sperm count declines dramatically.
Sperm count can also be influenced by:

- Testicular health
- Hormones
- Varicocele
- Infection
- Obesity
- Smoking
- Alcohol
- Certain medications
- Anabolic steroids
- Chemotherapy
- Genetic conditions
- Other medical conditions

Therefore, if a couple is experiencing difficulty conceiving, a sperm analysis test provides more useful information than simply looking at the man's age.

Does Sperm Quality Decrease After 40?
Some aspects of sperm quality may decline with increasing age.
Research and clinical guidance recognise associations between increasing paternal age and changes in semen parameters and sperm biology.
However, the effect is variable.
The important distinction is:

Age may influence sperm quality, but age does not determine an individual's sperm quality.
A semen analysis can provide actual information about the characteristics of a particular sample.
This is one reason sperm analysis after 40 may be considered when there is a fertility concern.

Does Age Affect Sperm Motility?

Sperm motility can change with age.
Motility is important because sperm need to move effectively to reach the egg.
A semen analysis may assess the percentage of sperm that are motile and provide additional information about movement depending on the laboratory's methodology.
Reduced motility can have many causes, so an abnormal result should not automatically be attributed to age.
A fertility specialist may consider:

- Age
- Medical history
- Hormones
- Lifestyle
- Previous infections
- Testicular health
- Other semen parameters

before determining the significance of the result.

Does Age Affect Sperm Morphology?

Sperm morphology refers to sperm shape and structure.
Age may be associated with changes in morphology in some men, but morphology also varies between individuals and between samples.
Importantly, morphology should not be interpreted as a simple “fertile” or “infertile” measurement.
The AUA/ASRM guideline notes that individual semen parameters such as concentration, morphology and motility are not, by themselves, highly predictive of fertility or diagnostic of infertility.
This is an important point for anyone reading a semen report.

What Is Sperm DNA Integrity?
Sperm carry genetic material that contributes to the embryo.
Ageing has been associated with changes in sperm DNA integrity and other genetic characteristics.
The AUA/ASRM guideline notes that advanced paternal age is associated with increased de novo germline mutations, sperm aneuploidy, structural chromosomal abnormalities and sperm DNA fragmentation.
However, this does not mean that every older man has abnormal sperm DNA or that pregnancy should be avoided solely because of paternal age.
It means that age can be one factor considered during appropriate reproductive counselling.

Advanced sperm testing is not automatically required for every man.
A fertility specialist should determine whether additional testing is clinically appropriate.

What Is a Semen Analysis?

A semen analysis, also called a sperm analysis test, is a laboratory investigation used to examine semen and sperm.
The WHO describes semen analysis as useful for investigating male fertility status and provides standardised methods intended to improve the quality and comparability of semen testing between laboratories.
Depending on the laboratory, the report may provide information about:

- Semen volume
- Sperm concentration
- Total sperm number
- Sperm motility
- Sperm morphology
- Other semen characteristics

A semen analysis is one of the first-line investigations used in the evaluation of male infertility.
The AUA/ASRM guideline recommends that initial male infertility evaluation include a reproductive history and one or more semen analyses.

What Does a Semen Analysis Test Show?

A semen analysis can answer several important questions.

How many sperm are present?

This is assessed through sperm concentration and/or total sperm number.

Are the sperm moving?

Motility provides information about sperm movement.

What do the sperm look like?

Morphology assesses sperm shape and structure.

How much semen is produced?

Semen volume is measured as part of the analysis.

But a semen analysis does not provide a complete answer to every fertility question.

It cannot independently determine:

- Whether a couple will conceive
- The fertility of the female partner
- Every hormonal abnormality
- Every anatomical problem
- All genetic causes
- Every possible sperm-function problem

WHO specifically notes that semen analysis cannot confirm fertility in an individual man because fertility depends on several factors, including the fertility potential of the partner.

Can One Abnormal Semen Analysis Mean Infertility?

No.
This is one of the most important things patients should understand.
Semen parameters are biological measurements and can vary between ejaculates.
An abnormal result therefore needs clinical interpretation.
The AUA/ASRM guideline states that semen parameters can vary substantially between ejaculates and recommends obtaining at least two semen analyses, ideally at least one month apart, particularly when the first test shows abnormal parameters.
Therefore, a patient should not assume that one abnormal report means permanent infertility.
A doctor may recommend repeat testing before making conclusions.

How Should Men Prepare for a Semen Analysis?
Preparation instructions can vary depending on the laboratory.
Patients may receive instructions regarding:

- Sexual abstinence
- Collection method
- Collection container
- Collection location
- Transport
- Time between collection and laboratory processing

Follow the instructions provided by the laboratory performing the test.
Do not rely solely on generic information from the internet because different tests and laboratory protocols may have specific requirements.
The WHO semen manual provides standardised laboratory procedures precisely because collection and processing can influence the quality and comparability of semen analysis.

Why Is Correct Sample Collection Important?

Semen analysis depends on obtaining an appropriate specimen.
If a portion of the ejaculate is accidentally lost, the sample may not accurately represent the complete ejaculate.
Patients should therefore:

1. Follow the laboratory's instructions.
2. Use the container provided or approved by the laboratory.
3. Ensure the complete sample is collected where instructed.
4. Provide the specimen within the specified time if home collection is permitted.
5. Tell the laboratory if there was any problem during collection.

The laboratory can then determine whether the sample is suitable for analysis.

What Causes Male Infertility Besides Age?

Age is only one possible factor.

Varicocele

A varicocele involves enlarged veins around the testicle and can affect sperm production in some men.

Hormonal Problems

Sperm production is regulated by a complex hormonal system involving the brain, pituitary gland and testes. Problems involving hormones such as FSH, LH or testosterone can affect fertility.

Testicular Problems

Conditions affecting the testicles can interfere with sperm production.

Infections

Certain infections can affect the reproductive tract.

Genetic Factors

Some genetic abnormalities can affect sperm production or reproductive function.

Previous Surgery

Some reproductive or pelvic surgeries can affect fertility.

Chemotherapy and Other Treatments

Certain medical treatments can affect sperm-producing cells.

Obesity and Metabolic Health

Metabolic health can influence hormonal and reproductive function.

Smoking

Smoking is associated with adverse reproductive and general health effects.

Excessive Alcohol Consumption

Heavy alcohol use can affect general health and may influence reproductive function.

Anabolic Steroids

Anabolic steroids can suppress natural reproductive hormone signalling and significantly reduce sperm production.

The WHO's current infertility guidance emphasises evidence-based evaluation and treatment of male-factor infertility, while the AUA/ASRM guideline highlights medical, lifestyle and age-related factors in male reproductive assessment.

Can Lifestyle Affect Male Fertility?

Lifestyle does not explain every case of infertility, but healthy habits can support overall reproductive health.

Maintain a Healthy Weight

Excess body weight can be associated with metabolic and hormonal changes.

Avoid Tobacco

Tobacco use is a recognised risk factor for poorer reproductive health.

Limit Excessive Alcohol

Reducing excessive alcohol consumption supports general health.

Exercise Regularly

Regular physical activity supports cardiovascular and metabolic health.

Sleep Adequately

Consistent sleep is important for general hormonal and metabolic health.

Manage Chronic Diseases

Conditions such as diabetes should be appropriately managed.

Avoid Unsupervised Testosterone or Steroids

Men planning pregnancy should speak with a doctor before using testosterone therapy or anabolic steroids. External testosterone can suppress sperm production.

Can Diabetes Affect Male Fertility?

Diabetes can affect male sexual and reproductive health. Potential complications can involve:

- Erectile function
- Nerve function
- Blood vessels
- Hormonal health
- Overall metabolic health

Men with diabetes who are experiencing difficulty conceiving should discuss the issue with a healthcare professional.

Depending on the circumstances, the doctor may recommend semen analysis, hormone testing or other investigations.

When Should Men Get a Semen Analysis?

A semen analysis may be appropriate when:

- A couple has difficulty conceiving.
- There is a history of male infertility.
- There is a known testicular problem.
- There has been reproductive surgery.
- There is concern about sperm production.
- Fertility treatment is being considered.
- A doctor identifies potential male-factor infertility.

WHO defines infertility as failure to achieve pregnancy after 12 months or more of regular unprotected sexual intercourse.

However, couples do not always need to wait exactly 12 months before discussing fertility.

Earlier evaluation may be appropriate when there are known risk factors or medical concerns.

The female partner's age is also important in determining how quickly a couple should seek evaluation. ASRM recommends evaluation after 12 months when the female partner is under 35 and after 6 months when she is 35 or older, assuming regular unprotected intercourse and no known cause.

Should Men Over 40 Get a Sperm Analysis?

Being over 40 does not automatically mean that a man needs a semen analysis.

However, if a man is trying to conceive and has concerns about fertility, age can be part of the reason to discuss testing with a doctor.

Testing may be particularly relevant when there are additional factors such as:

- Previous fertility problems
- Testicular disease
- Previous surgery
- Hormonal symptoms
- Diabetes
- Obesity
- Medication use
- Previous chemotherapy
- Long-term smoking
- Difficulty conceiving

A semen analysis gives objective information that age alone cannot provide.

What Other Male Fertility Tests May Be Recommended?

A doctor may recommend additional tests depending on the semen analysis and medical history.

Testosterone

Testosterone testing may be useful when hormonal problems are suspected.

FSH

Follicle-stimulating hormone can provide information about the hormonal regulation of sperm production.

LH

Luteinising hormone is involved in reproductive hormone regulation.

Prolactin

Prolactin may be evaluated in selected patients with hormonal or sexual symptoms.

Antisperm Antibodies

These may be investigated in selected situations, although they are not generally recommended as a routine first-line test for every infertile man.

Genetic Testing

Genetic testing may be appropriate for selected men with severe abnormalities in sperm production.

Scrotal Doppler

Ultrasound Doppler imaging can be used to evaluate scrotal structures when clinically indicated.

The important point is that not every patient requires every fertility test.

Testing should be based on the individual's medical history and clinical assessment.

For men searching for male fertility testing in Bengaluru, TruScan Diagnostics provides a dedicated fertility testing pathway.

TruScan's current Fertility Profile – Men & Women lists several male fertility investigations, including:

- Semen analysis
- Total testosterone
- Free testosterone
- LH
- FSH
- Prolactin
- Antisperm antibody testing
- Scrotal Doppler
- Karyotype and selected genetic investigation

The exact combination of investigations should be confirmed with the diagnostic provider and treating doctor because not every test is appropriate for every patient.

The current TruScan page states that its Fertility Profile is available at:

TruScan Diagnostics – JP Nagar 7th Phase
Manvi Arcade, WGHCS Layout, Bengaluru

TruScan Diagnostics – JP Nagar 2nd Phase
213, 9th Cross Road, KSRTC Layout, Bengaluru

TruScan currently states that both centres are open 24 hours for walk-ins and scheduled appointments.

The current package page also states that preparation requirements, including fasting where relevant, depend on the tests included and are confirmed during booking.
This creates a useful diagnostic pathway for men concerned about fertility:
Consultation → Appropriate fertility tests → Semen analysis and/or blood tests → Additional testing where clinically indicated → Doctor interpretation → Further management

---

Why a Fertility Profile Can Be More Useful Than a Single Test

A common mistake is to think that male fertility is simply a question of sperm count.
It is not.
A man may have an acceptable sperm concentration but have another abnormal parameter.
Conversely, one abnormal semen parameter does not necessarily mean that natural conception is impossible.
A broader fertility assessment can help identify whether there are additional issues involving:

- Sperm
- Hormones
- Testicular structures
- Reproductive tract
- Genetics
- Other health factors

This is why a doctor may recommend a combination of tests rather than relying on one number.

Male Fertility and Advanced Paternal Age

Paternal age deserves discussion when men plan to become fathers later in life.

The AUA/ASRM guideline recommends counselling couples with advanced paternal age, defined as 40 years or older in that guideline, about increased risks of adverse health outcomes for offspring.

These risks are generally population-level associations, not predictions about an individual child.

A man should therefore not assume that fatherhood after 40 or 50 is impossible or unsafe.

Instead, couples can discuss:

- Reproductive planning
- General health
- Family history
- Fertility testing
- Genetic counselling where appropriate
- Pregnancy-related risks

with qualified healthcare professionals.

What Does a Male Fertility Evaluation Usually Look Like?

A complete evaluation may involve several stages.

Stage 1: Medical and Reproductive History

The doctor may ask about:

- Previous pregnancies
- Duration of trying to conceive
- Sexual function
- Previous infections
- Surgeries
- Medications
- Lifestyle
- Family history
- Testicular problems

Stage 2: Physical Examination

The doctor may assess the reproductive system and look for signs such as testicular abnormalities or varicocele.

Stage 3: Semen Analysis

A semen analysis provides information about sperm and semen parameters.

Stage 4: Hormonal Testing

Hormone testing may be recommended when clinically indicated.

Stage 5: Imaging

Scrotal imaging may be considered when there is a specific clinical indication.

Stage 6: Genetic or Specialised Testing

Additional testing may be recommended for selected patients.

This structured approach is consistent with the AUA/ASRM recommendation that male infertility assessment include reproductive history and semen

analysis, followed by directed investigations where indicated.

Frequently Asked Questions About Male Fertility and Age

Does male fertility decrease with age?

Some semen parameters can decline with increasing age, but male fertility does not suddenly stop at a particular age.

Does age affect sperm count?

Age can be associated with changes in sperm concentration and total sperm number, but the effect varies significantly between individuals.

Does sperm quality decrease after 40?

Some aspects of sperm quality can change with age. However, actual sperm quality should be assessed through appropriate testing rather than assumed from age alone.

Can a 40-year-old man still have children?

Yes. Men can father children after 40. However, paternal age may be relevant to fertility counselling and reproductive risk assessment.

Should men get a semen analysis after 40?

Not simply because they are over 40. Testing may be appropriate when there is difficulty conceiving, a relevant medical history or another clinical reason.

What does a semen analysis test show?

It can assess semen volume, sperm concentration, total sperm number, motility and morphology, depending on the laboratory's testing protocol.

Can one abnormal semen analysis mean infertility?

No. Semen parameters vary between samples, and doctors may recommend repeat testing or further evaluation.

How long should a man abstain before a semen analysis?

The required abstinence period depends on the laboratory's protocol. Follow the instructions provided by the laboratory performing the test.

What is the difference between sperm count and sperm quality?

Sperm count refers primarily to the number or concentration of sperm. Sperm quality is a broader concept that can include motility, morphology and other characteristics.

Can smoking affect male fertility?

Yes. Tobacco use is associated with adverse reproductive and general health effects.

Can alcohol affect sperm quality?

Excessive alcohol consumption may negatively affect general health and reproductive function.

Does testosterone treatment affect sperm production?

External testosterone can suppress the body's natural reproductive hormone signalling and reduce sperm

production. Men trying to conceive should discuss testosterone treatment with a doctor.

Can diabetes affect male fertility?

Diabetes and its complications can affect sexual and reproductive health. Men with diabetes who have fertility concerns should seek appropriate medical evaluation.

What is a fertility profile?

A fertility profile is a combination of investigations designed to evaluate selected aspects of reproductive health. The exact tests vary by provider and patient.

Does TruScan offer male fertility testing?

Yes. TruScan currently lists a Fertility Profile for Men & Women that includes semen analysis, testosterone-related testing, LH, FSH, prolactin, antisperm antibody testing, scrotal Doppler and selected genetic testing among its male investigations.

Where can I get fertility testing in Bengaluru?

TruScan currently lists its Fertility Profile at its JP Nagar 7th Phase and JP Nagar 2nd Phase centres in Bengaluru. Patients should confirm the current test list, preparation requirements and appointment availability before visiting.

Final Takeaway

Male fertility and age are connected, but age alone does not determine fertility.

Men can remain capable of fathering children well into later adulthood. At the same time, increasing age may be

associated with changes in sperm concentration, motility, morphology, semen characteristics and sperm DNA.
This is why couples should avoid making assumptions based solely on age.
A semen analysis can provide useful information about sperm concentration, motility, morphology and other semen characteristics. If abnormalities are identified, a doctor may recommend repeat testing or additional investigations such as hormonal evaluation, genetic testing or imaging.
The WHO emphasises the importance of standardised semen laboratory procedures, while current WHO infertility guidance supports evidence-based evaluation and treatment of infertility.
For men seeking male fertility testing in Bengaluru, TruScan Diagnostics' current Fertility Profile includes semen analysis and a range of hormonal, reproductive, genetic and imaging investigations, with listed centres in JP Nagar 7th Phase and JP Nagar 2nd Phase.
If you and your partner are planning a pregnancy after 40, experiencing difficulty conceiving, or have concerns about sperm health, speak with a qualified fertility specialist. The right approach is to evaluate the individual—not simply the number on the calendar.
Understanding your reproductive health starts with appropriate evaluation, accurate testing and professional interpretation.

Meta Title: Male Fertility & Age: How Aging Affects Sperm Analysis Results

Meta Description: Does male fertility decline with age? Explore how paternal age impacts sperm count, motility, morphology, DNA integrity, and when to get a semen analysis.

H1 Tag: `<h1>How Age Affects Male Fertility and Sperm Analysis Test Results</h1>`

Blog 4

Understanding Breast Cancer Subtypes: How Molecular Testing Helps Guide Treatment

A breast cancer diagnosis can bring many questions. Patients may hear terms such as ER-positive, PR-positive, HER2-positive, HER2-negative, triple-negative, luminal A, luminal B, Ki-67 and molecular testing during conversations with their doctor or while reading a pathology report.

These terms can initially sound complicated, but they all relate to an important concept: breast cancer is not one single disease.

Different breast cancers can have different biological characteristics. Those characteristics can influence how a tumour behaves and which treatments may be appropriate.

This is why modern breast cancer diagnosis goes beyond simply answering the question, "Is this cancer?"

Doctors may also need to understand:

- **What type of breast cancer is present?**
- **What is the tumour grade?**
- **What is the stage?**
- **Does the tumour have estrogen receptors?**
- **Does it have progesterone receptors?**
- **Is HER2 present?**
- **How actively are the tumour cells dividing?**
- **Are additional molecular tests appropriate?**
- **Could the results influence treatment?**

The National Cancer Institute explains that breast cancer cells are commonly tested for estrogen receptor (ER), progesterone receptor (PR) and HER2, while other biomarkers such as Ki-67 may also provide clinically useful information.

Understanding these tests can help patients make sense of their pathology reports and understand why their oncology team may recommend specific treatment approaches.

What Are Breast Cancer Subtypes?

Breast cancer can be classified in different ways.

Doctors may describe a tumour based on:

- **Histological type**
- **Grade**

- **Stage**
- **Hormone receptor status**
- **HER2 status**
- **Molecular characteristics**
- **Other biomarkers**

When people talk about breast cancer molecular subtypes, they are generally referring to groups of tumours that share certain biological characteristics.

Commonly discussed groups include:

- **Luminal A**
- **Luminal B**
- **HER2-enriched**
- **Basal-like or triple-negative breast cancer**

In everyday clinical practice, doctors may also describe breast cancers using combinations such as:

- **Hormone receptor-positive / HER2-negative**
- **Hormone receptor-positive / HER2-positive**
- **Hormone receptor-negative / HER2-positive**
- **Triple-negative breast cancer**

These classifications are useful because they provide information about tumour biology and can contribute to treatment planning.

Why Is Breast Cancer Subtyping Important?

Two people can both have breast cancer but require very different treatment strategies.

For example, one tumour may depend on hormone signalling, while another may have increased HER2 activity.

Another tumour may not express ER, PR or HER2 at all.

These differences can influence whether treatments such as:

- **Hormone therapy**
- **Chemotherapy**
- **HER2-targeted therapy**
- **Immunotherapy**
- **Other targeted therapies**

may be appropriate.

However, biomarker results do not determine treatment by themselves.

Doctors also consider tumour stage, grade, size, lymph-node involvement, overall health, previous treatment and other clinical information.

What Are ER and PR?

ER stands for estrogen receptor.

PR stands for progesterone receptor.

These are receptors found on cells that can respond to hormones.

Some breast cancer cells contain estrogen and/or progesterone receptors.

When breast cancer cells contain estrogen receptors, the cancer is generally described as ER-positive.

When they contain progesterone receptors, it may be described as PR-positive.

A tumour can therefore have different combinations of ER and PR results.

For example:

- **ER-positive / PR-positive**
- **ER-positive / PR-negative**
- **ER-negative / PR-positive**
- **ER-negative / PR-negative**

The results help doctors understand the biology of the tumour.

What Does ER-Positive Breast Cancer Mean?

ER-positive breast cancer means that the cancer cells have estrogen receptors.

This is clinically important because hormone signalling can contribute to the growth of hormone receptor-positive breast cancers.

For appropriate patients, hormone-based treatment may therefore form an important part of treatment planning.

The National Cancer Institute explains that testing for estrogen and progesterone receptors can help determine whether hormone therapy may be useful.

However, an ER-positive result does not provide the complete treatment plan.

Doctors also need to consider:

- **Tumour stage**
- **Tumour grade**
- **HER2 status**
- **Lymph-node involvement**
- **Other pathological characteristics**
- **Patient health**

- **Previous treatment**

What Does PR-Positive Breast Cancer Mean?

PR-positive means that progesterone receptors have been detected in the tumour cells.

PR results are usually considered alongside ER results.

For example, a cancer that is both ER-positive and PR-positive has a different biological profile from a cancer that is ER-positive but PR-negative.

This information can contribute to the overall classification of the tumour.

What Is HER2?

HER2 stands for human epidermal growth factor receptor 2.

HER2 is a protein involved in cell growth and signalling.

Some breast cancers have increased HER2 expression or other HER2-related abnormalities.

These cancers may be classified as HER2-positive breast cancer.

HER2 testing is therefore an important component of breast cancer pathology.

What Does HER2-Positive Breast Cancer Mean?

A HER2-positive breast cancer has characteristics indicating increased HER2 activity according to established testing criteria.

HER2 status can have major treatment implications because HER2-targeted therapies are available for appropriate patients.

This is an example of how molecular or biomarker testing can contribute directly to personalised cancer treatment.

Instead of treating every breast cancer in exactly the same way, doctors can use tumour biology to help determine which therapies are most appropriate.

Why Is HER2 Testing Done?

HER2 testing helps doctors determine whether a tumour has HER2-related characteristics that could make HER2-directed treatment relevant.

The testing may be performed using laboratory techniques such as:

- **Immunohistochemistry**
- **In situ hybridisation in appropriate cases**

The exact testing approach depends on the pathology situation and established clinical protocols.

The NCI identifies HER2 testing as one of the key biomarker evaluations used in breast cancer.

What Is Triple-Negative Breast Cancer?

Triple-negative breast cancer, commonly abbreviated as TNBC, is generally defined by the absence of:

- **Estrogen receptors**
- **Progesterone receptors**
- **HER2**

In other words:

ER-negative + PR-negative + HER2-negative = Triple-negative breast cancer

This classification is important because hormone-based therapies and HER2-targeted therapies that are used for certain other breast cancers are not applicable in the same way to triple-negative disease.

Treatment therefore requires a different clinical approach.

The NCI specifically defines triple-negative breast cancer based on the absence of ER, PR and HER2.

Is Triple-Negative Breast Cancer More Aggressive?

Triple-negative breast cancers can have different biological and clinical behaviour from hormone receptor-positive breast cancers.

They may grow and spread more rapidly in some patients and have historically had fewer targeted treatment options.

However, treatment options for triple-negative breast cancer have expanded.

Depending on the individual case, treatment may involve combinations of:

- **Surgery**
- **Chemotherapy**
- **Immunotherapy**
- **Radiation**
- **Other systemic treatments**

Treatment should always be determined by the oncology team based on the specific cancer.

What Is Luminal A Breast Cancer?

Luminal A is one of the commonly discussed molecular subtypes.

It is generally associated with characteristics such as:

- **ER-positive status**
- **PR-positive status**

- **HER2-negative status**
- **Lower proliferation**
- **Often lower tumour grade**

Ki-67 may also be relatively low.

Luminal A cancers often have a different biological behaviour from cancers with higher proliferation.

However, classification is not based on one laboratory result.

Doctors consider the complete pathology and clinical picture.

What Is Luminal B Breast Cancer?

Luminal B breast cancers are generally hormone receptor-positive but may show characteristics associated with higher proliferation.

They may be:

- **HER2-negative or HER2-positive**
- **Associated with higher Ki-67**
- **PR-positive or PR-negative**

Compared with luminal A tumours, luminal B cancers may have biological characteristics associated with greater tumour proliferation.

This distinction can contribute to treatment planning.

What Is HER2-Enriched Breast Cancer?

HER2-enriched is a molecular subtype associated strongly with HER2 biology.

However, molecular classification and routine clinical HER2 testing are not necessarily identical concepts.

A patient's pathology report may simply describe HER2 status based on clinical testing.

The treating team interprets the result together with hormone receptor status and other tumour characteristics.

What Is Ki-67?

Ki-67 is a protein associated with cell proliferation.

In simple terms, it can provide information about how actively tumour cells are dividing.

A higher Ki-67 level can indicate a greater proportion of actively proliferating cells.

However, Ki-67 should not be treated as a standalone measure of prognosis or treatment response.

The National Cancer Institute notes that Ki-67 testing may provide information relevant to treatment decisions, but its interpretation needs to be considered alongside other tumour characteristics.

How Is Breast Cancer Molecular Testing Performed?

Biomarker testing is generally performed on tissue containing cancer cells.

The tissue may be obtained through:

- **Core needle biopsy**
- **Other types of biopsy**
- **Surgical removal of the tumour**
- **Tissue obtained during surgery**

The tissue is processed in a pathology laboratory.

Pathologists and laboratory professionals then use appropriate techniques to evaluate the relevant markers.

The NCI explains that biomarker testing for breast cancer is generally performed using tissue obtained during biopsy or surgery.

What Is Immunohistochemistry?

Immunohistochemistry, often abbreviated as IHC, is a laboratory technique used to identify specific proteins within tissue.

It uses antibodies that bind to particular proteins.

IHC can be used in breast cancer testing to assess markers such as:

- **ER**
- **PR**
- **HER2**
- **Ki-67**

The resulting staining pattern and intensity are interpreted according to established criteria.

This information becomes part of the pathology assessment.

What Is Molecular Testing?

Molecular testing refers more broadly to laboratory techniques that examine biological characteristics at the molecular level.

Depending on the cancer and clinical question, molecular testing can investigate:

- **DNA**
- **RNA**
- **Gene expression**
- **Gene alterations**
- **Protein expression**

- **Other molecular characteristics**

Not every breast cancer patient requires extensive molecular testing.

Testing should be selected according to the clinical question.

Is Molecular Testing the Same as Genetic Testing?

No.

This is one of the most important distinctions for patients to understand.

Tumour Biomarker Testing

This examines characteristics of the cancer cells.

Examples include:

- **ER**
- **PR**
- **HER2**
- **Ki-67**

Inherited Genetic Testing

This looks for genetic variants that may be inherited and potentially passed through families.

Examples can include genes such as:

- **BRCA1**
- **BRCA2**
- **Other hereditary cancer-associated genes**

A person can therefore have:

ER-positive breast cancer without having an inherited BRCA mutation.

Likewise, a person may have an inherited genetic risk even before developing cancer.

These tests answer different clinical questions.

Why Does Molecular Testing Matter in Precision Medicine?

Traditional cancer treatment often relied heavily on tumour location and histological appearance.

Modern oncology increasingly combines this information with molecular and biomarker information.

This approach is often described as precision medicine.

The objective is to understand the individual tumour and identify treatments that are more likely to be useful.

For breast cancer, this may mean using ER, PR, HER2 and other biomarkers to guide decisions.

The NCI describes biomarker testing as an important component of precision medicine because biomarkers can help doctors select treatments that may be more appropriate for particular cancers.

Can Breast Cancer Biomarkers Change?

Yes.

Cancer is biologically dynamic.

The characteristics of a tumour at the time of the original diagnosis may not always be identical to those of cancer that returns later.

For example, hormone receptor status can sometimes change between the original tumour and recurrent disease.

The NCI notes that hormone receptor status may change when breast cancer recurs.

Because of this, doctors may sometimes recommend a new biopsy and repeat biomarker testing when cancer returns.

Why Is Pathology Important in Breast Cancer?

Pathology provides the foundation for understanding what type of tumour is present.

A pathology report can contain information about:

- **Histological tumour type**
- **Tumour grade**
- **Tumour size**
- **Lymphovascular involvement**
- **Surgical margins**
- **Lymph-node findings**
- **ER**
- **PR**
- **HER2**
- **Ki-67**
- **Other relevant pathological features**

These findings help the oncology team develop an appropriate treatment strategy.

Pathology is therefore much more than simply confirming that cancer exists.

It helps describe what kind of cancer it is.

Grade vs Stage vs Molecular Subtype

These three concepts are often confused.

What Is Cancer Stage?

Stage describes the extent of cancer in the body.

It can include information about:

- **Primary tumour size**
- **Lymph nodes**
- **Spread to distant organs**

What Is Cancer Grade?

Grade describes characteristics of the cancer cells and how abnormal they appear under microscopic examination.

It can provide information about how aggressively the tumour may behave.

What Is Molecular Subtype?

Molecular or biomarker classification describes biological characteristics such as:

- **ER**
- **PR**
- **HER2**
- **Ki-67**
- **Other molecular markers**

All three can provide different and complementary information.

How Can Biomarker Testing Influence Treatment?

Biomarker testing can help doctors determine whether particular treatment strategies may be relevant.

For example:

Hormone Receptor-Positive Breast Cancer

Hormone therapy may be an important component of treatment for appropriate patients.

HER2-Positive Breast Cancer

HER2-targeted treatment may be considered when clinically appropriate.

Triple-Negative Breast Cancer

Treatment may involve chemotherapy, immunotherapy and/or other approaches depending on the specific situation.

Higher-Proliferation Tumours

Additional characteristics may influence decisions about systemic therapy.

The treatment plan is never determined by a single marker alone.

What Happens After Molecular Testing?

Once pathology and biomarker results are available, the treating team reviews them alongside:

- **Imaging**
- **Cancer stage**
- **Tumour grade**
- **Lymph-node status**

- **Patient age**
- **General health**
- **Menopausal status where relevant**
- **Previous treatment**
- **Family history**
- **Other laboratory findings**

The oncologist may then recommend an appropriate treatment strategy.

Depending on the case, this may involve:

1. **Surgery**
2. **Chemotherapy**
3. **Radiation**
4. **Hormone therapy**
5. **Targeted therapy**
6. **Immunotherapy**
7. **A combination of treatments**
8. **Active surveillance in selected circumstances**

What Is the Role of TruScan Diagnostics in Breast Cancer Testing?

TruScan Diagnostics is relevant to the laboratory and diagnostic component of breast cancer evaluation.

The SEO plan specifically identifies three TruScan service areas for this topic:

- **Pathology**
- **Molecular Biology**
- **Cancer Screening**

These services align naturally with the diagnostic pathway discussed in this article.

A typical pathway may involve:

Clinical consultation → Imaging → Biopsy → Pathology → Biomarker testing → Molecular testing where appropriate → Oncology consultation → Treatment planning

Pathology helps establish the characteristics of the tumour.

Biomarker testing can provide information about ER, PR and HER2.

Additional molecular testing may be considered when there is a specific clinical reason.

Cancer screening, meanwhile, belongs primarily to the earlier detection and preventive side of healthcare.

Patients should confirm with TruScan whether the specific pathology or molecular test recommended by their doctor is currently available.

Breast Cancer Screening vs Diagnostic Testing

These are two different concepts.

Screening

Screening is intended to identify possible disease before symptoms develop or at an earlier stage in appropriate populations.

Depending on age and risk factors, breast cancer screening may involve methods such as mammography.

Diagnostic Testing

Diagnostic testing is performed when there is:

- **A breast lump**
- **Abnormal imaging**
- **Nipple changes**
- **Suspicious findings**
- **Other clinical concerns**

A patient with a new breast lump should not simply wait for their next routine screening appointment.

They should seek appropriate medical evaluation.

What Breast Changes Should Be Checked?

Not every breast change means cancer.

However, new or persistent changes deserve medical attention.

These may include:

- **A new breast lump**
- **Persistent breast thickening**
- **Change in breast size or shape**
- **Skin dimpling**
- **Skin thickening**
- **Nipple inversion**
- **Unusual nipple discharge**
- **Persistent changes around the nipple**
- **An abnormal screening result**

- **Swelling around the breast or underarm**

Early evaluation can determine whether the change requires imaging, biopsy or further testing.

Who May Need More Detailed Molecular or Genetic Testing?

Not every breast cancer patient requires every available molecular test.

Additional testing may be considered based on:

- **Tumour characteristics**
- **Cancer stage**
- **Family history**
- **Age at diagnosis**
- **Previous cancers**
- **Treatment planning**
- **Recurrence**
- **Suspected inherited cancer risk**

A medical oncologist, breast surgeon, genetic counsellor or other appropriate specialist can determine whether additional testing is useful.

Frequently Asked Questions About Breast Cancer Subtypes

What are the molecular subtypes of breast cancer?

Commonly discussed molecular subtypes include luminal A, luminal B, HER2-enriched and basal-like/triple-negative breast cancer. Clinical classification also commonly uses hormone receptor and HER2 status.

What are ER, PR and HER2?

ER and PR are hormone receptors found on some breast cancer cells. HER2 is a protein involved in cell growth and signalling. Testing these markers helps characterise breast cancer.

What does ER-positive breast cancer mean?

It means the tumour cells contain estrogen receptors. Hormone receptor status can influence whether hormone-based treatment is appropriate.

What does PR-positive breast cancer mean?

It means progesterone receptors are detected in the tumour cells. PR is generally interpreted alongside ER and other tumour characteristics.

Why is HER2 testing done?

HER2 testing determines whether a breast tumour has HER2-related characteristics that may influence treatment decisions, including eligibility for HER2-targeted treatment.

What does HER2-positive breast cancer mean?

It means the tumour meets established criteria for HER2 positivity. HER2-positive cancers may be eligible for HER2-targeted therapies depending on the complete clinical situation.

What is triple-negative breast cancer?

Triple-negative breast cancer is generally negative for estrogen receptors, progesterone receptors and HER2.

Is triple-negative breast cancer treatable?

Yes. Treatment options are available, although the treatment strategy differs from that used for hormone receptor-positive or HER2-positive disease. The appropriate treatment depends on the individual cancer.

What is luminal A breast cancer?

Luminal A generally describes an ER-positive, HER2-negative tumour with relatively lower proliferation and often lower grade, although exact classification can vary.

What is luminal B breast cancer?

Luminal B generally refers to hormone receptor-positive cancers with characteristics associated with higher proliferation than luminal A cancers.

What is Ki-67?

Ki-67 is a marker associated with cell proliferation. It can provide information about how actively tumour cells are dividing but should not be interpreted independently.

Is molecular testing the same as genetic testing?

No. Molecular/biomarker testing examines characteristics of tumour cells, while inherited genetic testing looks for genetic variants that may be passed through families.

Can breast cancer biomarkers change?

Yes. Biomarker characteristics can sometimes differ when cancer recurs, which is why repeat testing may be considered in selected cases.

Does every breast cancer patient need molecular testing?

No. Testing should be selected according to the clinical question, tumour characteristics and treatment needs.

Where can breast cancer pathology testing be done in Bengaluru?

Patients can discuss the required investigation with their treating doctor and choose an appropriate diagnostic laboratory. TruScan Diagnostics identifies Pathology, Molecular Biology and Cancer Screening among its relevant service areas.

Final Takeaway

Understanding breast cancer subtypes can make a diagnosis and pathology report easier to understand.

ER, PR, HER2 and other biomarkers provide important information about the biology of a breast tumour. These results can contribute to cancer classification and help doctors determine which treatment approaches may be appropriate.

However, no single biomarker should be interpreted in isolation.

A complete breast cancer assessment may involve:

Pathology + Biomarker Testing + Molecular Testing where appropriate + Imaging + Stage + Grade + Clinical Evaluation

This combination allows the oncology team to develop a treatment strategy based on the individual patient's cancer rather than relying on a one-size-fits-all approach.

For patients undergoing breast cancer evaluation, TruScan Diagnostics has relevant service areas including Pathology, Molecular Biology and Cancer Screening, which align with different stages of the diagnostic journey.

If you notice a new breast lump, persistent breast change or an abnormal screening result, consult a qualified healthcare professional promptly. If breast cancer has already been diagnosed, ask your treating team which pathology, biomarker and molecular tests are appropriate for your specific case.

Accurate diagnosis is the foundation of informed cancer care—and understanding the biology of a tumour can help doctors make more personalised treatment decisions.

Meta Title: Breast Cancer Subtypes & Molecular Testing: A Guide to Pathology

Meta Description: Understand breast cancer subtypes, ER/PR status, HER2, Ki-67, and molecular testing. Learn how biomarker pathology guides personalized cancer treatment.

H1 Tag: <h1>Understanding Breast Cancer Subtypes: How Molecular Testing Helps Guide Treatment</h1>