■ India 28 States: Plant Pathology & Full Field Control Guide

Comprehensive Agronomy, 3-Tier Plant Anatomy, Epidemiology, Climate Drivers & State-Wise Pesticides Guide

---

Executive Summary: This technical guide integrates precision agricultural robotics, plant anatomy, and meteorological epidemiology across all 28 states of the Republic of India. Designed for closed-loop autonomous field management (FarmOS), it establishes targeted interventions matching biological ingress: **Worker Drone Swarms** for upper foliar canopy infections, and **Smart Solenoid Drip Valves** for subterranean root-zone bio-fertigation. Furthermore, it strictly enforces Government of India CIBRC regulations, blacklisting hazardous chemicals while establishing validated organic bio-pesticide protocols.

1. Three-Tier Plant Anatomy & Actuator Interventions

Plants possess three primary physiological tiers. In an autonomous agricultural system, each tier requires a distinct diagnostic and actuation methodology:

Plant Tier	Primary Functions & Organs	Vulnerable Pathogens	Automated Actuation Target
I. Bottom Part Root System	• Soil anchoring & moisture uptake • Mineral nutrient absorption (N, P, K) • *Taproot* (Mango, Neem) vs *Fibrous* (Paddy, Wheat)	• Root Rot (*Rhizoctonia*) • Damping-off (*Pythium*) • Bacterial Wilt (*Ralstonia*) • Fusarium Panama Wilt	**Smart Solenoid Drip Valves** • Subterranean root drenching • *Trichoderma viride* / *Pseudomonas* bio-fertigation
II. Middle Part Stem / Trunk	• Structural canopy elevation • Vascular transport: *Xylem* (water) and *Phloem* (sugars) • Woody bark vs herbaceous culms	• Sugarcane Red Rot • Rice Sheath Blight • Gummosis & Stem Cankers • Stem Borer larval tunnels	**Low-Canopy Drone Swarm** • Oblique-angle spray nozzles • Directed stem collar washing • Targeted biopesticide drench
III. Top Part Leaves & Canopy	• Photosynthetic carbon fixation • Stomatal transpiration cooling • Gas exchange & reproduction (flowers, fruit, seeds)	• Rusts (*Puccinia*) • Blights (*Alternaria* / *Phytophthora*) • Powdery Mildew • Leaf Curl Viruses (Vectors)	**Worker Drone Swarm** • Aerial micro-atomized foliar spray (80–150 µm droplets) • Maximum upper/lower leaf coverage

2. All 28 States of India — Agriculture & Pesticides Analysis

Comprehensive mapping of India's 28 states, detailing major economic crops, commonly registered chemical treatments, and statutory organic mandates (e.g. Sikkim 100% Organic State):

S.No	State	Zone	Major Crops	Commonly Used Pesticides / Regimen
1	**Kerala**	South	Coconut, Rubber, Tea, Cardamom, Black Pepper, Paddy	Copper Oxychloride, Chlorpyrifos, Carbendazim
2	**Tamil Nadu**	South	Paddy, Sugarcane, Groundnut, Banana, Cotton, Maize	Imidacloprid, Cartap Hydrochloride, Chlorantraniliprole, Mancozeb
3	**Andhra Pradesh**	South	Chilli, Cotton, Paddy, Tobacco, Maize	Acephate, Thiamethoxam, Profenofos, Diafenthiuron
4	**Telangana**	South	Cotton, Paddy, Maize, Chilli, Turmeric	Fipronil, Spinetoram, Monocrotophos
5	**Karnataka**	South	Coffee, Ragi, Paddy, Sugarcane, Sunflower, Pigeon Pea	Hexaconazole, Propiconazole, Cypermethrin
6	**Goa**	West	Cashew, Coconut, Paddy, Arecanut	Bordeaux Mixture, Carbaryl, Glyphosate
7	**Maharashtra**	West	Sugarcane, Cotton, Soybean, Grapes, Onion	Dimethoate, Sulphur, Emamectin Benzoate, Lambda-cyhalothrin
8	**Gujarat**	West	Cotton, Groundnut, Castor, Cumin	Acetamiprid, Diafenthiuron, Profenofos

9	**Rajasthan**	North-West	Mustard, Pearl Millet (Bajra), Green Gram, Cumin, Coriander	Malathion, Quinalphos, Dimethoate
10	**Punjab**	North	Wheat, Paddy (Basmati), Cotton, Potato	Pretilachlor, Tricyclazole, Glyphosate, Thiamethoxam
11	**Haryana**	North	Wheat, Paddy, Mustard, Sugarcane, Cotton	Chlorpyrifos, Carbendazim, Imidacloprid
12	**Himachal Pradesh**	North	Apple, Potato, Maize, Vegetables	Mancozeb, Diphos, Dodine, Lime Sulphur
13	**Uttarakhand**	North	Paddy, Wheat, Sugarcane, Fruits, Pulses	Copper Fungicides, Cypermethrin, Carbaryl
14	**Uttar Pradesh**	North	Sugarcane, Wheat, Potato, Mustard, Mango	Carbofuran, Chlorpyrifos, Mancozeb
15	**Bihar**	East	Paddy, Maize, Wheat, Jute, Litchi	Phorate, Imidacloprid, Mancozeb
16	**Jharkhand**	East	Paddy, Pulses, Maize, Vegetables	Chlorpyrifos, Carbendazim, Cypermethrin
17	**West Bengal**	East	Paddy, Jute, Tea, Potato, Mango	Cartap Hydrochloride, Hexaconazole, Neem Formulations
18	**Odisha**	East	Paddy, Pulses, Groundnut, Cashew, Sugarcane	Triazophos, Imidacloprid, Carbendazim
19	**Chhattisgarh**	Central	Paddy, Maize, Oilseeds, Vegetables	Cartap, Chlorpyrifos, Tricyclazole
20	**Madhya Pradesh**	Central	Soybean, Wheat, Chickpea, Mustard	Chlorantraniliprole, Thiamethoxam, Tebuconazole
21	**Sikkim** [100% Organic]	North-East	Cardamom, Ginger, Orange, Buckwheat	Bio-pesticides (Neem oil, Trichoderma, Vermicompost)
22	**Assam**	North-East	Tea, Paddy, Jute, Mustard	Copper Oxychloride, Ethion, Propargite
23	**Arunachal Pradesh** [100% Organic]	North-East	Paddy, Maize, Orange, Pineapple, Ginger	Neem Oil, Copper Hydroxide, Bio-fungicides
24	**Nagaland** [100% Organic]	North-East	Paddy, Maize, King Chilli, Cardamom	Organic Bio-pesticides, Lime Sulphur
25	**Manipur**	North-East	Paddy, Maize, Mustard, Fruits	Neem Formulations, Carbendazim
26	**Mizoram** [100% Organic]	North-East	Paddy, Ginger, Turmeric, Bamboo, Fruits	Bio-pesticides, Organic Compost Controls
27	**Tripura**	North-East	Paddy, Rubber, Tea, Pineapple, Bamboo	Copper Oxychloride, Chlorpyrifos
28	**Meghalaya** [100% Organic]	North-East	Paddy, Potato, Lakadong Turmeric, Ginger	Organic Bio-pesticides, Bordeaux Mixture

3. Disease Etiology: Fungi vs. Bacteria vs. Viruses & Climate Dynamics

Plant diseases follow distinct anatomical infection routes (Bottom vs. Top) governed critically by microclimatic triggers:

A. Fungal Diseases (Spore Pathogens — ~70% of Plant Diseases)

• **Bottom-Level Ingress (Soil/Roots):** Pathogens like *Rhizoctonia solani*, *Pythium*, and *Fusarium oxysporum* survive as sclerotia or chlamydospores in soil debris. They infect fibrous root tips and move upward into vascular xylem, causing damping-off, root rot, and Panama wilt.
• **Top-Level Ingress (Canopy):** Airborne spores (conidia, urediniospores) of *Puccinia* (Rust) and *Alternaria* (Early Blight) land on upper foliage. Spores germinate in microscopic moisture droplets, penetrating stomata.
• **Climate Master Role:** Fungi require high relative humidity (**RH > 80%**) and 4–8 hours of uninterrupted **Leaf Wetness Duration (LWD)** at moderate temperatures (**18°C–26°C**). Light wind currents disperse spores across acreage.

B. Bacterial Diseases (Water-Borne Vascular Invaders)

• **Bottom-Level Ingress:** *Ralstonia solanacearum* (Bacterial Wilt) and *Pectobacterium* (Soft Rot) enter root wounds caused by nematodes or tilling, clogging the xylem with extracellular slime.
• **Top-Level Ingress:** *Xanthomonas oryzae* (Rice Bacterial Blight) enters leaf tips via hydathodes or stomata, spreading via water droplets.
• **Climate Master Role:** Bacteria flourish in **warm, sultry climates (28°C–35°C)** paired with driving monsoon rains. High windstorms cause micro-abrasions on leaves that provide instant entry portals for bacterial ooze.

C. Viral Diseases (Vector-Borne Systemic Parasites)

• **Top-Level Ingress via Vectors:** Viruses cannot penetrate plants unaided. Sap-sucking insect vectors (Whiteflies, Thrips, Aphids) feed on succulent terminal shoots and tender crown leaves, injecting viral particles directly into the phloem sieve stream. Examples include Chilli Leaf Curl Virus (ChLCV) and Cotton Leaf Curl Virus (CLCuV).
• **Climate Master Role: Hot, dry spells (32°C–40°C)** drastically accelerate insect vector reproduction cycles. Drought conditions force insects to feed aggressively on crops to obtain water, causing exponential viral epidemic spread.

4. CIBRC Safety Compliance: Banned Chemicals & Organic Alternatives

The Central Insecticide Board & Registration Committee (CIBRC) of India strictly regulates hazardous pesticides. The PhytoScan system enforces an automated blacklist and prescribes certified bio-pesticide alternatives:

Chemical Name	Legal Regulatory Status	Hazard Class & Health Impact	Approved Bio-Pesticide Alternative
Endosulfan	BANNED Supreme Court Order	Extremely Hazardous Organochlorine; endocrine disruptor, neurotoxin, bio-accumulative.	**Neem Oil (Azadirachtin 10000 ppm)** + *Beauveria bassiana*
Phorate	BANNED / RESTRICTED S.O. 3951(E)	WHO Class Ia; extreme mammalian oral toxicity; rapid cholinesterase inhibitor.	**Neem Cake (250 kg/ha)** + *Trichoderma viride* soil incorporation
Methyl Parathion	BANNED CIBRC Gazette	WHO Class Ia; lethal toxicity to farmers, non-target pollinators, and wildlife.	**Panchagavya foliar spray (3%)** + *Bacillus thuringiensis*
Dichlorvos (DDVP)	BANNED / PHASED OUT	WHO Class Ib; high vapor toxicity; suspected human carcinogen.	**Dashparni Ark** or Neem Seed Kernel Extract (NSKE 5%)
Alachlor	BANNED S.O. 2486(E)	Class II Carcinogenic herbicide; rapid groundwater leaching.	**Organic straw mulching** & manual rotary weeding

5. Autonomous Precision Field Architecture (Sense → Zone → Actuate)

The integrated PhytoScan FarmOS architecture achieves 100% field control through a closed-loop cyber-physical pipeline:

1. **Multi-Modal Sensing:** Master Scout Drones conduct boustrophedon aerial surveys while ground IoT sensors monitor soil moisture (%), pH, and N-P-K levels in real time.
2. **Severity Indexing:** Computer vision models quantify necrotic surface area into *None (<5%)*, *Mild (5–20%)*, *Moderate (20–50%)*, and *Severe (>50%)*.
3. **Meteorological Interlock:** An automated safety interlock continuously samples wind speed. If wind exceeds **15 km/h** or rain is detected, aerial drone spraying is automatically halted to prevent off-target chemical drift under CIBRC standards.
4. **Dual Actuation Execution:** Top foliar outbreaks trigger precision Worker Drone Swarm spraying; soil moisture deficits and root rot trigger Smart Solenoid Drip Valves for targeted subterranean fertigation.
5. **Autonomous Navigation Export:** Directly exports standardized MAVLink / QGroundControl (.waypoints) flight files for physical deployment on Pixhawk, DJI Agras, and ArduPilot agricultural aircraft.