

## Subject: Engineering Mathematics

DPP-03

## Chapter: Vector Calculus

Topic: Divergence & Curl of Vector Function , Line, surface & Volume  
Integral

- Curl of vector  $\vec{v}(x, y, z) = 2x^2\hat{i} + 3z^2\hat{j} + y^3\hat{k}$  at  $x = y = z = 1$  is  
 (a)  $-3\hat{i}$  (b)  $3\hat{i}$   
 (c)  $3\hat{i} - 4\hat{j}$  (d)  $3\hat{i} - 6\hat{k}$
- If  $\vec{r} = x\hat{a}_x + y\hat{a}_y + z\hat{a}_z$  and  $|\vec{r}| = r$ , then  $\text{div}(r^2\nabla(\ln r)) = \underline{\hspace{1cm}}$ .
- A vector  $\vec{P}$  is given by  $\vec{P} = x^3y\vec{a}_x - x^2y^2\vec{a}_y - x^2yz\vec{a}_z$ . Which one of the following statements is TRUE?  
 (a)  $\vec{P}$  is solenoidal, but not irrotational  
 (b)  $\vec{P}$  is irrotational, but not solenoidal  
 (c)  $\vec{P}$  is neither solenoidal nor irrotational  
 (d)  $\vec{P}$  is both solenoidal and irrotational
- The velocity field of an incompressible flow is given by  
 $V = (a_1x + a_2y + a_3z)\hat{i} + (b_1x + b_2y + b_3z)\hat{j} + (c_1x + c_2y + c_3z)\hat{k}$   
 and  $a_1 = 2$  &  $c_3 = -4$ . The value of  $b_2$  is \_\_\_\_\_.
- $\nabla \times \nabla \times P$  (where  $P$  is a vector) is equal to  
 (a)  $P \times \nabla \times P - \nabla^2 P$   
 (b)  $\nabla^2 P + \nabla(\nabla \times P)$   
 (c)  $\nabla^2 P + \nabla \times P$   
 (d)  $\nabla(\nabla \cdot P) - \nabla^2 P$
- The curl of vector  $A = e^{xy}\hat{i} + \sin xy\hat{j} + \cos^2 xz\hat{k}$  is  
 (a)  $ye^{xy}\hat{i} + x \cos xy\hat{j} - 2x \sin 2xz\hat{k}$   
 (b)  $z \sin 2xz\hat{i} + (y \cos xy - xe^{xy})\hat{k}$   
 (c)  $z \sin 2xz\hat{i} + (x \cos xy - xe^{xy})\hat{k}$   
 (d)  $xye^{xy}\hat{i} + xy \cos xy\hat{j} - 2xz \sin 2xz\hat{k}$
- If  $A = (3y^2 - 2z)\hat{i} - 2x^2z\hat{j} + (x + 2y)\hat{k}$ , the value of  $\nabla \times \nabla \times A$  at  $P(-2, 3, -1)$  is  
 (a)  $-(6\hat{i} + 4\hat{j})$  (b)  $8(\hat{i} + \hat{j})$   
 (c)  $-8(\hat{i} + \hat{j})$  (d)  $0$
- The directional derivative of function  $\Phi = xy + yz + zx$  at point  $P(3, -3, -3)$  in the direction toward point  $Q(4, -1, -1)$  is  
 (a)  $-3$  (b)  $1$   
 (c)  $-2$  (d)  $0$
- The maximum value of the directional derivative of the function  $\phi = 2x^3 + 3y^2 + 5z^2$  at a point  $(1, 1, -1)$  is  
 (a)  $10$  (b)  $-4$   
 (c)  $\sqrt{152}$  (d)  $152$
- The grad.  $\nabla \times A$  of a vector field  $A = x^2y\hat{i} + y^2z\hat{j} - 2xz\hat{k}$  is  
 (a)  $2xy + 2yz - 2x$   
 (b)  $x^2y + y^2z - 2xz$   
 (c)  $2x^2y + 2y^2z - 2xz$   
 (d)  $0$

## Answer Key

1. (a)
2. (3)
3. (a)
4. (2)
5. (d)

6. (b)
7. (a)
8. (c)
9. (c)
10. (d)



Any issue with DPP, please report by clicking here:- <https://forms.gle/t2SzQVvQcs638c4r5>

For more questions, kindly visit the library section: Link for web: <https://smart.link/sdfez8ejd80if>



PW Mobile APP: <https://smart.link/7wwosivoicgd4>