


[HOME](#) | [SEARCH](#) | [SITE MAP](#)
[GEO Publications](#) | [FAQ](#) | [MIAME](#) | [Email GEO](#)
[NCBI](#) > [GEO](#) > **Accession Display** [?](#)

 Not logged in | [Login](#) [?](#)

 Scope: Format: Amount: GEO accession:
Series GSE62120
[Query DataSets for GSE62120](#)

Status	Public on Mar 01, 2015
Title	Time dynamics of quantitative protein and mRNA levels reveals extensive translational regulation after stress
Platform organisms	Schizosaccharomyces pombe ; Saccharomyces cerevisiae
Sample organism	Schizosaccharomyces pombe
Experiment type	Expression profiling by array
Summary	<p>Eukaryotic cells are constantly challenged by the presence of reactive oxygen species, which play an important role in aging and human disease progression. In particular, acute oxidative stress can lead to extensive damage to cellular DNA, proteins, and lipids and can trigger a response that remodels the transcriptional and translational state of the cell. Although a number of previous studies have profiled the relative changes in mRNA and protein and more studies revealing the dynamics of transcription and translation in response to stress are starting to emerge, a quantitative view of this response has been lacking. Here, we have applied quantitative methods to characterize the time dynamics of mRNA and protein levels in the oxidative stress response of the fission yeast <i>Schizosaccharomyces pombe</i>, which has allowed us to perform dynamic modeling of responsive genes in units of copies per cell. Analysis of the resulting time dynamics provided a new genome-wide view of the scale, timing and rates of transcription and translation in the transient response. The majority of dynamic genes were observed to be responsive in their mRNA or protein levels alone implying extensive translational regulation. Nevertheless, modeling of genes with responsive mRNA and protein levels showed that protein levels could, in a majority of these cases, be accurately predicted with constant translation and decay rates while a minority benefited from explicit translation delay parameters. A number of independent features, e.g. measures of codon bias, ribosome occupancy, etc., were found to be less correlated to maximally perturbed protein levels than steady-state levels. Codon bias measures were more correlated than mRNA levels to quantitative protein levels at both perturbed and unperturbed states. Measures of translation activity, on the other hand, were only significantly correlated at steady state.</p>
Overall design	In total 32 samples: 11 for stressed time series R1, 11 for stressed time series R2, 5 for control time series C1 and 5 for control time series C2
Contributor(s)	Papadakis MA , Schmidt A , Marsellach FX , De Masi F , Pedersen HK , Slodkiewicz G , Beyer A , Bähler J , Aebersold R , Workman CT
Citation missing	<i>Has this study been published? Please login to update or notify GEO.</i>
Submission date	Oct 07, 2014
Last update date	Mar 02, 2015
Contact name	Federico De Masi
E-mail	fdemasi@cbs.dtu.dk
Organization name	Technical University of Denmark
Department	Institute of Systems Biology
Lab	Regulatory Genomics
Street address	Kemitorvet 208
City	Lyngby

Samples (32) [+ More...](#)

GSM1519986	R1 Stress induced T=0
GSM1519987	R1 Stress induced T=5
GSM1519988	R1 Stress induced T=10

BioProject [PRJNA263246](#)

SOFT formatted family file(s)
MINiML formatted family file(s)
Series Matrix File(s)

SOFT [?](#)
MINiML [?](#)
TXT [?](#)

Supplementary file	Size	Download	File type/resource
GSE62120_RAW.tar	30.6 Mb	(http) (custom)	TAR (of CEL)
<i>Raw data provided as supplementary file</i>			
<i>Processed data included within Sample table</i>			