






	$p(stop)$
①	0.04
②	0.08
③	0.28
④	0.47
⑤	0.93
⑥	1.00

	$\mathcal{X}$	①	②	③	④	⑤
①	✓					
②	✓	✓				
③	✓	✓				
④	✓	✓		✓		
⑤	✓	✓	✓		✓	
⑥	✓	✓	✓		✓	✓

Problem $\mathcal{X}$: A group needs 30 bottles of water for a trip. **Store A** sells packs of 6 bottles for \$4 each. **Store B** sells packs of 10 bottles for \$7 each, but gives a \$3 discount if the total before discount is at least \$20. Which store is cheaper, and how much cheaper is it?




① The group needs at least 30 bottles, so each store must provide enough full packs to reach 30 bottles.




② **Store A** needs $30 \div 6 = 5$ packs, costing $5 \times 4 = 20$ dollars.



③ **Store B** needs $30 \div 10 = 3$ packs, with a pre-discount cost of $3 \times 7 = 21$ dollars.



④ Since Store B's pre-discount cost is at least \$20, the \$3 discount applies, so **Store B** costs $21 - 3 = 18$ dollars.



⑤ Comparing the two totals, Store A costs \$20 and Store B costs \$18.



⑥ Therefore, Store B is cheaper by $20 - 18 = 2$ dollars.

