

Name: Gonzalo Ortega Carpintero

Exercise 1.

Proof. Let X be a set, and let U_i for $i \in I$ be topological spaces, where $X = \bigcup U_i$. Let τ_i be the topology of each U_i . For every $i, j \in I$, the topologies τ_i, τ_j on U_i and U_j restricted to $U_i \cap U_j$ agree. A definition of restricted topology, also known as the relative topology can be found in Section 6 of [2]. We can define a topology over X by considering the union of all open sets of all U_i 's. That is

$$\mathcal{T} = \{V \subseteq X : V \cap U_i \in \tau_i, \forall i \in I\}.$$

Let's check that \mathcal{T} is actually a topology. We have $\emptyset \in \mathcal{T}$ as for all $i \in I$, $\emptyset \cap U_i = \emptyset \in \tau_i$. The same happens with X , as for all $i \in I$, $X \cap U_i = U_i \in \tau_i$. Therefore $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$.

Suppose $V_1, V_2 \in \mathcal{T}$, then

$$(V_1 \cap V_2) \cap U_i = (V_1 \cap U_i) \cap (V_2 \cap U_i) \in \tau_i,$$

as $V_1 \cap U_i$ and $V_2 \cap U_i$ are open sets of U_i , and the intersection of open sets is open. Thus $V_1 \cap V_2 \in \mathcal{T}$.

In the same manner, if $\{V_\alpha\}_{\alpha \in A}$ is a collection of sets of \mathcal{T} , we have

$$\left(\bigcup_{\alpha} V_{\alpha} \right) \cap U_i = \bigcup_{\alpha} (V_{\alpha} \cap U_i) \in \tau_i$$

as each $V_{\alpha} \cap U_i$ is open in U_i . We have then $\bigcup_{\alpha} V_{\alpha} \in \mathcal{T}$, and we can conclude that \mathcal{T} is a topology of X .

We have proved existence, let's prove now the uniqueness of \mathcal{T} . Suppose we have two topologies \mathcal{T}_1 and \mathcal{T}_2 of X that restrict to the topology of each U_i . Let τ_i be the restricted topology of \mathcal{T}_1 on each U_i and τ_j be the restricted topology of \mathcal{T}_2 on each U_j . And let τ'_i, τ'_j be the restricted topologies of τ_i and τ_j on $U_i \cap U_j$. We then have that

$$V \in \mathcal{T}_1 \Leftrightarrow V \cap U_i \in \tau_i \Leftrightarrow V \cap U_i \cap U_j \in \tau'_i \underset{\text{hypothesis}}{\Leftrightarrow} V \cap U_i \cap U_j \in \tau'_j \Leftrightarrow V \cap U_j \in \tau_j \Leftrightarrow V \in \mathcal{T}_2.$$

Therefore, $\mathcal{T}_1 = \mathcal{T}_2$, so there is exactly one topology on X that restricts to the topology on each U_i . \square

Exercise 3.

If $f \in K[x_0, x_1, x_2]$ be a non-constant homogeneous polynomial and let $f = \prod_{i=1}^n g_i^{m_i}$ be the decomposition of f in irreducible polynomials.

(a)

Proof.

(\Rightarrow) If $V(f)$ is irreducible, suppose that we can not write f as a power of a irreducible polynomial. Then, there exist $g_1, g_2 \in K[x_0, x_1, x_2]$, with $g_1 \neq g_2$, such that $f = g_1 g_2$. But then we have $V(f) = V(g_1 g_2) = V(g_1) \cup V(g_2)$ as seen in Remark 3.9 of [1]. This is a contradiction as $V(f)$ is irreducible so it need to be $f = g^m$ for some g irreducible.

(\Leftarrow) If $f = g^m$, with g irreducible, then we have $V(f) = V(g^m) = V(g)$. $V(g)$ needs to be irreducible, because if not it could be expressed as the union of two curves $V(g_1)$ and $V(g_2)$, having $V(g) = V(g_1) \cap V(g_2) = V(g_1 g_2)$, and it would be $g = g_1^m g_2^m$ contradicting the fact of g being irreducible. Thus, $V(g^m)$ is irreducible. \square

(b)

Proof. For $n = 1$, $f = g_1^{m_1}$, so $g_1^{m_1}$ is homogeneous and so g_1 , as f is homogeneous. Suppose $h = \prod_{i=1}^{n-1} g_i^{m_i}$ homogeneous. If g_n was not homogeneous, $g_n^{m_n}$ neither would be, and the product $hg_n^{m_n}$ would not be homogeneous as the product of a homogeneous and a non homogeneous polynomials can not be homogeneous. But $hg_n^{m_n} = \prod_{i=1}^n g_i^{m_i} = f$ contradicting the fact of f been homogeneous. Thus g_n must be homogeneous and, because of induction, each g_i is homogeneous.

As each g_i is an homogeneous irreducible polynomial, because of (a), $V(g_i)$ is a irreducible curve. Therefore, $V(f) = \cup_i V(g_i)$ is a decomposition in irreducible curves of $V(f)$. \square

Exercise 4.

Proof. Let $I < K[x_0, x_1, x_2]$ be a homogeneous ideal. Let $K[x_0, x_1, x_2]/I$ be finite dimensional. If $V(I) \neq \emptyset \subset \mathbb{P}^2$ then $\exists p = [p_0 : p_1 : p_2] \in \mathbb{P}^2$ such that $\forall f \in I, f(p) = 0$. Without lose of generality we can asume $p = [1 : p_1 : p_2]$ (the argument will follow analogously choosing any other coordinate to be different from 0). Thus $\forall n \in \mathbb{N}, x_0^n \notin I$.

Therefore we have an infinite family $\{x_0^n + I\}_{n \in \mathbb{N}}$ of elements of $K[x_0, x_1, x_2]/I$. If the family elements were linearly dependent it would exist $n_i \in \mathbb{N}$ and a finite family of indices $J \subset \mathbb{N}$ such that $x_0^{n_i} = \sum_{j \in J} a_j(x_0^{n_j} + I)$. Let $g = x_0^{n_i} - \sum_{j \in J} a_j(x_0^{n_j} + I) \in I$. Because of the equivalences of the homogeneous ideal definition seen in class, we have $g_i \in I$ for each homogeneous part g_i of g . In particular $x_0^{n_i} \in I$ forming a contradiction. Thus, $\{x_0^n + I\}_{n \in \mathbb{N}}$ is a linearly independent infinite family of $K[x_0, x_1, x_2]/I$ so the quotient can not be infinite dimensional contradicting the exercise hypothesis. It then need to be $V(I) = \emptyset$. \square

References

- [1] Andreas Gathmann, *Plane Algebraic Curves*, Class Notes RPTU Kaiserslautern 2023.
- [2] Stephen Willard, *General Topology*, Addison-Wesley 1970.