

Universidad Autónoma de Madrid

FINAL MASTER THESIS

STRUCTURE AND STABILITY THEOREMS
IN TOPOLOGICAL DATA ANALYSIS
DRAFT

Author:

Gonzalo Ortega Carpintero

Tutor:

Manuel Mellado Cuerno

13/03/2025

Abstract

This is a draft version of a in-work Master's thesis on TDA, compiled for the application to the PhD position in area of topological data analysis in VU University.

Key words

Contents

| | | |
|----------|--|-----------|
| 1 | Preliminaries | 1 |
| 2 | Structure Theorem | 2 |
| 2.1 | Structure theorem for finitely generated modules over a principal ideal domain | 2 |
| 2.2 | Structure theorem for persistence diagrams | 2 |
| 3 | Interleaving Stability Theorem | 5 |
| 4 | Edelsbrunner & Harer’s (Hausdorff) Stability Theorem | 15 |
| 4.1 | Hausdorff Stability | 16 |
| 4.2 | Bottleneck Stability | 19 |
| 5 | Gromov-Hausdorff’s Stability Theorem | 23 |
| 6 | Vectorizations’ Stability Theorems | 24 |
| 6.1 | Persistence landscapes | 24 |
| 6.2 | Persistence images | 24 |
| 6.3 | Euler curves | 24 |
| | Bibliography | 24 |

1

Preliminaries

The contents of this chapter are based on [\[4\]](#), [\[5\]](#) and [\[6\]](#).

Definition 1.0.1 (Graded ring).

Definition 1.0.2 (Graded ideal).

Definition 1.0.3 (Graded module).

Definition 1.0.4 (Persistence module, finite type).

Definition 1.0.5 (Module morphism, shift).

Definition 1.0.6 (Interval module).

Definition 1.0.7 (Direct sum of persistence modules).

Definition 1.0.8 (Barcode).

Definition 1.0.9 (δ -interleaving modules).

Definition 1.0.10 (Interleaving distance).

Definition 1.0.11 (Multiset matching).

Definition 1.0.12 (δ -matching barcodes).

Definition 1.0.13 (Bottleneck distance).

2

Structure Theorem

2.1 Structure theorem for finitely generated modules over a principal ideal domain

2.2 Structure theorem for persistence diagrams

The Structure Theorem for persistence modules is referred to as the “first miracle” of persistence homology [4]. This algebraic property allows to express a persistence module of finite type as a direct sum of finitely many interval modules.

Its proof requires the algebraic structure theorem for finitely generated modules over a principal domain. Due to the lengthy concepts needed to prove it, we will refer to it as a well known fact. An introduction to module theory and a detailed proof of the theorem of Fact 2.2.1 can be found at [3, Chapter IV].

Fact 2.2.1 (Structure theorem for finitely generated modules over a principal ideal domain). *[3, Chapter IV, Theorem 6.12] Let M be a finitely generated module over a principal ideal domain R . There exist a finite sequence of proper ideals $(d_1) \supseteq (d_2) \supseteq \cdots \supseteq (d_n)$*

such that

$$M \cong \bigoplus_{i=1}^n R/(d_i).$$

In addition to Fact 2.2.1, we will use the following simple algebraic statement.

Proposition 2.2.2 (Proposition 4.6, [6]). *An ideal $I \subseteq R$ is graded if and only if it is generated by homogeneous elements.*

Proof. First, if I is a graded ideal $I = \bigoplus_p I^p$ and is generated by $\bigcup_p I^p$. Then, each

$$I^p = I \cap R^p \subseteq R^p$$

is a subset of homogeneous elements. Therefore, I is generated by homogeneous elements.

Now, let I be generated by a set X of homogeneous elements. For sure, $I \cap R^p \subseteq I$, so we just need to prove the converse inclusion. As I is generated by X , its elements $u \in I$ are of the form

$$u = \sum_i r_i x_i s_i, \quad (2.1)$$

for $r_i, s_i \in R$ and $x_i \in X$. And as $I \subseteq R$, also,

$$u = \sum_p u_p,$$

for $u_p \in R^p$. For every term in (2.1), we have

$$r_i = \sum_j r_{i,j}, \quad s_i = \sum_l s_{i,l},$$

with each $r_{i,j}, s_{i,l}$ being homogeneous. Therefore, combining all we have that

$$u = \sum_i \sum_{j,l} r_{i,j} x_i s_{i,l}. \quad (2.2)$$

Each term in (2.2) is homogeneous as is a product of homogeneous elements. Thus u_p is the sum of those terms, and u has degree p . Therefore $u_p \in I$ and $I \subseteq \bigoplus_p I \cap R^p$. \square

Theorem 2.2.3 (Structure). [6, Proposition 4.8] *Let (V, π) be a persistence module. There exist a barcode $\text{Bar}(V, \pi)$, with $\mu: \text{Bar}(V, \pi) \rightarrow \mathbb{N}$, the multiplicity of the barcode intervals, such as there is a unique direct sum decomposition*

$$V \cong \bigoplus_{I \in \text{Bar}(V)} \mathbb{F}(I)^{\mu(I)}. \quad (2.3)$$

Proof. V is of finite type, so it is a finite $\mathbb{F}[x]$ -module. As \mathbb{F} is a field, $\mathbb{F}[x]$ is a principal ideal domain, therefore, V is a finitely generated module over a principal ideal domain. Using Fact 2.2.1, V can be decompose in the direct sum of its free and torsion subgroups, $F \oplus T$. Thus, we have

$$F = \bigoplus_{i \geq q} x^i \cdot \mathbb{F}$$

$$T = \bigoplus_{i \geq q} R^i / I^i.$$

Each $x^i \cdot \mathbb{F}$ is isomorphic to ideals of the form (x^q) . By Proposition 2.2.2, each R^i / I^i is isomorphic to some quotient of graded ideals of the form $(x^p)/(x^r)$. Note that the free subgroup can be seen as a particular case of the torsion group taking $r = 0$. Thus V can be decompose as described in (2.3). \square

3

Interleaving Stability Theorem

In this section we are going to give a detailed proof of the first stability theorem for persistence homology. This theorem is referred to as the “geometry miracle” of persistent homology, as it allows to describe an isometry between persistence modules and barcodes [4]. This shows that *small* changes in a data sets will perform *small* changes in their persistence modules, and therefore small changes in how persistent homology groups vary through time. The theorem claims that given two persistence modules, the distance between them using the interleaving distance, is the same as the distance between their barcodes using the bottleneck distance.

Theorem 3.0.1 (Stability). *[5, Theorem 2.2.8] There is an isometry between a persistence module with the interleaving distance and its barcode with the bottleneck distance. That is, given two persistence modules V and W , it holds that*

$$d_{int}(V, W) = d_{bot}(\text{Bar}(V), \text{Bar}(W)).$$

For the presented proof we have followed [5]. Hence, we will divide the proof into proving two complementary inequalities separately. This implies checking that if there exists a δ -matching between two given barcodes, then there exists a δ -interleaving morphism between, Proposition 3.0.3. Also, we need to check that, if there exists a δ -interleaving morphism between two persistence modules, then there exists a δ -matching between their barcodes, 3.0.11.

The first claim can be deduced from the Structure Theorem in a rather direct way,

proving first the case where our modules are just interval modules.

Lemma 3.0.2. [5, Exercise 2.2.7] *Let I, J be two δ -matched intervals. Then, their corresponding interval modules $(\mathbb{F}(I), \pi)$ and $(\mathbb{F}(J), \theta)$ are δ -interleaved.*

Proof. Let $I = (a, b]$, $J = (c, d]$. If ρ is the δ -matching between them, then $\rho(I) = J$ and, following Definition 1.0.12, $(a, b] \subseteq (c - \delta, d + \delta]$ and $(c, d] \subseteq (a - \delta, b + \delta]$, with $b - a > 2\delta$ and $d - c > 2\delta$. Then, the morphisms

$$\begin{aligned} \phi_\delta: \mathbb{F}(I) &\rightarrow \mathbb{F}(J)_\delta & \text{and} & & \psi_\delta: \mathbb{F}(J) &\rightarrow \mathbb{F}(I)_\delta \\ \phi_\delta(\mathbb{F}(I)_t) &\mapsto \mathbb{F}(J)_{(t+\delta)} & & & \psi_\delta(\mathbb{F}(J)_t) &\mapsto \mathbb{F}(I)_{(t+\delta)} \end{aligned}$$

are well defined as for any $t \in (a, b]$, $t + \delta \in (c, d]$, as $a + \delta > c$ and $b + \delta \leq d$. In the same way, for any $t \in (c, d]$, $t + \delta \in (a, b]$. Thus,

$$\psi_\delta \circ \phi_\delta(\mathbb{F}(I)_t) = \psi_\delta(\mathbb{F}(J)_{(t+\delta)}) = \mathbb{F}(I)_{(t+2\delta)} = \pi_{t \leq t+2\delta}(\mathbb{F}(I)_t)$$

and

$$\phi_\delta \circ \psi_\delta(\mathbb{F}(J)_t) = \phi_\delta(\mathbb{F}(I)_{(t+\delta)}) = \mathbb{F}(J)_{(t+2\delta)} = \theta_{t \leq t+2\delta}(\mathbb{F}(J)_t).$$

Therefore, ϕ_δ and ψ_δ are a pair of δ -interleaving morphisms. \square

Once we are able to build a δ -interleaving between two δ -matched interval modules, we will use the Structure Theorem for persistence modules to generalize the construction for arbitrary persistence modules. This will prove useful to prove the first inequality needed to prove Theorem 3.0.1.

Proposition 3.0.3. [5, Theorem 3.0.1] *Given two persistence modules V, W , if there is a δ -matching between their barcodes, then there is a δ -interleaving morphism between them.*

Proof. Suppose that $\rho: \text{Bar}(V) \rightarrow \text{Bar}(W)$ is a δ -matching between the barcodes of V and W . By the Structure Theorem 2.2.3, V and W decompose in a finite direct sum of interval modules

$$V \cong \bigoplus_{I \in \text{Bar}(V)} \mathbb{F}(I), \quad W \cong \bigoplus_{J \in \text{Bar}(W)} \mathbb{F}(J).$$

We can express $V = V_Y \oplus V_N$, $W = W_Y \oplus W_N$ denoting

$$\begin{aligned} V_Y &\cong \bigoplus_{I \in \text{coim } \rho} \mathbb{F}(I), & V_N &\cong \bigoplus_{I \in \text{Bar}(V) \setminus \text{coim } \rho} \mathbb{F}(I), \\ W_Y &\cong \bigoplus_{J \in \text{im } \rho} \mathbb{F}(J), & W_N &\cong \bigoplus_{J \in \text{Bar}(W) \setminus \text{im } \rho} \mathbb{F}(J). \end{aligned}$$

The V_Y, W_Y modules separate the “yes, matched” intervals, from the V_N, W_N “not matched” intervals. For every interval $I \in \text{Bar}(V_Y)$, I is δ -matched to an interval $J \in \text{Bar}(W_Y)$ by $\rho(I) = J$. Thus, by Lemma 3.0.2, for all pair I, J of matched intervals, there exist a pair of δ -interleaved morphisms

$$\phi_\delta: \mathbb{F}(I) \rightarrow \mathbb{F}(J)_\delta \quad \text{and} \quad \psi_\delta: \mathbb{F}(J) \rightarrow \mathbb{F}(I)_\delta$$

which induce the pair of δ -interleaved morphisms

$$\phi_\delta: V_Y \rightarrow W_{Y\delta} \quad \text{and} \quad \psi_\delta: W_Y \rightarrow V_{Y\delta}.$$

Not matched intervals are of length smaller than 2δ , therefore both, V_N and V_Y are δ -interleaved with the empty set. We can now construct the δ -interleaving morphism $\phi: V \rightarrow W$ such as $\phi|_{V_Y} = \phi_Y$ and $\phi|_{V_N} = 0$. In a similar way, we also construct $\psi: W \rightarrow V$. \square

With Proposition 3.0.3 we have proven the first halve of Stability Theorem 3.0.1. Now we need to prove that we can build a δ -interleaving morphism from a δ -matching. To verify this claim we need several previous lemmas that will lead us to prove Proposition 3.0.3.

First, we will introduce some notation. Let $(V, \pi), (W, \theta)$ be two persistence modules and let $I = (b, d]$ be an interval with $d \in \mathbb{R} \cup \{+\infty\}$. Denote the set of bars of $\text{Bar}(V)$ that start before b end exactly at d as

$$\text{Bar}_{I-}(V) := \{(a, d] \in \text{Bar}(V) : a \leq b\}.$$

Analogously, denote the set of bars that start at b and end after d as

$$\text{Bar}_{I+}(V) := \{(b, c] \in \text{Bar}(V) : c \geq d\}.$$

Lemma 3.0.4. [5, Proposition 3.1.1] *Let $I = (b, d]$ be an interval. It exists an injective morphism $\iota: (V, \pi) \rightarrow (W, \theta)$, then $\#(\text{Bar}_{I-}(V)) \leq \#(\text{Bar}_{I-}(W))$. Where $\#(\cdot)$ denotes the cardinal operator.*

Proof. For $b < s < d < r$, denote the set of elements in V_d which come from all V_s and disappear in all V_r as

$$E_{I-} = \bigcap_{b < s < d} \text{im } \pi_{s \leq d} \cap \bigcap_{r > d} \ker \pi_{d \leq r} \subseteq V_d.$$

It holds that $\dim E_{I-}(V) = \#(\text{Bar}_{I-}(V))$. For every morphism $p: (V, \pi) \rightarrow (W, \theta)$ the following diagram commutes

$$\begin{array}{ccc} V_s & \xrightarrow{\pi_{s \leq r}} & V_r \\ p_s \downarrow & & \downarrow p_r \\ W_s & \xrightarrow{\theta_{s \leq r}} & W_r \end{array}$$

This implies that

$$p_r(\text{im } \pi_{s \leq r}) \subseteq \text{im } \theta_{s \leq r}, \quad p_s(\ker \pi_{s \leq r}) \subseteq \ker \theta_{s \leq r}.$$

Taking $r = d$, $b < s < d$ in the first inclusion, and $s = d$, $r > d$ in the second, we have that

$$p_d(\text{im } \pi_{s \leq d}) \subseteq \text{im } \theta_{s \leq d}, \quad p_d(\ker \pi_{d \leq r}) \subseteq \ker \theta_{d \leq r},$$

and

$$p_d(E_{I-}(V)) \subseteq E_{I-}(W).$$

If we now take p as the injective morphism of the hypothesis, $p = \iota$, we get

$$\dim E_{I-}(V) \leq \dim E_{I-}(W).$$

□

Lemma 3.0.5. [5, Exercise 3.1.3] *Let $I = (b, d]$ be an interval. It exists a surjective morphism $s: (V, \pi) \rightarrow (W, \theta)$, then $\#(\text{Bar}_{I+}(V)) \geq \#(\text{Bar}_{I+}(W))$.*

Proof. Analogously to the proof of Lemma 3.0.4 we now define

$$E_{I+}(V) = \bigcap \text{im } \pi_{d \leq r}.$$

Therefore $\dim E_{I+}(V) = \#(\text{Bar}_{I+}(V))$, and recalling the diagram used for the previous proof, and using the fact that is commutative, we have that

$$p_r(\text{im } \pi_{s \leq r}) \supseteq \text{im } \theta_{s \leq r}.$$

Taking $s = d$ we then have that

$$p_d(E_{I+}(V)) \supseteq E_{I+}(W).$$

And finally, taking the surjective morphism $p = s$ we have that

$$\dim E_{I-}(V) \geq \dim E_{I-}(W).$$

□

To build our δ -matching we first define two induced matchings, by an injection and by a surjection respectively. First, suppose that there exists an injection $\iota: V \rightarrow W$. For every $c \in \mathbb{R} \cup \{\infty\}$, sort the bars $(a_i, c] \in \text{Bar}(V)$, $i \in \{1, \dots, k\}$ by decreasing length order,

$$(a_1, c] \supseteq (a_2, c] \supseteq \dots \supseteq (a_k, c], \text{ with } a_1 \leq a_2 \leq \dots \leq a_k.$$

Sort in the same manner the bars $(b_j, c] \in \text{Bar}(V)$, $j \in \{1, \dots, l\}$,

$$(b_1, c] \supseteq (b_2, c] \supseteq \dots \supseteq (b_k, c], \text{ with } b_1 \leq b_2 \leq \dots \leq b_k.$$

As there is an injection between V and W , Lemma 3.0.4 assures that the amount of bars in $\text{Bar}(V)$ is lower than the amount in $\text{Bar}(W)$, i.e., $k \leq l$. We define the *injective induced matching* $\mu_{inj}: \text{Bar}(V) \rightarrow \text{Bar}(W)$ matching, for each $c \in \mathbb{R} \cup \{\infty\}$, the intervals from both lists by decreasing length.

Lemma 3.0.6. [5, Proposition 3.1.5] *If there exists an injection $\iota: (V, \pi) \rightarrow (W, \theta)$, then the induced matching $\mu_{inj}: \text{Bar}(V) \rightarrow \text{Bar}(W)$ satisfies:*

1. $\text{coim } \mu_{inj} = \text{Bar}(V)$,
2. $\mu_{inj}(a, c] = (b, c], \forall b \leq a, \forall (a, d] \in \text{Bar}(V)$.

Proof. Applying Lemma 3.0.4 with the interval $(a_k, c]$, we have that for each $c \in \mathbb{R} \cup \{\infty\}$, $\# \text{Bar}_{(a_k, c]}(V) \leq \# \text{Bar}_{(a_k, c]}(W)$, having $k \leq l$ as we note earlier. This means that every bar in $\text{Bar}(V)$ is matched to some bar in $\text{Bar}(W)$. Hence $\text{coim } \mu_{inj} = \text{Bar}(V)$. Moreover, as the matching is carried out in length descending order, for each $i \in \{1, \dots, k\}$, $\mu_{inj}(a_i, c] = (b_i, c]$, and applying Lemma 3.0.4, now with the interval $(a_i, c]$, and making the same reasoning, $a_i \leq b_i$. \square

Now we suppose that there exists a surjection $\sigma: V \rightarrow W$. For every $a \in \mathbb{R}$, sort the intervals $(a, c_i] \in \text{Bar}(V)$, $i \in \{1, \dots, k\}$ by decreasing length order as before,

$$(a, c_1] \supseteq (a, c_2] \supseteq \dots \supseteq (a, c_k], \text{ with } c_1 \geq c_2 \geq \dots \geq c_k,$$

and again in the same manner, sort the intervals $(a, d_j] \in \text{Bar}(V)$, $j \in \{1, \dots, l\}$,

$$(a, d_1] \supseteq (a, d_2] \supseteq \dots \supseteq (a, d_l], \text{ with } d_1 \geq d_2 \geq \dots \geq d_l.$$

We define the *surjective induced matching* $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(W)$ matching, for each $a \in \mathbb{R}$, the intervals from both lists by decreasing length.

Lemma 3.0.7. [5, Exercise 3.1.8] *If there exists a surjection $s: (V, \pi) \rightarrow (W, \theta)$, then the induced matching $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(W)$ satisfies:*

1. $\text{im } \mu_{sur} = \text{Bar}(W)$,
2. $\mu_{sur}(a, c] = (a, d], \forall c \geq d, \forall (a, d] \in \text{Bar}(V)$.

Proof. Using Lemma 3.0.5 with the interval $(b, d_k]$ for each $b \in \mathbb{R}$, we get that, as there exists a surjection between the modules, now $k \geq l$. Therefore, every bar in $\text{Bar}(W)$ is matched to some bar in $\text{Bar}(V)$ and $\text{im } \mu_{sur} = \text{Bar}(W)$. In an analogue way to the previous lemma, as the intervals in both lists are matched in a decreasing manner, for every $j \in \{1, \dots, l\}$, $\mu_{sur}(a, c_j] = (a, d_j]$, and if we now apply Lemma 3.0.5, we get that $c_j \geq d_j$. \square

Hence, with the injective and the surjective induced matchings at hand, for a general morphism f , we can define the *induced matching* $\mu(f): \text{Bar}(V) \rightarrow \text{Bar}(W)$, as the composition $\mu_{inj} \circ \mu_{sur}$, defined as $\text{im } \mu_{sur} = \text{Bar}(\text{im } f) = \text{coim } \mu_{inj}$.

The following lemma verifies that, in fact, the mapping between persistence modules with its morphisms and barcodes with induced matchings (either the injective or the surjective versions) has functorial properties between the two categories.

Lemma 3.0.8. [5, Claim 3.1.13] *Let U, V and W persistence diagrams and f, g, h morphisms between them defined as in the following diagram:*

$$\begin{array}{ccccc} U & \xrightarrow{f} & V & \xrightarrow{g} & W \\ & & \searrow & \nearrow & \\ & & h & & \end{array} .$$

If all f, g, h are all injections, or all surjections, then the corresponding diagram formed by the barcodes of the modules, and their respective matchings commutes as well.

$$\begin{array}{ccccc} \text{Bar}(U) & \xrightarrow{\mu_*(f)} & \text{Bar}(V) & \xrightarrow{\mu_*(g)} & \text{Bar}(W) \\ & & \searrow & \nearrow & \\ & & \mu_*(h) & & \end{array} .$$

Where μ_* denotes μ_{inj} or μ_{sur} accordingly.

Proof. Let f, g, h injective morphisms, by the definition of the injective induced matching and Lemma 3.0.4 for any $d \in \mathbb{R} \cup \{+\infty\}$, there exist $k \leq l \leq q$ such that the barcodes of U, V, W consist on the following bars:

$$\begin{aligned} \text{Bar}(U) &: (a_1, d] \supset \dots \supset (a_k, d] \\ \text{Bar}(V) &: (b_1, d] \supset \dots \supset (b_k, d] \supset \dots \supset (b_l, d] \\ \text{Bar}(W) &: (c_1, d] \supset \dots \supset (c_k, d] \supset \dots \supset (c_l, d] \supset \dots \supset (c_q, d]. \end{aligned}$$

Therefore, for any d the diagram commutes as

$$\mu_{inj}(f)(a_i, d] = (b_i, d], \quad \mu_{inj}(g)(b_i, d] = (c_i, d], \quad \mu_{inj}(h)(a_i, d] = (c_i, d]$$

for $1 \leq i \leq k$. If f, g, h were surjective morphisms, an analogue reasoning using the surjective induced matching definition and Lemma 3.0.5 completes the proof. \square

Finally, we can claim the two main lemmas from which we will construct our desired δ -matching.

Lemma 3.0.9. [5, Lemma 3.2.1] *Let $(V, \pi), (W, \theta)$ be δ -interleaved persistence modules, with δ -interleaving morphisms $\phi: V \rightarrow W_\delta$ and $\psi: W \rightarrow V_\delta$. Let $\phi: V \rightarrow \text{im } \phi$ be a surjection and $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(\text{im } \phi)$ the induced matching. Then*

1. $\text{coim } \mu_{sur} \supseteq \text{Bar}(V)_{\geq 2\delta}$,
2. $\text{im } \mu_{sur} = \text{Bar}(\text{im } \phi)$ and
3. $\mu_{sur}(b, d] = (b, d'], \forall (b, d] \in \text{coim } \mu_{sur}, d' \in [d - 2\delta, d]$.

Proof. 1. To check the first part, we observe that, in the following diagram, the three morphisms are surjective as ϕ and $\pi_{t \leq t+2\delta}$ are defined onto their images, and the diagram commutes,

$$\begin{array}{ccccc} V & \xrightarrow{\phi} & \text{im } \phi & \xrightarrow{\psi_\delta} & \text{im } \pi_{t \leq t+2\delta} \\ & \searrow & & \nearrow & \\ & & \pi_{t \leq t+2\delta} & & \end{array}.$$

Therefore, because of Lemma 3.0.8 the barcode diagram also commutes:

$$\begin{array}{ccccc} \text{Bar}(V) & \xrightarrow{\mu_{sur}(\phi)} & \text{Bar}(\text{im } \phi) & \xrightarrow{\mu_{sur}(\psi_\delta)} & \text{Bar}(\text{im } \pi_{t \leq t+2\delta}) \\ & \searrow & & \nearrow & \\ & & \mu_{sur}(\pi_{t \leq t+2\delta}) & & \end{array}.$$

By the definition of the surjective induced matching,

$$\text{coim } \mu_{sur}(\pi_{t \leq t+2\delta}) = \text{Bar}(V)_{\geq 2\delta}.$$

For each starting point $a \in \mathbb{R}$, we have that

$$\text{Bar}(\text{im } \pi_{t \leq t+2\delta}) = \{(a, b - 2\delta]: (a, b] \in \text{Bar}(V), b - a > 2\delta\}.$$

Sorting all bars of $\text{Bar}(V)$ and of $\text{Bar}(\text{im } \pi_{t \leq t+2\delta})$ in length-not-increasing order and matching the bars though the longest-first order, each bar $(a, b] \in \text{Bar}(V)$ is matched with the bar $(a, b - 2\delta] \in \text{Bar}(\text{im } \pi_{t \leq t+2\delta})$ while $b - a > 2\delta$. The smaller bars are not matched. Thus,

$$\text{coim } \mu_{sur}(\phi) \supseteq \text{coim } \mu_{sur}(\text{im } \pi_{t \leq t+2\delta}) = \text{Bar}(V)_{\geq 2\delta}.$$

2. The second part is just a reformulation of Lemma 3.0.4.

3. Let $(b, d] \in \text{coim}$. There are two cases:

On one hand, if $d - b \leq 2\delta$, $(b, d]$ is matched to $(b, d']$ where $d \geq d'$, by definition of μ_{sur} . Also, $d' > b$ and, as in this case we have $b \geq d - 2\delta$, we have $d' > d - 2\delta$. Therefore, $d' \in [d - 2\delta, d]$.

On the other hand, if $d - b > 2\delta$, $(b, d]$ is matched to $(b, d']$ by $\mu_{sur}(\phi)$, with $(b, d'] \in W_{\leq 2\delta}$. We can therefore use Lemma 3.0.7 to check that $d' \geq d$. In the same manner, $(b, d']$ is matched to $(b, d'']$ by $\mu_{sur}(\psi)_\delta$ with $d'' \geq d'$. Finally, using the commutativity of the following diagram, we have that $(b, d'') = (b, d - 2\delta]$, making $d' \in [d - 2\delta, d]$.

$$\begin{array}{ccccc}
 \text{Bar}(V)_{\geq 2\delta} & & \text{Bar}(\text{im } \phi) & & \text{Bar}(\text{im } \pi_{t \leq t+2\delta}) \\
 \Psi & & \Psi & & \Psi \\
 (b, d] & \xrightarrow{\mu_{sur}(\phi)} & (b, d'] & \xrightarrow{\mu_{sur}(\psi)_\delta} & (b, d''] \\
 & & & & \parallel \\
 & & & & (b, d - 2\delta] \\
 & \searrow \mu_{sur}(\pi_{t \leq t+2\delta}) & & &
 \end{array}$$

□

Lemma 3.0.10. [5, Proposition 3.2.2] Let $(V, \pi), (W, \theta)$ be δ -interleaved persistence modules, with δ -interleaving morphisms $\phi: V \rightarrow W_\delta$ and $\psi: W \rightarrow V_\delta$. Let $\phi: V \rightarrow \text{im } \phi$ be a injection and $\mu_{inj}: \text{Bar}(\text{im } \phi) \rightarrow \text{Bar}(W_\delta)$ the induced matching. Then

1. $\text{coim } \mu_{sur} = \text{Bar}(\text{im } \phi)$,
2. $\text{im } \mu_{inj} \supseteq \text{Bar}(W_\delta)_{\geq 2\delta}$ and
3. $\mu_{inj}(b, d'] = (b', d']$, $(b, d'] \in \text{coim } \mu_{inj}$, $b' \in [b - 2\delta, b]$.

Proof. 1. Immediate using Lemma 3.0.6.

2. As $\phi_\delta \circ \psi = \theta_{t \leq t+2\delta}$ the following diagram commutes:

$$\begin{array}{ccccc}
 W & \xrightarrow{\psi} & \text{im } \psi & \xrightarrow{\phi_\delta} & W_{2\delta} \\
 & \searrow \theta_{t \leq t+2\delta} & & \nearrow &
 \end{array}$$

This implies that $\text{im } \theta_{t \leq t+2\delta} \subseteq \text{im } \phi_\delta \subseteq W_{2\delta}$, so there are some injections j and i which make the following diagram commute as well:

$$\begin{array}{ccccc}
 \text{im } \theta_{t \leq t+2\delta} & \xrightarrow{j} & \text{im } \phi_\delta & \xrightarrow{i} & W_{2\delta} \\
 & \searrow \theta_{t \leq t+2\delta} & & \nearrow &
 \end{array}$$

As all morphisms in the diagram above are injections, we can use the functorial properties of Lemma 3.0.8 having a commutative diagram of the barcodes of each of the previous persistence modules:

$$\begin{array}{ccccc} \text{Bar}(\text{im } \theta_{t \leq t+2\delta}) & \xrightarrow{\mu_{inj}(j)} & \text{Bar}(\text{im } \phi_\delta) & \xrightarrow{\mu_{inj}(i)} & \text{Bar}(W_{2\delta}) \\ & & & \nearrow & \\ & & & \mu_{inj}(\theta_{t \leq t+2\delta}) & \end{array}.$$

We have that

$$\begin{aligned} \text{Bar}(\text{im } \theta_{t \leq t+2\delta}) &= \{(b, d - 2\delta) : (b, d] \in \text{Bar}(W), b < d - 2\delta\}, \\ \text{Bar}(W_{2\delta}) &= \{(b - 2\delta, d - 2\delta) : (b, d] \in \text{Bar}(W)\} \text{ and} \\ \mu_{inj}(\theta_{t \leq t+2\delta})((b, d - 2\delta)) &= (b - 2\delta, d - 2\delta) \end{aligned}$$

Therefore $\text{im}_\mu \text{inj}(i) \supseteq \text{im } \mu_{inj}(\psi_{t \leq t+2\delta}) = \text{Bar}(W_{2\delta})_{2\delta}$. This, undoing the shifts made, make the prove.

3. Let $(b, d] \in \text{Bar}(\text{im } f_\delta)$ such as for some b' , $\mu_{inj}(b, d] = (b', d] \in \text{Bar}(W)$. Because of Lemma 3.0.6, $b' \leq b$. There are again two cases:

If $d - b \leq 2\delta$, then $b' \geq d - 2\delta \geq b > b - 2\delta$ and $b' \in [b - 2\delta, b]$.

Else, if $d - b > 2\delta$, there exists an interval $(b' + 2\delta, d] \in \text{Bar}(\text{im } \theta_{t \leq t+2\delta})$ such that

$$\mu_{inj}(\theta_{t \leq t+2\delta})(b' + 2\delta, d] = \mu_{inj}(i)(b, d] = (a, d].$$

Thus, $b \leq b' + 2\delta$ and $b' \in [b - 2\delta, b]$.

□

At last, we can now prove the other part of the Stability theorem. For so, we will construct a δ -matching out of a δ -interleaving morphism.

Proposition 3.0.11. [5, Theorem 3.0.2] *Given two persistence modules V , W , with a δ -interleaving morphism between them, then there is a δ -matching between their barcodes.*

Proof. Let $\mu(\phi) = \mu_{inj} \circ \mu_{sur}$ and let $\Phi_\delta: \text{Bar}(W_\delta) \rightarrow \text{Bar}(W)$ be the *shift map* that carries each bar $(a, b]$ into $(a + \delta, b + \delta]$. The composition $\Phi_\delta \circ \mu(\phi)$ is a matching between

$\text{Bar}(V)$ and $\text{Bar}(W)$. Hence, using Lemma 3.0.10 and 3.0.9, we get the following diagram:

$$\begin{array}{ccccc}
 \text{Bar}(V) & & \text{Bar}(W_\delta)_{\geq 2\delta} & & \text{Bar}(W)_{\geq 2\delta} \\
 \cup \downarrow & & \cap \downarrow & & \cap \downarrow \\
 \text{Bar}(V)_{\geq 2\delta} & \xrightarrow{\mu_{sur}} & \text{Bar}(\text{im } f) & \xrightarrow{\mu_{inj}} & \text{im } \mu_{inj} & \xrightarrow{\Psi_\delta} & \text{Bar } B(W) \\
 \Psi \downarrow & & \Psi \downarrow & & \Psi \downarrow & & \Psi \downarrow \\
 (b, d] & \longmapsto & (b, d'] & \longmapsto & (b', d'] & \longmapsto & (b' + \delta, d' + \delta]
 \end{array}$$

The diagram shows that, by Lemma 3.0.9, a bar $(b, d] \in \text{Bar}(V)_{\geq 2\delta}$ is sent to $\mu_{sur}(b, d] = (b, d'] \in \text{Bar}(\text{im } \phi)$ with $d' \in [d - 2\delta, d]$. Then, by Lemma 3.0.9, it is sent to $\mu_{sur}(b, d'] = (b', d']$ with $b' \in [b - 2\delta, b]$. At last, using the shift morphism Φ_δ it is carried to $(b' + \delta, d' + \delta]$.

This shows that any bar in $\text{Bar}(V)_{\geq 2\delta}$ is matched. In the same manner it can be seen that any bar in $\text{Bar}(W)_{\geq 2\delta}$ is matched. Thus, we have that

$$\begin{cases} d - 2\delta \leq d' \leq d \\ b - 2\delta \leq b' \leq b \end{cases} \Rightarrow \begin{cases} d - \delta \leq d' + \delta \leq d + \delta \\ b - \delta \leq b' + \delta \leq b + \delta \end{cases},$$

and therefore, $\Phi_\delta \circ \mu(\phi)$ is a δ -matching between $\text{Bar}(V)$ and $\text{Bar}(W)$. \square

The constructions made by Proposition 3.0.3 and Proposition 3.0.11 assure that given a δ -interleaving morphism we can build a δ -matching, and conversely, given a δ -matching we can build a δ -interleaving morphism. This means that if one of the two exists, it fixes a δ . Both the interleaving distance and the bottleneck distance try to minimize this δ , so once fixed for one of them, the other needs an smaller or equal δ' . Thus, with each of the propositions we can prove one of the inequalities needed to reach the isomorphism between the space of persistence diagrams and the space of their barcodes.

Theorem 3.0.1 (Stability). *[5, Theorem 2.2.8] There is an isometry between a persistence module with the interleaving distance and its barcode with the bottleneck distance. That is, given two persistence modules V and W , it holds that*

$$d_{int}(V, W) = d_{bot}(\text{Bar}(V), \text{Bar}(W)).$$

Proof. Suppose $d_{int}(V, W) = \delta$. Proposition 3.0.11 assures there exist a δ -matching between $\text{Bar}(V)$ and $\text{Bar}(W)$. As $d_{bot}(V, W)$ is the infimum δ for which exists a δ -matching, $d_{bot}(V, W) \leq d_{int}(V, W)$. On the other hand, Proposition 3.0.3 leads, with the same reasoning, to $d_{int}(V, W) \leq d_{bot}(V, W)$. Thus, it has to be $d_{int}(V, W) = d_{bot}(\text{Bar}(V), \text{Bar}(W))$. \square

4

Edelsbrunner & Harer's (Hausdorff) Stability Theorem

Persistence diagrams help summarize the information given by the homology groups of a filtration over a certain data set. They represent the birth and death of every feature in an easy to analice format scattering points over the upper half of the plane \mathbb{R}^2 and its diagonal Δ . Once we have computed the diagrams given by two datasets we can measure distances between them using the bottleneck distance, enabling us to decide wether two diagrams are close to each other. However, this would be kind of useless if the bottleneck distance were not stable. If minor differences in original data would cause great changes in the bottleneck distance between the corresponding persistence diagrams, then this data summary method would be as good as any other random method.

Fortunately for us, David Cohen-Steiner, Herbert Edelsbrunner and John Harer, proved in their 2005 paper that bottleneck distance over persistence diagrams is indeed stable [2]. This means that, when comparing the bottleneck distance between the persistence diagrams formed by the pre images of two tame functions, the first will be always lower or equal that the Lebesgue L_∞ norm between the two functions.

Along this chapter we will follow Edelsbrunner et al. paper [2] to prove Theorem 4.0.4. To simplify notation, along this chapter we will adopt the following. Let $H_k(X)$ be the k -th singular homology group of a topological space X . The dimension of $H_k(x)$ is denoted by the k -th Betti number $\beta_k(X) := \dim H_k(x)$.

Let $f: X \rightarrow \mathbb{R}$, $x < y \in \mathbb{R}$. Denote the k -th homology group of the pre-image by f of an interval $(-\infty, x]$ as $F_x := H_k(f^{-1}(-\infty, x])$. Denote the inclusion map from the k -th homology group F_x to the k -th homology group F_y as $f_x^y: F_x \rightarrow F_y$. Finally, denote $F_x^y := \text{im } f_x^y$.

Note that if $y = \infty$, F_x^y is the trivial group. Also, if $x = \infty$, then $y = \infty$ too.

Definition 4.0.1 (Homological critical value). Let X be a topological space and let $f: X \rightarrow \mathbb{R}$. A **homological critical value** of f a number $a \in \mathbb{R}$ such that there exists $k \in \mathbb{Z}$ such that for all $\varepsilon > 0$, the morphism $H_k(f^{-1}(-\infty, a - \varepsilon)) \rightarrow H_k(f^{-1}(-\infty, a + \varepsilon))$ is not an isomorphism.

Definition 4.0.2 (Tame function). A function $f: X \rightarrow \mathbb{R}$ is said to be **tame** if it has a finite number of homological critical values, and for all $z \in \mathbb{Z}$, and for all $a \in \mathbb{R}$, $\dim F_a < \infty$.

Definition 4.0.3 (Multiplicity). Let $f: X \rightarrow \mathbb{R}$ be tame, and $(a_i)_{i=1, \dots, n}$ be its homological critical values. Take $(b_i)_{i=1, \dots, n}$ be an interleaved sequence of non critical values such that $b_{i-1} < a_i < b_i$ for all $i = 1, \dots, n$. Define $b_{-1} = a_0 = -\infty$, $b_{n+1} = a_{n+1} = \infty$. The **multiplicity** of $(a_i, a_j) \in D(f)$, denoted μ_i^j is

$$\mu_i^j := \beta_{b_{i-1}}^{b_j} - \beta_{b_i}^{b_j} + \beta_{b_i}^{b_{j-1}} - \beta_{b_{i-1}}^{b_{j-1}}.$$

The **total multiplicity** of a multiset A , denoted $\#(A)$ is the sum of the multiplicities of every element in A .

Note that the total multiplicity of a multiset is the the generalized concept of cardinality of a normal set. While the cardinality of a set counts the number of elements in the set, the multiplicity of a multiset counts how many elements, different or not, are there in the multiset.

Theorem 4.0.4 (Main Theorem, [2]). *Let X be a triangulable space, and $f, g: X \rightarrow \mathbb{R}$ continuous tame functions. Then,*

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_{\infty}$$

4.1 Hausdorff Stability

We will denote the closed upper left quadrant of a point $(x, y) \in \mathbb{R}^2$ as $Q_x^y := [-\infty, x] \times [y, \infty]$.

Lemma 4.1.1 (*k-Triangle Lemma*, [2]). *Let $f: X \rightarrow \mathbb{R}$ be a tame function, $x < y \in \mathbb{R}$ be non critical values of f . Then the multiplicity μ of the persistence diagram of f in the closed upper left quadrant is*

$$\mu = \#(D(f) \cap Q_x^y) = \beta_x^y.$$

Proof. Let $x = b_i$, $y = b_{j-1}$.

$$\mu = \sum_{k \leq i \leq j \leq l} \mu_k^l = \sum_{k \leq i \leq j \leq l} \beta_{b_{k-1}}^{b_l} - \beta_{b_k}^{b_l} + \beta_{b_k}^{b_{l-1}} - \beta_{b_{k-1}}^{b_{l-1}} \quad (4.1)$$

$$= \beta_{b_{-1}}^{b_{n+1}} - \beta_{b_i}^{b_{n+1}} + \beta_{b_i}^{b_{j-1}} - \beta_{b_{j-1}}^{b_{j-1}} = \beta_{b_k}^{b_{l-1}} = \beta_x^y. \quad (4.2)$$

The first two equalities in (4.1) are just the definition of total multiplicity. In (4.2), note that every other term in the sum cancels. Then note that $\beta_{b_{-1}}^{b_{n+1}} = \dim F_{-\infty}^\infty$, $\beta_{b_i}^{b_{n+1}} = \dim F_x^\infty$ and $\beta_{b_{j-1}}^{b_{j-1}} = \dim F_{-\infty}^y$. All of them are the dimension of the trivial group, therefore, equal to 0. This leaves only one remaining term and completes the proof. \square

Denote the **upper left quadrants** $Q := Q_b^c = [-\infty, b] \times [c, \infty]$, $Q_\varepsilon := Q_{b-\varepsilon}^{c+\varepsilon} = [-\infty, b-\varepsilon] \times [c+\varepsilon, \infty]$.

Lemma 4.1.2 (*Quadrant Lemma*, [2]). *Let $f, g: X \rightarrow \mathbb{R}$ be two tame functions. With the notation above, the following inequality holds,*

$$\#(D(f) \cap Q_\varepsilon) \leq \#(D(g) \cap Q).$$

Proof. Let $\varepsilon := \|f - g\|_\infty$. Hence, considering the pre-image of the functions, we have the following inclusions

$$f^{-1}((-\infty, x]) \subseteq g^{-1}((-\infty, x + \varepsilon)), \quad (4.3)$$

$$g^{-1}((-\infty, x]) \subseteq f^{-1}((-\infty, x + \varepsilon)). \quad (4.4)$$

Name $\varphi_x: F_x \rightarrow G_{x+\varepsilon}$ to the inclusion map induced by (4.3) and $\psi_x: G_x \rightarrow F_{x+\varepsilon}$ to the inclusion map induced by (4.4). Let $b < c \in \mathbb{R}$. With the described maps, we can form commutative diagram (4.5) where we observe that

$$\text{im}(f_{c-\varepsilon}^{c+\varepsilon}) = F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c \circ g_b^c(G_b) = \psi_c(G_b^c).$$

$$\begin{array}{ccc} F_{b-\varepsilon} & \xrightarrow{f_{b-\varepsilon}^{c+\varepsilon}} & F_{c+\varepsilon} \\ \varphi_{b-\varepsilon} \downarrow & & \uparrow \psi_c \\ G_b & \xrightarrow{g_b^c} & G_c \end{array} \quad (4.5)$$

Last inclusion is enough for the requirements of this proof, never the less, we will make one more note that will be useful latter on, through the proof of Lemma 4.1.3. Fit the maps so that they describe commutative diagram (4.6), showing that

$$\psi_c(G_b^c) = \psi_c \circ g_b^c(G_b) = f_{b+\varepsilon}^{c+\varepsilon} \circ \psi_b(G_b) \subseteq F_{b+\varepsilon}^{c+\varepsilon}.$$

$$\begin{array}{ccc} F_{b+\varepsilon} & \xrightarrow{f_{b+\varepsilon}^{c+\varepsilon}} & F_{c+\varepsilon} \\ \varphi_b \uparrow & & \uparrow \psi_c \\ G_b & \xrightarrow{g_b^c} & G_c \end{array} \quad (4.6)$$

From both diagrams we finally obtain the inclusion chain

$$F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c(G_b^c) \subseteq F_{b+\varepsilon}^{c+\varepsilon}. \quad (4.7)$$

By Lemma 4.1.1, we are able to count the elements in the intersection of the diagrams with the upper left quadrants. Hence

$$\begin{aligned} \#(D(f) \cap Q_\varepsilon) &= \beta_{b-\varepsilon}^{c+\varepsilon} = \dim F_{b-\varepsilon}^{c+\varepsilon}, \\ \#(D(g) \cap Q) &= \beta_b^c = \dim G_b^c. \end{aligned}$$

As if one homology group is contain in an other, the dimension of the first must be lower or equal to the one of the second. Also, the dimension is invariant under inclusion maps. Thus, the first inclusion of (4.7) asserts that $F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c(G_b^c)$ and therefore we have proven that $\dim F_{c-\varepsilon}^{c+\varepsilon} \leq \dim G_b^c$. \square

Before next lemma we will introduce some new notation. Let $f: X \rightarrow \mathbb{R}$ fe a tame function. Let $w < x < y < z \in \mathbb{R}$ be numbers different from critical values of f . Recall that $F_x = H_k(f^{-1}(-\infty, x])$, $f_x^y: F_x \rightarrow F_y$ and $F_x^y = \dim f_x^y$. We denote

$$f_x^{y,z} := f_y^z|_{F_x^y}, \quad F_x^{y,z} := \dim f_x^{y,z}.$$

Note, from linear algebra, that $\dim F_x^{y,z} = \dim F_x^y - \dim F_x^z$. Note too that $F_w^y \subseteq F_x^y$. Therefore, $\ker F_w^y \subseteq \ker F_x^y$ and we can define the quotient

$$F_{w,x}^{y,z} := F_x^{y,z} / F_w^{y,z}.$$

Let $a < b < c < d \in \mathbb{R}$. Denote the **rectangles** $R := [a, b] \times [c, d]$, $R_\varepsilon := [a + \varepsilon, b - \varepsilon] \times [c + \varepsilon, d - \varepsilon]$.


Lemma 4.1.3 (Box Lemma, [2]). *With the notation abobe, the following inequality holds,*

$$\#(D(f), \cap R_\varepsilon) \leq \#(D(g) \cap R).$$

Proof. Note that we can assume that $a + \varepsilon < b - \varepsilon$ and $c + \varepsilon < d - \varepsilon$. Otherwise there would not be rectangle R_ε . Also note that

$$\begin{aligned}\#(D(f) \cap R_\varepsilon) &= \dim F_{a+\varepsilon, b-\varepsilon}^{c+\varepsilon, d-\varepsilon} \\ \#(D(g) \cap R) &= \dim G_{a,b}^{c,d}\end{aligned}$$

To make our prove we draw diagram 4.1. Lets analice every element of the diagram.



First of all, the middle upside arrows, are

$$u_2 = f_{a+\varepsilon}^{c+\varepsilon, d-\varepsilon}, \quad u_3 = f_{b-\varepsilon}^{c+\varepsilon, d-\varepsilon}.$$

Right arrows r_1, r_2, r_3, r_4 represent the inclusions from its respective vector spaces to their destination, in the same manner. The objective is to define que respective quotients to define $G_{a,b}^{c,d}$. Recall the inclusion maps defined in the proof of Lemma 4.1.2, $\varphi_x: F_x \rightarrow G_{x+\varepsilon}$ and $\psi_x: G_x \rightarrow F_{x+\varepsilon}$. We define

$$E_b^c := \psi_c^{-1}(F_{b-\varepsilon}^{c+\varepsilon, d-\varepsilon}) \cap G_b^c, \quad E_a^c := G_a^c \cap E_b^c.$$

We have then that the outer upside arrows are the respective restrictions

$$u_1 = g_a^{c,d}|_{E_a^c} \quad u_4 = g_b^{c,d}|_{E_b^d}$$

□

Theorem 4.1.4 (Hausdorff Stability).

$$d_H(D(f), D(g)) \leq \|f - g\|_\infty.$$

4.2 Bottleneck Stability

In this section we are going to prove Theorem 4.0.4.

Definition 4.2.1 (Very close tame functions). Let $f, g: X \rightarrow \mathbb{R}$ be tame functions. We define

$$\delta_f = \min\{\|p - q\|_\infty : p \in D(f) \setminus \Delta, q \in D(f), p \neq q\}.$$

We say that g is **very close** to f if $\|f - g\|_\infty < \delta_f/2$.

Lemma 4.2.2 (Easy Bijection Lemma, [2]). Let $f, g: X \rightarrow \mathbb{R}$ be tame functions, where g is very close to f . Then, following holds,

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_\infty.$$

Proof. Let $p := (a_i, a_j) \in D(f) - \Delta$ be a point in the diagram of f that is not in the diagonal, and let $\mu := \beta_i^j$ denote its multiplicity. Let S_ϵ be the square of center p and radius $\epsilon = \|f - g\|_\infty$. That is, the square of side 2ϵ . By definition of the square S_ϵ we have that the number of points of its intersection with the diagram of g must be grater or equal than the multiplicity at p . Hence, by the Box Lemma 4.1.3 we have

$$\mu \leq \#(D(g) \cap S_\epsilon) \leq \#(D(f) \cap S_{2\epsilon}).$$

As g is very close to f , we have $2\epsilon \leq \delta_f$. Hence p is the only point of $D(f)$ in S_ϵ , and therefore the previous inequality is in fact an equivalence. If there was a point in the intersection which was not in μ then it would be inside the square S_ϵ and meaning the distance $\|f - g\|$ would be smaller. That is

$$\mu = \#(D(g) \cap S_\epsilon).$$

Hence we can map every point in $D(g) \cap S_{\epsilon \text{psilon}}$ with p . We can then repeat this process for every other $p \in D(f) \setminus \Delta$. After this process, every point of $D(g)$ which have not been matched jet, must be at distance grater than ϵ from $D(f) \setminus \Delta$. By Theorem 4.1.4, every unmatched point must be at distance at most ϵ from the diagonal Δ . Hence, if we map each of this points to Δ , we have built a bijection between $D(f)$ and $D(g)$ that moves each point at most ϵ . \square

Definition 4.2.3. Let \hat{f}, \hat{g} be two picewise linear functions over a simplicial complex K . Let $\lambda \in [0, 1]$. A **convex combination** of \hat{f} and \hat{g} is a function of the form

$$h_\lambda := (1 - \lambda)\hat{f} + \lambda\hat{g}.$$

Lemma 4.2.4 (Interpolation Lemma, [2]). Let K be a simplicial complex. Take two picewise linear functions $\hat{f}, \hat{g}: K \rightarrow \mathbb{R}$. Then, following holds,

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \|\hat{f} - \hat{g}\|_\infty.$$

Proof. Let $c := \|\hat{f} - \hat{g}\|_\infty$. For every $\lambda \in [0, 1]$, define $\delta(\lambda) := \delta_{h_\lambda} > 0$. Let J_λ denote open intervals around each δ as follows, and consider the set C of all J_λ be

$$C := \left\{ J_\lambda := \left(\lambda - \frac{\delta(\lambda)}{4c}, \lambda + \frac{\delta(\lambda)}{4c} \right) \right\}.$$

The set C is an open cover of the interval $[0, 1]$. Let C' be the minimal subcover of C . As $[0, 1]$ is compact, the subcover C' must be finite. Consider then $\lambda_1 < \lambda_2 < \dots < \lambda_n$ the midpoints of the intervals in C' . As C' is minimal, the each intersection $J_{\lambda_i} \cap J_{\lambda_{i+1}}$ is not empty. Hence

$$\lambda_i + \lambda_{i+1} \leq \frac{\delta(\lambda_i) + \delta(\lambda_{i+1})}{4c} \leq \frac{\max\{\delta(\lambda_i), \delta(\lambda_{i+1})\}}{2c}.$$

Therefore, by definition of c and each h_{δ_i} , it holds

$$\|h_{\delta_i} - h_{\delta_{i+1}}\|_\infty = c(\lambda_{i+1} - \lambda_i) \leq \frac{\max\{\delta(\lambda_i), \delta(\lambda_{i+1})\}}{2}.$$

This implies that h_{δ_i} is very close to $h_{\delta_{i+1}}$ or viceversa. Then, by Lemma 4.2.2, for every $1 \leq i \leq n-1$,

$$d_{\text{bot}}(D(h_{\lambda_i}), D(h_{\lambda_{i+1}})) \leq \|h_{\lambda_i} - h_{\lambda_{i+1}}\|_\infty. \quad (4.8)$$

Let $\lambda_0 = 0$ and $\lambda_{n+1} = 1$. Then $h_{\lambda_0} = \hat{f}$ is very close to h_{λ_1} and $h_{\lambda_1} = \hat{g}$ is very close to h_{λ_n} and therefore (4.8) also holds for $i = 0$ and $i = n+1$. Finally, using the triangle inequality we have

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \sum_{i=0}^n d_{\text{bot}}(D(h_{\lambda_i}), D(h_{\lambda_{i+1}})) \leq \sum_{i=0}^n \|h_{\lambda_i} - h_{\lambda_{i+1}}\|_\infty = \|\hat{f} - \hat{g}\|_\infty.$$

□

For the final prove lets recall

Definition 4.2.5 (Star of a simplicial complex). Let σ be a simplex in a simplicial complex L . The **star** $\text{St}(\sigma)$ of σ is the set of simplices in L which contain σ as a face. The **star of a subset** K of L , denoted $\text{St}(K)$, is the union of the stars of each simplex of K .

Theorem 4.0.4 (Main Theorem, [2]). Let X be a triangulable space, and $f, g: X \rightarrow \mathbb{R}$ continuous tame functions. Then,

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_\infty$$

Proof. As X is triangulable, there exists a finite simplicial complex L and a homeomorphism $\Phi: L \rightarrow X$. Hence a persistence diagram is invariant under this change of variables. That is, $f \circ \Phi: L \rightarrow \mathbb{R}$ is tame and $D(f \circ \Phi) = D(f)$. Since f and g are continuous and L is compact, for every $\delta > 0$ there exists a subdivision K of L such that for every u, v points of a common simplex $\sigma \in K$,

$$\begin{aligned} |f \circ \Phi(u) - f \circ \Phi(v)| &\leq \delta, \\ |g \circ \Phi(u) - g \circ \Phi(v)| &\leq \delta. \end{aligned}$$

Let $\hat{f}, \hat{g}: \text{St}(K) \rightarrow \mathbb{R}$ be the piecewise linear interpolations of $f \circ \Phi$ and $g \circ \Phi$ on K . By construction of K and the definition of the L_∞ -norm, these interpolations satisfy

$$\begin{aligned} \|\hat{f} - f \circ \Phi\|_\infty &\leq \delta, \\ \|\hat{g} - g \circ \Phi\|_\infty &\leq \delta. \end{aligned}$$

Hence, by Lemma 4.2.4 and the triangle inequality

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \|\hat{f} - \hat{g}\|_\infty \leq \|f \circ \Phi - g \circ \Phi\|_\infty + 2\delta \leq \|f - g\|_\infty + 2\delta.$$

Now, we can take some δ such that $\delta \leq \delta_f/2$ so \hat{f} is very close to f . This allows to use Lemma 4.2.2 to make a bijection that satisfy

$$d_{\text{bot}}(D(f), D(\hat{f})) \leq d_{\text{bot}}(D(f \circ \Phi), D(\hat{f})) \leq \delta.$$

Analogously, also assuring $\delta < \delta_g$ we also have

$$d_{\text{bot}}(D(g), D(\hat{g})) \leq d_{\text{bot}}(D(g \circ \Phi), D(\hat{g})) \leq \delta,$$

and therefore, by triangle inequality again,

$$d_{\text{bot}}(D(f), D(g)) \leq d_{\text{bot}}(D(f), D(\hat{f})) + d_{\text{bot}}(D(\hat{f}), D(\hat{g})) + d_{\text{bot}}(D(\hat{g}), D(g)) \leq 4\delta.$$

As this holds for any δ smaller than δ_f and δ_g , taking the limit when δ tends to 0, we complete the proof. \square

5

Gromov-Hausdorff's Stability Theorem

[1]

6

Vectorizations' Stability Theorems

6.1 Persistence landscapes

6.2 Persistence images

6.3 Euler curves

Bibliography

- [1] BURAGO, D., BURAGO, Y., AND IVANOV, S. *A Course in Metric Geometry*. American Mathematical Society, 2001.
- [2] EDELSBRUNNER, H., COHEN-STEINER, D., AND HARER, J. Stability of persistence diagrams. *University of Chicago* (2005).
- [3] HUNGERFORD, T. W. *Algebra*. Springer, 2000.
- [4] NANDA, V. Computational algebraic topology, lecture notes. *University of Oxford* (2020).
- [5] POLTEROVICH, L., ROSEN, D., SAMVELYAN, K., AND ZHANG, J. *Topological Persistence in Geometry and Analysis*. American Mathematical Society, 2020.
- [6] WANG, K. G. The basic theory of persistent homology. *University of Chicago* (2012).