

Universidad Autónoma de Madrid

FINAL MASTER THESIS

STRUCTURE AND STABILITY THEOREMS  
IN TOPOLOGICAL DATA ANALYSIS  
DRAFT

Author:

Gonzalo Ortega Carpintero

Tutor:

Jaime Santos Rodríguez

13/05/2025

## **Abstract**

This is a draft version of a in-work Master's thesis in TDA.

## **Key words**

- Topological Data Analysis
- Persistent homology
- Topological summary
- Stability

## **Acknowledgments**

To Manuel M. Cuerno because of all his help and advise, though the paperwork circumstances did not let him be my thesis tutor.

# Contents

<b>Introduction</b>	<b>1</b>
<b>1 Preliminaries</b>	<b>3</b>
1.1 Persistent homology . . . . .	4
1.2 Persistence modules and interleaving distance . . . . .	7
1.3 Barcodes and the bottleneck distance . . . . .	10
1.4 Persistence diagrams and the Wasserstein distance . . . . .	11
1.5 The Hausdorff and Gromov-Hausdorff distances . . . . .	14
<b>2 Structure Theorem</b>	<b>19</b>
2.1 Algebraic structure theorem . . . . .	19
2.2 Structure theorem for persistence diagrams . . . . .	20
<b>3 Interleaving Stability Theorem</b>	<b>23</b>
3.1 First inequality . . . . .	23
3.2 Second inequality . . . . .	25
<b>4 Hausdorff Stability Theorem</b>	<b>33</b>
4.1 Hausdorff Stability . . . . .	34
4.2 Bottleneck Stability . . . . .	38
<b>5 Gromov-Hausdorff's Stability Theorem</b>	<b>42</b>
5.1 Gromov-Hausdorff stability . . . . .	43
5.2 A generalization with endowed functions . . . . .	46
<b>6 Vectorizations' Stability Theorems</b>	<b>49</b>
6.1 Persistence landscapes . . . . .	49
6.2 Persistence images . . . . .	52
6.3 Euler curves . . . . .	54
<b>Bibliography</b>	<b>55</b>

# Introduction

Topological Data Analysis (TDA) is an emerging field at the intersection of mathematics, statistics, and computer science that leverages tools from algebraic topology to extract robust, interpretable features from complex, high-dimensional data. Traditional data analysis techniques often struggle to capture the intrinsic shape and connectivity of modern datasets—ranging from biological networks and sensor arrays to financial time series and high-dimensional point clouds. TDA addresses this gap by focusing on the topological invariants of data, which provide a mathematically rigorous description of its global structure. Central to this approach is the idea that the “shape” of data—characterized by holes, voids, and connected components—reveals critical insights impervious to noise and geometric distortions.

The cornerstone of TDA is persistent homology, a multiscale extension of classical homology. While homology groups capture topological features (e.g., connected components, loops, cavities) of a space at a fixed scale, persistent homology tracks the birth, persistence, and death of these features across a continuum of scales. This process is formalized through filtrations: nested sequences of topological spaces (e.g., simplicial complexes) built from data, parameterized by a scale parameter (e.g., distance thresholds). As the scale evolves, homology groups induce a structure known as a persistence module—an algebraic object encoding how topological features evolve. The Structure Theorem (Chapter 2) classifies these modules, decomposing them into elementary interval modules. This decomposition underpins two powerful visual and analytical summaries: barcodes and persistence diagrams.

A fundamental challenge in TDA is ensuring that topological summaries are stable: small perturbations in input data should induce only controlled changes in the output. Stability theorems form the bedrock of reliable applications, guaranteeing that TDA-based inferences are robust to noise, discretization artifacts, and measurement errors. This thesis provides a unified treatment of four pillars of TDA stability:

- **Interleaving Stability** (Chapter 3): Establishes an isometry between the algebraic interleaving distance for persistence modules and the combinatorial bottleneck distance for barcodes.
- **Hausdorff Stability** (Chapter 4): Bounds the bottleneck distance between persistence diagrams of functions by their  $L^\infty$ -distance, crucial for functional data.

- Gromov-Hausdorff Stability (Chapter 5): Extends stability to metric spaces, linking the bottleneck distance of Vietoris–Rips or Čech complexes to the Gromov–Hausdorff distance between spaces.
- Vectorization Stability (Chapter ??): Proves robustness for practical summaries like persistence landscapes, images, and Euler curves, enabling integration with machine learning workflows.

This work synthesizes foundational results from algebraic topology, category theory, and metric geometry to present a cohesive theoretical framework for TDA. By elucidating the stability properties of topological summaries, we aim to fortify the mathematical foundations of data analysis and empower applications across scientific domains—from understanding protein folding to quantifying shape in high-dimensional datasets. The subsequent chapters develop these ideas rigorously, beginning with the preliminaries of persistent homology and culminating in modern vectorization techniques.

# Chapter 1

## Preliminaries

Topological Data Analysis is a wide topic, where results from all Mathematical branches are put together in order to compute topological invariants among real world data. Basic knowledge in both, Algebra and Analysis, as well as Geometry is needed to approach the theoretical results that strengthen the main theory that defines proves the needed structures and their stability. Therefore, in order to settle a common base and notation unifying the diverse procedures among bibliography, this chapter introduces the main definitions and basic results needed to tackle the rest of the thesis.

Section 1.1 makes a brief recall of simplicial homology in order to define persistent homology modules. Section 1.2 then introduces persistence modules, the algebraic tool used to study the structure of persistence homology groups. Section 1.3 introduces a tool to summarize the data from the persistence homology groups, the barcodes. With this summary it is possible to compute how homologically different are two datasets, or how different are two filtrations among the same data. To make this measure, the interleaving and bottleneck distance are introduced. The first one measures distances between persistence modules, while the later one measures distance between the barcodes of those modules. One key aspect of them, is that they are actually equivalent, as its later proved in Chapter 3.

A visual way of presenting barcodes is through persistence diagrams. Both barcodes and persistence diagrams are equivalent manners of presenting the homological features of data. To give a different perspective about how to measure distances between persistence diagrams, Section 1.4 introduces the Wasserstein distance and redefines the bottleneck distance as a particular case.

Finally, Section 1.5 recalls the Hausdorff distance to measure distance between subsets of metric spaces, and generalizes it presenting the Gromov-Hausdorff distance, which allows to measure distances between sets located in different metric spaces. This will be useful to delimit bottleneck distance in order to proof its stability.

Section 1.1 primarily follows [16]. Sections 1.2 and 1.3 are based on the presentation at [14]. Section 1.4 takes the definitions given at [3] and Section 1.5 follows [4].

## 1.1 Persistent homology

In most Algebraic Topology introductory texts, homology is first presented in the form of homology groups (see [12], [10]). We are going to generalize homology groups to homology  $R$ -modules based on the notes by Wang [16]. This makes possible to take coefficients over an arbitrary commutative ring  $R$  and will lead us to introduce persistent homology and the persistence modules. Lets start recalling what an  $R$ -module is, to be followed by a brief presentation of simplicial homology.

**Definition 1.1.1** ( $R$ -module, Definition IV.1.1.1 [11]). Let  $R$  be a commutative ring. An  $R$ -**module** is an abelian group  $(M, +)$  with an operation  $\cdot : R \times M \rightarrow M$  such that for all  $r, s \in R$  and for all  $x, y \in M$ ,

- (i)  $(rs) \cdot x = r(s \cdot x)$ ,
- (ii)  $(r + s) \cdot x = r \cdot x + s \cdot x$ ,
- (iii)  $r \cdot (x + y) = r \cdot x + r \cdot y$ .

If  $R$  has a multiplicative identity 1, then  $M$  is said to be a **unitary  $R$ -module** and

- (iv)  $1 \cdot x = x$ .

If  $R$  is a division ring, that is, a ring with identity where every non zero element is a unit, then a unitary  $R$ -module is called a **left  $R$ -vector space**. Note that in this case,  $R$  is in fact a field. Also note that, when we take  $R = (\mathbb{Z}, +, \cdot)$ , a  $\mathbb{Z}$  module is actually an abelian group.

**Definition 1.1.2** (Affinely independent tuple). A  $(k + 1)$ -tuple of points  $(x_0, \dots, x_k)$  in  $\mathbb{R}^n$ , where each  $x_i \in \mathbb{R}^n$  is said to be **affinely independent** if the set of vectors  $\{x_j - x_0 \mid 1 \leq j \leq k\}$  is linearly independent.

**Definition 1.1.3** ( $k$ -simplex). A  $k$ -**simplex** is an ordered  $(k + 1)$ -tuple of affinely independent points  $\sigma = (x_0, \dots, x_k)$ . The **vertices** of  $\sigma$  are the points of the set  $V = \{x_0, \dots, x_k\}$ . Every simplex  $\sigma$  induces a total ordering  $\preceq_\sigma$  on its set of vertices  $V$ , where if  $p \leq q$ , then  $x_p \preceq_\sigma x_q$  in  $V$ . For  $m \leq n$ , an  $(m + 1)$ -tuple  $(y_0, \dots, y_m)$ , where each  $y_j \in V$  and if  $j \leq k$ , then  $y_j \preceq_\sigma y_k$ , is called an  $m$ -**face** of  $\sigma$ .

**Definition 1.1.4** (Abstract simplicial complex). An **abstract simplicial complex**  $K$  is a finite collection of simplices such that for every face of a simplex in  $K$  is also a simplex in  $K$ .

**Definition 1.1.5** (Siplicial  $k$ -chain). Let  $R$  be a commutative ring with additive identity 0 and multiplicative identity 1 and let  $K$  be a simplicial complex. A **simplicial  $k$ -chain** is a formal sum of  $k$ -simplices,

$$\sum_{i=1}^N r_i \sigma_i,$$

such that  $r_i \in R$  and  $\sigma_i \in K$ .

Note that that set of simplicial  $k$ -chains with formal addition over  $R$  form a  $R$ -module. Denote that  $R$ -module as  $K_k$ .

**Definition 1.1.6** (Boundary map). Let  $K$  be a simplicial complex, and let  $K_k$  and  $K_{k-1}$  be two  $R$ -modules from  $K$ . For every  $k$ -simplex  $\sigma = (x_0, \dots, x_k) \in K$ , the **boundary** map  $\partial_k: K_k \rightarrow K_{k-1}$  is defined as

$$\partial_k(\sigma) = \sum_{i=0}^k (-1)^i (x_0, \dots, \hat{x}_i, \dots, x_k),$$

where  $(x_0, \dots, \hat{x}_i, \dots, x_k)$  denotes the  $(k-1)$ -face of  $\sigma$  obtained by removing the vertex  $x_i$ .

As the boundary map can be extends linearly to arbitrary  $k$ -chains, it forms an  $R$ -module homomorphism  $\partial_k: K_k \rightarrow K_{k-1}$ .

**Definition 1.1.7** (Chain complex). A **chain complex**  $(K, d)$  is a sequence of  $R$ -modules  $A_k$  with boundary maps  $d_k: A_k \rightarrow A_{k-1}$ , such that the composition  $d_{k-1} \circ d_k = 0$  for all  $k \leq 0$ .

**Lemma 1.1.8** (Lemma 2.7, [16]). *The collection of  $R$ -modules  $K_k$  connected by boundary maps  $\partial_k$  forms a chain complex  $(K, \partial)$ .*

*Proof.* ... □

**Corolary 1.1.9.** *The boundary maps  $\partial_k$  satisfy that for every  $k \leq 0$ ,  $\text{im } \partial_{k+1} \subseteq \ker \partial_k$ .*

*Proof.* ... □

**Definition 1.1.10** (Homology module). Let  $K$  be a simplicial complex. The  **$k$ -th homology module** of  $K$  is the  $R$ -module given by

$$H_k(K) := \frac{\ker \partial_k}{\text{im } \partial_{k+1}}.$$



Homology can be extended to every topological space, not just those that are triangulable by introducing singular homology. However, in this work, we will restrict ourselves to simplicial homology. This choice is motivated by the nature of the data typically encountered in topological data analysis. The spaces involved are usually "sufficiently nice" and therefore triangulable. Moreover, real-world data intended for computational analysis is finite by nature, and hence, triangulable. Furthermore, simplicial homology of a finite simplicial complex  $K$  is computable (see [12][Chapter 4, Theorem 11.5]) what makes it a perfect tool for computational studies.

**Definition 1.1.11** (Filtration). Let  $K$  be a simplicial complex and let  $T$  be a complete ordered index set. A **filtration** of  $K$  is a nested sequence of subcomplexes  $F_i K$ ,  $i \in T$ , such that for every  $i \in T$ ,

$$F_i \subseteq F_{i+1}.$$

We will denote the **inclusion simplicial maps** by  $g_i: F_i \hookrightarrow F_{i+1} K$ .

Filtrations are a natural way of obtaining simplicial complexes from datasets, as it is possible to set the first element of the filtration to be the points in the datasets, and then join in various steps each of the point to form a simplicial complex that connects every point by simplices of different dimensions. Naturally, depending on the procedure we follow we could get quite different filtration for one same dataset.

Considering a filtration of a simplicial complex, we can compute the homology groups at each step of the filtration. Hence, for each  $k \geq 0$ , there are induced linear maps  $H_k g_i: H_k(F_i K) \rightarrow H_k(F_{i+1} K)$  that fit together into a sequence of vector spaces

$$H_k(F_0 K) \xrightarrow{H_k g_0} H_k(F_1 K) \xrightarrow{H_k g_1} \dots \xrightarrow{H_k g_{n-2}} H_k(F_{n-1} K) \xrightarrow{H_k g_{n-1}} H_k(F_n K).$$

As homology is functorial, when we consider the inclusion map formed by the composition of several others, the induced map at homology level also becomes a composition of the corresponding maps. That is, for any pair  $i \leq j$  of indices, where  $i, j = 1, \dots, n$ , the map induced on homology by the inclusion map  $g_{i \leq j}: F_i K \hookrightarrow F_j K$ , is given by the composition

$$H_k g_{i \leq j} = H_k g_{j-1} \circ H_k g_{j-2} \circ \dots \circ H_k g_{i+1} \circ H_k g_i.$$

Such maps  $H_k g_{i \leq j}$  give really relevant information about our data, as they permit us to relate the  $k$ -th homology groups of every subcomplex  $F_i K$  that appear in a filtration of  $K$ .

Now if  $X$  is a triangulable topological space, and  $f: X \rightarrow T$ , we can extend the concept of filtration to be attained to  $X$ . For clarity, fix  $T = \mathbb{R}$ . Hence, for each  $a \in \mathbb{R}$ , the pre image of the interval  $(-\infty, a]$  by  $f$  is some subspace  $X_a$  of  $X$ . That is,

$$f^{-1}((-\infty, a]) = X_a.$$

As  $X$  is triangulable, we will look for functions that satisfy that each  $X_a$  is triangulable too. And therefore we can compute its homology module.

**Definition 1.1.12** (Homological critical value). Let  $X$  be a topological space and let  $f: X \rightarrow \mathbb{R}$ . A **homological critical value** of  $f$  is a number  $a \in \mathbb{R}$  such that there exists  $k \in \mathbb{Z}$  such that for all  $\varepsilon > 0$ , the morphism  $H_k(f^{-1}(-\infty, a - \varepsilon]) \rightarrow H_k(f^{-1}(-\infty, a + \varepsilon])$  is not an isomorphism.

**Definition 1.1.13** (Tame function). A function  $f: X \rightarrow \mathbb{R}$  is said to be **tame** if it has a finite number of homological critical values, and for all  $z \in \mathbb{Z}$ , and for all  $a \in \mathbb{R}$ ,  $\dim F_a < \infty$ .

**Definition 1.1.14** (Persistent homology module). Let  $X$  be a triangulable topological space and let  $f: X \rightarrow \mathbb{R}$ . For every fixed  $k \in \mathbb{Z}$ , we denote

$$F_x := H_k(f^{-1}(-\infty, x]),$$

and for every  $x \leq y$ , we denote the induced inclusion map from  $F_x$  to  $F_y$  as

$$f_x^y: F_x \rightarrow F_y.$$

The **persistent homology module** associated to  $x$  and  $y$  is the subspace of  $F_y$  given by

$$F_x^y := \text{im } f_x^y.$$

The **persistent Betti numbers** are defined as

$$\beta_x^y := \dim(F_x^y).$$

## 1.2 Persistence modules and interleaving distance

To enable a more rigorous and structured study, we introduce the category of persistence modules along with their morphisms. This categorical framework provides the foundation for analyzing the stability and structure of persistent homology.

**Definition 1.2.1** (Persistence module). Let  $\mathbb{F}$  be a field and let  $T$  be a totally ordered set. Let  $V = \{V_t\}_{t \in T}$  be a collection of  $\mathbb{F}$ -vector spaces. A  $T$ -indexed **persistence module** is a pair  $(V, \pi)$  such that  $\pi = \{\pi_{s \leq t}\}$  is a collection of linear maps  $\pi_{s \leq t}: V_s \rightarrow V_t$  that verifies that for all  $r, s, t \in T$ ,

$$\pi_{r \leq s} \circ \pi_{s \leq t} = \pi_{r \leq t}.$$

**Definition 1.2.2** (Morphism between persistence modules). Let  $T$  be a totally ordered set. Let  $(V, \pi), (W, \theta)$  be two persistence modules. A **morphism** between persistence

modules  $p: (V, \pi) \rightarrow (W, \theta)$  is a family of linear maps  $p_t: V_t \rightarrow W_t$  such that for all  $s \leq t$  the following diagram commutes:

$$\begin{array}{ccc} V_s & \xrightarrow{\pi_{s \leq t}} & V_t \\ p_s \downarrow & & \downarrow p_t \\ W_s & \xrightarrow{\theta_{s \leq t}} & W_t \end{array}$$

If a morphism  $i$  verifies that for all  $t \in T$ ,  $i_t: V_t \rightarrow V_t$  is the identity, then  $i$  is the **identity morphism**. If there exists two morphisms  $p: (V, \pi) \rightarrow (W, \theta)$  and  $q: (W, \theta) \rightarrow (V, \pi)$  such that the compositions  $p \circ q$  and  $q \circ p$  are both the identity morphism, then  $p$  and  $q$  are **isomorphisms** of persistence modules. In this case,  $(V, \pi)$  and  $(W, \theta)$  are said to be **isomorphic** persistence modules.

For now on, to simplify notation, we will limit our totally order set to be the real numbers,  $T = \mathbb{R}$ . Also, when there is no possible confusion, we might denote the persistence module  $(V, \pi)$  by just its collection of vector spaces  $V$ .

This section defines a distance between persistence modules that will allow us to differentiate how far are two modules from each other.

**Definition 1.2.3** (Persistence module shift). Let  $(V, \pi)$  be a persistence module and let  $\delta \in \mathbb{R}$ . The  $\delta$ -**shift** of  $(V, \pi)$  is the persistence module  $(V_\delta, \pi_\delta)$  defined by taking

$$(V_\delta)_t := V_{t+\delta}, \quad (\pi_\delta)_{s \leq t} := \pi_{s+\delta \leq t+\delta}.$$

**Proposition 1.2.4** (Exercise 1.2.3, [14]). Let  $\delta > 0$ . Let  $(V, \pi), (V_\delta, \pi_\delta)$  be a persistence module and its shift. The map  $\phi_\delta: (V, \pi) \rightarrow (V_\delta, \pi_\delta)$ , defined as

$$\phi_\delta(V_t) := \pi_{t \leq t+\delta}(V_t) = V_{t+\delta},$$

is a persistence module morphism.

*Proof.* As  $\delta > 0$ , then  $t \leq t + \delta$ . Hence

$$\begin{aligned} \phi_\delta \circ \pi_{t \leq t+\delta}(V_t) &= \phi_\delta(V_{t+\delta}) = V_{t+\delta+\delta} = V_{t+2\delta}, \\ \pi_{t+\delta \leq t+2\delta} \circ \phi_\delta(V_t) &= \pi_{t+\delta \leq t+2\delta}(V_{t+\delta}) = V_{t+2\delta}. \end{aligned}$$

□

**Definition 1.2.5** (Shift morphism). The persistence module morphism  $\phi_\delta$  defined as in Proposition 1.2.4 is named  $\delta$ -**shift morphism**.

**Definition 1.2.6** ( $\delta$ -interleaved modules). Let  $(V, \pi), (W, \theta)$  be two persistence modules and let  $\delta > 0$ .  $V$  and  $W$  are  **$\delta$ -interleaved** if there exists two persistence module morphisms  $\phi: V \rightarrow W_\delta$  and  $\psi: W \rightarrow V_\delta$  such that the following diagrams commute:

$$\begin{array}{ccc} V & \xrightarrow{\phi} & W_\delta \xrightarrow{\psi_\delta} V_{2\delta} \\ & \searrow \pi_{2\delta} & \nearrow \\ & & \end{array}, \quad \begin{array}{ccc} W & \xrightarrow{\psi} & V_\delta \xrightarrow{\phi_\delta} W_{2\delta} \\ & \searrow \theta_{2\delta} & \nearrow \\ & & \end{array}.$$

Persistence modules are a vast abstract algebraic tool. In order to make it more manageable, we give it some more structure, restricting the dimension of the vector spaces. Also, we limit as the amount of different up to isomorphism vector spaces there are. Note that this is analogous to the tame functions we used in order to define persistent homology.

**Definition 1.2.7** (Tame persistence module). A persistence module  $(V, \pi)$  over  $\mathbb{R}$  is **tame** if

- (i) For all  $t \geq 0$ ,  $\dim(V_t)$  is finite.
- (ii) For any  $\varepsilon > 0$ , there exists a finite subset  $K \subset \mathbb{R}$  such that for all  $t \in \mathbb{R} \setminus K$ , the map  $\pi_{t-\varepsilon \leq t+\varepsilon}: V_{t-\varepsilon} \rightarrow V_{t+\varepsilon}$  is not an isomorphism.

**Definition 1.2.8** (Interleaving distance). Let  $(V, \pi)$  and  $(W, \theta)$  be two tame persistence modules. The **interleaving distance** between them is defined as

$$d_{\text{int}}(V, W) := \inf\{\delta > 0 \mid V \text{ and } W \text{ are } \delta\text{-interleaved}\}.$$

Tameness property ensures that the persistence modules have manageable and well-behaved structures, which are essential for the infimum in the definition of the interleaving distance to exist and be meaningful. If there was some infinite dimensional vector space  $V_t$  or  $W_t$ , they could lead to some pathological behavior, making it difficult to define or compute distances meaningfully. On the other hand, if there were infinite many different isomorphisms, then the infimum in the definition of the interleaving distance might not exist or could be zero for modules that are not "close" in any intuitive sense.

**Proposition 1.2.9.** *The interleaving distance between two tame persistence modules is actually a distance.*

*Proof.* ... □

**Definition 1.2.10** (Interval module). Let  $I = (a, b]$  be an interval with  $b \leq \infty$  and let  $\mathbb{F}$  be a field. An **interval module**  $\mathbb{F}(I)$  is a persistence module defined as

$$\mathbb{F}(I)_t := \begin{cases} \mathbb{F} & \text{if } t \in I, \\ 0 & \text{else,} \end{cases} \quad \pi_{s \leq t} = \begin{cases} \text{Id} & \text{if } t \in I, \\ 0 & \text{else.} \end{cases}$$

**Definition 1.2.11** (Direct sum of persistence modules). Let  $(V, \pi)$  and  $(V', \pi')$  be two persistence modules. Their **direct sum**  $(W, \theta)$  is a persistence module where

$$W_t := V_t \oplus V'_t, \text{ the direct sum of both vector spaces, and}$$

$$\theta_{s \leq t} := \pi_{s \leq t} \oplus \pi'_{s \leq t}.$$

## 1.3 Barcodes and the bottleneck distance

Barcodes are the main representation element that will allow us to resume the information given by the persistence homology groups of a dataset.

**Definition 1.3.1** (Barcode). A **barcode**  $B$  is a finite multiset of intervals. That is, a collection  $\{(I_i, m_i)\}$  of intervals  $I_i$  with multiplicities  $m_i \in N$ , where each interval  $I_i$  is either finite of the form  $(a, b]$  or infinite of the form  $(a, \infty)$ . Each interval  $I_i$  is named to be a **bar** of  $B$ . The first number,  $a$  is named the **birth** of the barcode and its second number is its **death**.

Chapter 2 proves the Structure Theorem of persistence modules, that asserts that it is possible to express any persistence module as a direct sum of the interval modules given by the bars of a barcode. This is a key result in our purpose of introduce measures between the homological features of datasets.

Given an interval  $I = (a, b]$ , and some  $\delta \geq 0$ , we will denote

$$I^\delta := (a - \delta, b + \delta].$$

**Definition 1.3.2** (Multiset matching). Let  $X$  and  $Y$  be two multi-sets and let  $X' \subseteq X$ ,  $Y' \subseteq Y$ . A **matching** between them is a bijection  $\mu: X' \rightarrow Y'$ . The elements in  $X'$  and  $Y'$  are said to be **matched** by  $\mu$ .

Note that  $X' = \text{coim}(\mu)$  and  $Y' = \text{im}(\mu)$ . Also note that as  $X$  and  $Y$  are multisets, it might happen that one same element appears several times in one of the multisets, and that some, but not all of its copies are matched to some element in the other multiset.

**Definition 1.3.3** ( $\delta$ -matching barcodes). A delta matching between two barcodes  $B$  and  $C$  is a multiset matching that verifies

1.  $B_{2\delta} \subseteq \text{coim}(\mu)$ ,
2.  $C_{2\delta} \subseteq \text{im}(\mu)$ ,
3. If  $\mu(I) = J$ , then  $I \subseteq J^\delta$  and  $J \subseteq I^\delta$ .

There are various ways of defining the bottleneck distance, all of them equivalent to one another. We first give the natural definition that comes up following the use of  $\delta$ -matchings.

**Definition 1.3.4** (Bottleneck distance). The **bottleneck distance** between two barcodes  $B$  and  $C$  is the infimum over all  $\delta \in \mathbb{R}$  such that there exists a  $\delta$ -matching between  $B$  and  $C$ .

## 1.4 Persistence diagrams and the Wasserstein distance

Barcodes can be seen as multisets of pair of points  $(a, b)$ , where  $b > a$ . Hence we could treat each interval  $(a, b]$  in a barcode as a point  $(a, b)$  in the Euclidean plane. We will denote the strict upper triangular region of the Euclidean plane as

$$\mathbb{R}_{<}^2 := \{(x, y) \in \mathbb{R}^2 : x < y\},$$

and the diagonal of the plane as

$$\Delta := \{(x, y) \in \mathbb{R}^2 : x = y\}.$$

**Definition 1.4.1** (Persistence diagram). Let  $I$  be a countable multiset. A *persistence diagram* is a function  $D : I \rightarrow \mathbb{R}_{<}^2$ .

Persistence diagrams are just a way of presenting barcodes as subsets of the plane, so we could introduce more distances between them. Note that while formally a persistence diagram is a function, usually it is used to also denote the set of points and their multiplicities in the plane. Note that when we do so, it is equivalent to name a barcode than a persistence diagram. Hence, the bottleneck distance as defined in Definition 1.3.4 can be also used to measure distances between persistence diagrams.

While definition 1.3.4 is purely algebraic, this section gives an alternative analytical definition for the bottleneck distance making an adaptation of the classic Wasserstein distance, used originally to measure distances between probability measures.

Instead of probability measures, now we are actually dealing with countable sets of points in  $\mathbb{R}$ . We will make use of the  $l^p$  norm at countable spaces to measure the distance between matched pairs and the distance between unmatched pairs and the diagonal  $\Delta$ . For a more detailed explanation of Lebesgue measures check [15][Definition 3.7]. This norm is named after Pafnuty Chebyshev.

**Definition 1.4.2** (Chebyshev distance). Let  $a, b \in \mathbb{R}^2$  with  $a = (a_x, a_y)$  and  $b = (b_x, b_y)$ . The *Chebyshev distance* is defined as

$$d_\infty(a, b) := \|a - b\|_\infty := \max\{|a_x - b_x|, |a_y - b_y|\}.$$

To define our adapted Wasserstein distance we need to check how Chebyshev distance measures distances between points of  $\mathbb{R}_{<}^2$  and  $\Delta$ .

**Proposition 1.4.3.** *If  $a = (a_x, a_y) \in \mathbb{R}_{<}^2$ , then  $d_\infty(a, \Delta) = \inf_{t \in \Delta} d_\infty(a, t) = \frac{a_y - a_x}{2}$ .*

*Proof.* The  $t$  which minimizes the distance is the midpoint of  $a_x$  and  $a_y$ , that is  $t = (\frac{a_x + a_y}{2}, \frac{a_x + a_y}{2})$ . Then,

$$\left| a_x - \frac{a_x + a_y}{2} \right| = \left| \frac{a_x - a_y}{2} \right| = \left| \frac{a_y - a_x}{2} \right| = \left| a_y - \frac{a_x + a_y}{2} \right|,$$

and as  $a_y > a_x$  we have

$$d_\infty(a, t) = \left| \frac{a_y - a_x}{2} \right| = \frac{a_y - a_x}{2}.$$

□

We now verify that the upper triangular region of the Euclidean plane with the Chebyshev distance adapted to measure distances in  $\Delta$  is a metric space.

**Proposition 1.4.4.** *The function  $d_\infty$  is a distance in  $\mathbb{R}_{<}^2$  with the diagonal  $\Delta$ .*

*Proof.* For points  $a, b \in \mathbb{R}_{<}^2 \subset \mathbb{R}^2$ ,  $d_\infty$  is a distance as usual Lebesgue norms are well defined. See [15][Chapter 3]. To verify that the metric requirements are fulfilled for  $d_\infty(a, \Delta)$ , it is enough to consider  $t = \frac{a_y - a_x}{2}$  as in Proposition 1.4.3. □

**Definition 1.4.5** (Partial matching). Let  $D_1 : I_1 \rightarrow \mathbb{R}_{<}^2$  and  $D_2 : I_2 \rightarrow \mathbb{R}_{<}^2$  be persistence diagrams. A *partial matching* between  $D_1$  and  $D_2$  is the triple  $(I'_1, I'_2, f)$  such that  $f : I'_1 \rightarrow I'_2$  is a bijection with  $I'_1 \subseteq I_1$  and  $I'_2 \subseteq I_2$ .

**Definition 1.4.6** ( $p$ -cost). Let  $D_1 : I_1 \rightarrow \mathbb{R}_{<}^2$  and  $D_2 : I_2 \rightarrow \mathbb{R}_{<}^2$  be persistence diagrams. Let  $(I'_1, I'_2, f)$  be a partial matching between them. If  $p < \infty$ , the  $p$ -cost of  $f$  is defined as

$$\begin{aligned} \text{cost}_p(f) := & \left( \sum_{i \in I'_1} d_\infty(D_1(i), D_2(f(i)))^p \right. \\ & + \sum_{i \in I_1 \setminus I'_1} d_\infty(D_1(i), \Delta)^p \\ & \left. + \sum_{i \in I_2 \setminus I'_2} d_\infty(D_2(i), \Delta)^p \right)^{\frac{1}{p}}. \end{aligned}$$

For  $p = \infty$ , the  $\infty$ -cost of  $f$  is defined as

$$\text{cost}_\infty(f) := \max \left\{ \sup_{i \in I'_1} d_\infty(D_1(i), D_2(f_i)), \right. \\ \left. \sup_{i \in I_1 \setminus I'_1} d_\infty(D_1(i), \Delta), \right. \\ \left. \sup_{i \in I_2 \setminus I'_2} d_\infty(D_2(i), \Delta) \right\}.$$

**Definition 1.4.7** ( $p$ -Wasserstein distance). Let  $D_1, D_2$  be persistence diagrams. Let  $1 \leq p \leq \infty$ . Define

$$\tilde{\omega}_p(D_1, D_2) = \inf \{ \text{cost}_p(f) : f \text{ is a partial matching between } D_1 \text{ and } D_2 \}.$$

Let  $\emptyset$  denote the unique persistence diagram with empty indexing set. Let  $(\text{Dgm}_p, \omega_p)$  be the space of persistence diagrams  $D$  that satisfy  $\tilde{\omega}_p(D, \emptyset) < \infty$  modulo the equivalence relation  $D_1 \sim D_2$  if  $\tilde{\omega}_p(D_1, D_2) = 0$ . The metric  $\omega_p$  is called the  $p$ -Wasserstein distance.

**Definition 1.4.8** (Bottleneck distance). In the conditions of Definition 1.4.7, if  $p = \infty$ , the metric  $\omega_\infty$  is called the *bottleneck distance*.

**Proposition 1.4.9.** *There is only one matching between  $D : I \rightarrow \mathbb{R}_{<}^2$  and  $\emptyset$ . Hence, if  $p \leq \infty$ ,*

$$\tilde{\omega}_p(D, \emptyset) = \left( \sum_{i \in I} d_\infty(D(i), \Delta)^p \right)^{\frac{1}{p}},$$

and, if  $p = \infty$ ,

$$\tilde{\omega}_\infty(D, \emptyset) = \sup_{i \in I} d_\infty(D(i), \Delta)$$

*Proof.* Let  $I' \subseteq D$ . If  $f$  is a partial matching between  $D$  and  $\emptyset$ , means that  $f(I') = \emptyset$  is a bijection. That is only possible if  $I' = \emptyset$  too. Therefore  $I \setminus I' = I \setminus \emptyset = I$  and following Definition 1.4.6 we conclude our proof.  $\square$

Next proposition will prove that, in indeed, the space of persistence diagrams with the  $p$ -Wasserstein distance  $(\text{Dgm}_p, \omega_p)$  is a metric space. Its proof is usually omitted in literature, as it based on the simple fact that  $d_\infty$  is a distance. We will give, however, an step by step version here.

**Proposition 1.4.10.**  $\omega_p$  is a distance on the space  $(\text{Dgm}_p, \omega_p)$ .

*Proof.* Let  $D_1, D_2, D_3 \in \text{Dgm}_p$ , with  $1 \leq p \leq \infty$ . First of all,  $\omega_p(D_1, D_2) \geq 0$  because  $d_\infty \geq 0$ .  $\omega_p(D_1, D_2) = 0$  if and only if  $\tilde{\omega}_p(D_1, D_2) = 0$ . Thus, because of the equivalence relationship used to define  $\omega_p$ , it has to be  $D_1 \sim D_2$ .



To check symmetry, note that every partial matching  $f$  is bijective, therefore  $f^{-1}$  is a partial matching. But, for all  $i \in I'_1$ , exists  $j \in I'_2$  such that  $f(i) = j$  and

$$d_\infty(D_1(i), D_2(f(i))) = d_\infty(D_2(f(i)), D_1(i)) = d_\infty(D_2(j), D_1(f^{-1}(j))).$$

Then,  $\text{cost}_p(f) = \text{cost}_p(f^{-1})$  and we have

$$\begin{aligned} \omega_p(D_1, D_2) &= \inf\{\text{cost}_p(f) : f \text{ is a partial matching between } D_1 \text{ and } D_2\} \\ &= \inf\{\text{cost}_p(f^{-1}) : f^{-1} \text{ is a partial matching between } D_2 \text{ and } D_1\} \\ &= \omega_p(D_2, D_1). \end{aligned}$$

Finally, lets prove the triangle inequality. If  $f : I'_1 \rightarrow I'_2$  is a partial matching between  $D_1$  and  $D_2$  and  $g : I'_2 \rightarrow I'_3$  is a partial matching between  $D_2$  and  $D_3$ ,  $g \circ f : I'_1 \rightarrow I'_3$  is a partial matching between  $D_1$  and  $D_3$  as both  $f$  and  $g$  are bijective. Computing the cost of the matchings for  $p < \infty$ , we notice that

$$\begin{aligned} &\sum_{i \in I'_1} d_\infty(D_1(i), D_2(f(i))) + \sum_{i \in I_1 \setminus I'_1} d_\infty(D_1(i), \Delta) + \sum_{i \in I_2 \setminus I'_2} d_\infty(D_2(i), \Delta) \\ &+ \sum_{i \in I'_2} d_\infty(D_2(i), D_3(g(i))) + \sum_{i \in I_2 \setminus I'_2} d_\infty(D_2(i), \Delta) + \sum_{i \in I_3 \setminus I'_3} d_\infty(D_3(i), \Delta) \\ &\geq \sum_{i \in I'_1} d_\infty(D_1(i), D_3(g \circ f(i))) + \sum_{i \in I_1 \setminus I'_1} d_\infty(D_1(i), \Delta) + \sum_{i \in I_3 \setminus I'_3} d_\infty(D_3(i), \Delta) \end{aligned}$$

as  $d_\infty(D_1(i), D_2(f(i))) + d_\infty(D_2(f(i)), D_3(g(f(i)))) \geq d_\infty(D_1(i), D_3(g \circ f(i)))$  using the triangle inequality of  $d_\infty$ . Therefore, for all partial matchings  $f$  and  $g$  as described, we have  $\text{cost}_p(f) + \text{cost}_p(g) \geq \text{cost}_p(g \circ f)$ . Using the same reasoning, for  $p = \infty$  we also obtain  $\text{cost}_\infty(f) + \text{cost}_\infty(g) \geq \text{cost}_\infty(g \circ f)$ . Hence, we have verified that

$$\omega_p(D_1, D_2) + \omega_p(D_2, D_3) \geq \omega_p(D_1, D_3).$$

□

## 1.5 The Hausdorff and Gromov-Hausdorff distances

The Hausdorff distance is a way of measuring distances of different sets contained into a same metric space. This concept can be generalized defining a metric which allow us to measure distances between different metric spaces.

**Definition 1.5.1** (Hausdorff distance). Let  $(M, d)$  be a metric space, and let  $A \subseteq M$ ,  $B \subseteq M$  two compact subspaces of  $M$ . Define the  $r$ -**neighborhood** of a set  $S \subset M$  as

$$U_r(S) := \{x \in S \mid d(x, S) \leq r\}.$$

The **Hausdorff distance** can be defined as

$$d_H(A, B) := \inf \{r > 0 \mid A \subset U_r(B) \text{ and } B \subset U_r(A)\}.$$

**Definition 1.5.2** (Isometric metric spaces). Let  $(X, d_X), (Y, d_Y)$  be metric spaces.  $X$  and  $Y$  are said to be **isometric** if there exists a bijective map  $f : X \rightarrow Y$  such that distances are preserved. That is, for all  $x_1, x_2 \in X$ ,

$$d_X(x_1, x_2) = d_Y(f(x_1), f(x_2)).$$

**Definition 1.5.3** (Gromov-Hausdorff distance). Let  $(X, d_X), (Y, d_Y)$  be metric spaces. The **Gromov-Hausdorff distance** is defined as

$$d_{GH} := \inf \{r > 0 \mid \exists (Z, d_Z) \text{ metric space such that, } \exists X', Y' \subseteq Z, d_H(X', Y') < r\},$$

where  $X', Y'$  are isometric spaces to  $X$  and  $Y$  respectively.

**Lemma 1.5.4** (Proposition 7.3.16, [4]). *Gromov-Hausdorff distance satisfy the triangle inequality. That is, for any metric spaces  $X_1, X_2, X_3$  it is verified that*

$$d_{GH}(X_1, X_3) \leq d_{GH}(X_1, X_2) + d_{GH}(X_2, X_3).$$

*Proof.* Let  $d_{12}$  be a metric over  $X_1 \cup X_2$  and let  $d_{23}$  be a metric over  $X_2 \cup X_3$ . Over  $X_1 \cap X_3$ , define

$$d_{13} := \begin{cases} d_{X_1}(x_1, x_3) & \text{if } x_1, x_3 \in X_1, \\ d_{X_2}(x_1, x_3) & \text{if } x_1, x_3 \in X_3, \\ \inf_{x_2 \in X_2} \{d_{12}(x_1, x_2) + d_{23}(x_2, x_3)\} & \text{if } x_1 \in X_1, x_3 \in X_3. \end{cases}$$

For the first two cases we clearly have a metric. For the third one observe that taking  $x_1 \in X_1, x_3 \in X_3$  and some  $x \in X_1$  we have

$$\begin{aligned} d_{13}(x_1, x) + d_{13}(x, x_3) &= d_{X_1}(x_1, x) + \inf_{x_2 \in X_2} \{d_{12}(x, x_2) + d_{23}(x_2, x_3)\} \\ &= \inf_{x_2 \in X_2} \{d_{12}(x_1, x) + d_{12}(x, x_2) + d_{23}(x_2, x_3)\} \\ &\geq \inf_{x_2 \in X_2} \{d_{12}(x_1, x_2) + d_{23}(x_2, x_3)\} \\ &= d_{13}(x_1, x_3). \end{aligned}$$

This implies, taking the corresponding metric  $d_{ij}$  where  $i, j = 1, 2, 3$ , that

$$d_H(X_1, X_3) \leq d_H(X_1, X_2) + d_H(X_2, X_3),$$

and, taking the infimum over the metrics  $d_{12}$  and  $d_{23}$  we have

$$d_{GH}(X_1, X_3) \leq d_{GH}(X_1, X_2) + d_{GH}(X_2, X_3).$$

□

To check that Gromov-Hausdorff distance is actually a distance we first give a useful characterization in 1.5.7. It is expressed in terms of correspondance distortions.

**Definition 1.5.5** (Correspondance between sets). Given two sets  $X$  and  $Y$ , a **correspondance** between them is a set  $R \subseteq X \times Y$  verifying that for every  $x \in X$ , there exists at least one  $y \in Y$  such that  $(x, y) \in R$  and, for every  $y \in Y$ , there exists an  $x \in X$  such that  $(x, y) \in R$ .

**Definition 1.5.6** (Distortion of a correspondance). Let  $(X, d_X), (Y, d_Y)$  be two metric spaces, and let  $R$  be a correspondance between them. The **distortion** of  $R$  is defined as

$$\text{dis}(R) := \sup \{|d_X(x, x') - d_Y(y, y')| : (x, y), (x', y') \in R\}.$$

**Proposition 1.5.7** (Theorem 7.3.25, [4]). *Let  $(X, d_X), (Y, d_Y)$  be two metric spaces. The Gromov-Hausdorff distance between them can be characterized as*

$$d_{\text{GH}}((X, d_X), (Y, d_Y)) = \frac{1}{2} \inf_R \text{dis}(R).$$

*Proof.* Given  $r \geq d_{\text{GH}}(X, Y)$ , for some metric space  $(Z, d_Z)$ , we can take  $X', Y' \subseteq Z$  such that  $X'$  and  $Y'$  are isometric embeddings of  $X$  and  $Y$  respectively and  $d_{\text{H}}(X', Y') < r$  in  $Z$ . Thus, we can see every element of  $X$  and  $Y$  as elements of  $Z$  through some isometry. Therefore, we can define the correspondance

$$R := \{(x, y) \in X \times Y : d_Z(x, y) < r\}.$$

The set  $R$  is actually a correspondance because the fact that  $d_{\text{H}}(X', Y') < r$  implies that for every  $x \in X$  and every  $y \in Y$ ,  $d_Z(x, y) < r$ , so every  $x$  and every  $y$  have some correspondance. Now, let  $(x, y), (x', y') \in R$ . Using the triangle inequality of  $Z$  we have

$$\begin{aligned} \text{dis}(R) &\leq |d_X(x, x') - d_Y(y, y')| \\ &= |d_Z(x, x') - d_Z(y, y')| \\ &\leq |d_Z(x, y) + d_Z(y, x') - d_Z(y, y')| \\ &\leq d_Z(x, y) + d_Z(x', y) + d_Z(y, y') \\ &\leq d_Z(x, y) + d_Z(x', y') \leq 2r. \end{aligned}$$

This shows

$$2d_{\text{GH}}(X, Y) \geq \inf_R \text{dis}(R).$$

To see the reverse inequality, let  $R$  be any correspondance, and let's take  $\text{dis}(R) = 2r$ . Let's construct a metric space  $(Z, d_Z)$  formed by the disjoint union of spaces  $Z = X \cup Y$ .

For every  $z_1, z_2 \in Z$ , we define  $d_Z$  as

$$d_Z(z_1, z_2) := \begin{cases} d_X(z_1, z_2) & \text{if } z_1, z_2 \in X, \\ d_Y(z_1, z_2) & \text{if } z_1, z_2 \in Y, \\ \inf\{d_X(z_1, x') + r + d_Y(z_2, y') : (x', y') \in R\} & \text{if } z_1 \in X, z_2 \in Y. \end{cases}$$

By definition, it is clear that  $d_Z$  respects isometrically both  $d_X$  and  $d_Y$ . By the same reason  $d_Z(z_1, z_2) = d_Z(z_2, z_1)$  and  $d_Z(z_1, z_2) \geq 0$  for every  $z_1, z_2 \in Z$ , where  $d_Z(z_1, z_2) = 0$  only if either  $z_1 = z_2$  or  $r = 0$ . To check the triangle inequality we take  $z_1, z_2, z_3 \in Z$ . If either all three are elements of  $X$ , or all three are elements of  $Y$ , the inequality is verified as it is granted in  $X$  and  $Y$  with  $d_X$  and  $d_Y$  respectively. In case  $z_1, z_2 \in X$  and  $z_3 \in Y$  we can take some  $y \in Y$  such that  $(z_2, y) \in R$ . Thus, we have

$$\begin{aligned} d_Z(z_1, z_2) + d_Z(z_2, z_3) &\geq d_X(z_1, z_2) + d_X(z_2, z_2) + r + d_Y(z_3, y) \\ &\geq d_X(z_1, z_2) + r + d_Y(z_3, y) \\ &\geq d_Z(z_1, z_3). \end{aligned}$$

Analogously, the argument follows for  $z_1 \in X$  and  $z_2, z_3 \in Y$ . Thus, all is left to prove is to check  $d_H(X, Y) < r$ .

...

□

**Definition 1.5.8** (Distortion of a map). Let  $(X, d_X), (Y, d_Y)$  be two metric spaces and let  $f: X \rightarrow Y$  an arbitrary map. The **distortion** of  $f$  is defined as

$$\text{dis}(f) := \sup_{x_1, x_2 \in X} |d_Y(d(x_1), f(x_2)) - d_X(x_1, x_2)|.$$

**Definition 1.5.9** ( $\varepsilon$ -isometry). Let  $X$  and  $Y$  be two metric spaces and let  $\varepsilon > 0$ . A  **$\varepsilon$ -isometry** between two metric spaces is a map  $f: X \rightarrow Y$  such that  $\text{dis}(f) \leq \varepsilon$ . The image  $f(X)$  is called an  **$\varepsilon$ -net**.

**Proposition 1.5.10** (Theorem 7.3.28.1, [4]). Let  $X$  and  $Y$  be two metric spaces and let  $\varepsilon > 0$ . If  $d_{\text{GH}} < \varepsilon$ , then there exists a  $2\varepsilon$ -isometry from  $X$  to  $Y$ .

*Proof.* Let  $R$  be a correspondance between  $X$  and  $Y$ . By 1.5.7 it holds that  $\text{dis}(R) < 2\varepsilon$ . For every  $x \in X$  we choose some  $y \in Y$  such that  $(x, y) \in R$  and define  $f(x) := y$ . This defines a map  $f: X \rightarrow Y$ . We then have

$$\text{dis}(f) \leq \text{dis}(R) < 2\varepsilon.$$

...

□

Up to this moment we have seen that Gromov-Hausdorff distance defines a pseudo-metric over the set of metric spaces. Note that if  $X$  and  $Y$  are isometric, directly of the definition we get,  $d_{\text{GH}}(X, Y) = 0$ . To make Gromov-Hausdorff distance an actual metric we need to ask one more thing to our metric spaces. That is, to be compact. Denote

$$\mathcal{X} := \{(X, d_X) : (X, d_X) \text{ is a metric compact space}\}.$$

**Theorem 1.5.11** (Theorem 7.3.30, [4]). *Gromov-Hausdorff distance is in fact a metric over the space of isometry classes of compact metric spaces.*

*Proof.* We just seen that if  $X$  and  $Y$  are isometric, directly of the definition we get,  $d_{\text{GH}}(X, Y) = 0$ . By definition, Gromov-Hausdorff distance is nonnegative and symmetric and, by Lemma 1.5.4, it verifies the triangle inequality. It only remains to prove that given two metric spaces  $X, Y \in \mathcal{X}$ , if  $d_{\text{GH}}(X, Y) = 0$  then  $X$  and  $Y$  are isometric.

Let  $X, Y \in \mathcal{X}$  such that  $d_{\text{GH}}(X, Y) = 0$ . By Proposition 1.5.10, there exists a sequence of maps  $f_n : X \rightarrow Y$  such that  $\text{dis}(f_n) \rightarrow 0$ . As  $X$  is compact, we can fix a countable dense set  $S \subset X$ .

...

□

In order to extend the scope of Gromov-Hausdorff distance we can endow our compact metric spaces with real-valued functions, which will still maintain good stability properties as we will see in Chapter 5.

Denote the collection of such spaces as

$$\mathcal{X}_1 := \{(X, d_X, f) : (X, d_X) \in \mathcal{X}, f_X : X \rightarrow \mathbb{R} \text{ continuous}\}.$$

**Definition 1.5.12.** Let  $X, Y \in \mathcal{X}_1$ . We extend the **Gromov-Hausdorff distance over  $\mathcal{X}_1$**  as

$$d_{\text{GH}}^1((X, d_X, f_X), (Y, d_Y, f_Y)) = \inf_R \max \left\{ \frac{1}{2} \text{dis}(R), \|f_X - f_Y\|_{\ell^\infty} \right\}.$$

An analogous adaptation of Theorem 1.5.11 and the previous results proofs that  $d_{\text{GH}}^1$  defines a metric over the set of isomorphism classes of  $\mathcal{X}_1$ .

## Chapter 2

# Structure Theorem

The Structure Theorem for persistence modules is referred to as the “first miracle” of persistence homology [13]. This algebraic property allows to express a persistence module of finite type as a direct sum of finitely many interval modules. This enables to express a persistence module by the bars in a barcode, which latter allows to compare bottleneck distance between the barcodes, with the interleaving distance between the modules.

The proof of the theorem requires the algebraic structure theorem for finitely generated modules over a principal domain. Section 2.1 introduces the algebraic structure theorem for finitely generated modules over a principal ideal domain. Section 2.2 then gives a proof of the Structure Theorem for persistence diagrams.

### 2.1 Algebraic structure theorem

**Definition 2.1.1** (Graded ring). Let  $R$  be a ring. It is said that  $R$  is a **graded ring** if it can be decomposed into a direct sum of additive groups

$$R = \bigoplus_{n=1}^{\infty} R_n = R_1 \oplus R_2 \oplus \dots$$

such that for all  $n, m \geq 0$ ,

$$R_n \cdot R_m \subseteq R_{n+m}.$$

**Example 2.1.2** (Example 4.4, [16]). Let  $\mathbb{F}$  be a field, the polynomial ring over it,  $\mathbb{F}[x]$ , is a graded ring. It can be decomposed as

$$\mathbb{F}[x] = \bigoplus_{i=0}^{\infty} x^i \cdot \mathbb{F} = \bigoplus_{i=0}^{\infty} \{cx^i \mid c \in \mathbb{F}\}.$$

Also, polynomial multiplication verifies the degree of the product of two monomials is the sum of the degrees of the factors.

**Definition 2.1.3** (Graded ideal). Let  $R$  be a graded ring. A **graded ideal** is a two sided ideal  $I \subseteq R$  that can be decomposed into a direct sum

$$I = \bigoplus_{n=0}^{\infty} I_n$$

where for each  $n \geq 0$ ,  $I_n = I \cap R_n$ .

**Example 2.1.4.** Note that in Example 2.1.2, every ideal is a graded ideal. No see an example of non graded ideals take the two-dimensional polynomial ring over a field,  $R = \mathbb{F}[x, y]$ . There are also many graded ideals in  $R$ , take, for example, the ideal generated by the two dimensional monomials  $I = \langle x^2, xy \rangle$ .

Take now an ideal generated by non homogeneous elements as  $I' = \langle x^2 + y \rangle$ . Then it happens that  $x^2 \in I'_2 = I' \cap R^2$  and  $y \in I'_1 = I' \cap R^1$ , but either  $x^2 \in I'$  nor  $y \in I'$ .

**Definition 2.1.5** (Graded module, Definition 4.7 [16]). Let  $M$  be a left module over a graded ring  $R$ . It is said that  $M$  is a **left graded module** if it can be decomposed into a direct sum

$$M = \bigoplus_{n=1}^{\infty} M_n$$

if for each  $n, m \geq 0$ ,  $R_n M_m \subseteq M_{n+m}$ .

**Theorem 2.1.6** (Chapter IV, Theorem 6.12, [11]). *Let  $M$  be a finitely generated module over a principal ideal domain  $R$ . There exist a finite sequence of proper ideals  $(d_1) \supseteq (d_2) \supseteq \dots \supseteq (d_n)$  such that*

$$M \cong \bigoplus_{i=1}^n R/(d_i).$$

Due to the lengthy concepts needed to prove it, we will refer to it as a well known fact. An introduction to module theory and a detailed proof of the theorem of Theorem 2.1.6 can be found at [11, Chapter IV].

## 2.2 Structure theorem for persistence diagrams

In addition to Theorem 2.1.6, we will use the following simple algebraic statement.

**Proposition 2.2.1** (Proposition 4.6, [16]). *An ideal  $I \subseteq R$  is graded if and only if it is generated by homogeneous elements.*

*Proof.* First, if  $I$  is a graded ideal  $I = \bigoplus_p I^p$  and is generated by  $\bigcup_p I^p$ . Then, each

$$I^p = I \cap R^p \subseteq R^p$$

is a subset of homogeneous elements. Therefore,  $I$  is generated by homogeneous elements.

Now, let  $I$  be generated by a set  $X$  of homogeneous elements. For sure,  $I \cap R^p \subseteq I$ , so we just need to prove the converse inclusion. As  $I$  is generated by  $X$ , its elements  $u \in I$  are of the form

$$u = \sum_i r_i x_i s_i, \quad (2.1)$$

for  $r_i, s_i \in R$  and  $x_i \in X$ . And as  $I \subseteq R$ , also,

$$u = \sum_p u_p,$$

for  $u_p \in R^p$ . For every term in (2.1), we have

$$r_i = \sum_j r_{i,j}, \quad s_i = \sum_l s_{i,l},$$

with each  $r_{i,j}, s_{i,l}$  being homogeneous. Therefore, combining all we have that

$$u = \sum_i \sum_{j,l} r_{i,j} x_i s_{i,l}. \quad (2.2)$$

Each term in (2.2) is homogeneous as is a product of homogeneous elements. Thus  $u_p$  is the sum of those terms, and  $u$  has degree  $p$ . Therefore  $u_p \in I$  and  $I \subseteq I \cap R^p$ .  $\square$

**Theorem 2.2.2** (Proposition 4.8, [16]). *Let  $(V, \pi)$  be a persistence module. There exist a barcode  $\text{Bar}(V, \pi)$ , with  $\mu: \text{Bar}(V, \pi) \rightarrow \mathbb{N}$ , the multiplicity of the barcode intervals, such as there is a unique direct sum decomposition*

$$V \cong \bigoplus_{I \in \text{Bar}(V)} \mathbb{F}(I)^{\mu(I)}. \quad (2.3)$$

*Proof.*  $V$  is of finite type, so it is a finite  $\mathbb{F}[x]$ -module. As  $\mathbb{F}$  is a field,  $\mathbb{F}[x]$  is a principal ideal domain, therefore,  $V$  is a finitely generated module over a principal ideal domain. Using Fact 2.1.6,  $V$  can be decompose in the direct sum of its free and torsion subgroups,  $F \oplus T$ . Thus, we have

$$F = \bigoplus_{i \geq q} x^i \cdot \mathbb{F}$$

$$T = \bigoplus_{i \geq q} R^i / I^i.$$



Each  $x^i \cdot \mathbb{F}$  is isomorphic to ideals of the form  $(x^q)$ . By Proposition 2.2.1, each  $R^i/I^i$  is isomorphic to some quotient of graded ideals of the form  $(x^p)/(x^r)$ . Note that the free subgroup can be seen as a particular case of the torsion group taking  $r = 0$ . Thus  $V$  can be decompose as described in (2.3).  $\square$

## Chapter 3

# Interleaving Stability Theorem

This Chapter gives a detailed proof of the first stability theorem for persistence homology. This theorem is referred to as the “Geometry miracle” of persistent homology, as it allows to describe an isometry between persistence modules and barcodes [13]. This shows that *small* changes in a data sets will perform *small* changes in their persistence modules, and therefore small changes in how persistent homology groups vary through time. The theorem claims that given two persistence modules, the distance between them using the interleaving distance, is the same as the distance between their barcodes using the bottleneck distance.

**Theorem 3.0.1** (Interleaving Stability, Theorem 2.2.8, [14]). *There is an isometry between a persistence module with the interleaving distance and its barcode with the bottleneck distance. That is, given two persistence modules  $V$  and  $W$ , it holds that*

$$d_{\text{int}}(V, W) = d_{\text{bot}}(\text{Bar}(V), \text{Bar}(W)).$$

For the presented proof we have followed [14]. Hence, we will divide the proof into proving two complementary inequalities separately. This implies checking that if there exists a  $\delta$ -matching between two given barcodes, then there exists a  $\delta$ -interleaving morphism between their persistence modules, Proposition 3.1.2. Also, we need to check that, if there exists a  $\delta$ -interleaving morphism between two persistence modules, then there exists a  $\delta$ -matching between their barcodes, 3.2.8.

### 3.1 First inequality

The first claim can be deduced from the Structure Theorem in a rather direct way, proving first the case where our modules are just interval modules.

**Lemma 3.1.1** (Exercise 2.2.7, [14]). *Let  $I, J$  be two  $\delta$ -matched intervals. Then, their corresponding interval modules  $(\mathbb{F}(I), \pi)$  and  $(\mathbb{F}(J), \theta)$  are  $\delta$ -interleaved.*

*Proof.* Let  $I = (a, b]$ ,  $J = (c, d]$ . If  $\rho$  is the  $\delta$ -matching between them, then  $\rho(I) = J$  and, following Definition 1.3.3,  $(a, b] \subseteq (c - \delta, d + \delta]$  and  $(c, d] \subseteq (a - \delta, b + \delta]$ , with  $b - a > 2\delta$  and  $d - c > 2\delta$ . Then, the morphisms

$$\begin{aligned} \phi_\delta: \mathbb{F}(I) &\rightarrow \mathbb{F}(J)_\delta & \text{and} & & \psi_\delta: \mathbb{F}(J) &\rightarrow \mathbb{F}(I)_\delta \\ \phi_\delta(\mathbb{F}(I)_t) &\mapsto \mathbb{F}(J)_{(t+\delta)} & & & \psi_\delta(\mathbb{F}(J)_t) &\mapsto \mathbb{F}(I)_{(t+\delta)} \end{aligned}$$

are well defined as for any  $t \in (a, b]$ ,  $t + \delta \in (c, d]$ , as  $a + \delta > c$  and  $b + \delta \leq d$ . In the same way, for any  $t \in (c, d]$ ,  $t + \delta \in (a, b]$ . Thus,

$$\psi_\delta \circ \phi_\delta(\mathbb{F}(I)_t) = \psi_\delta(\mathbb{F}(J)_{(t+\delta)}) = \mathbb{F}(I)_{(t+2\delta)} = \pi_{t \leq t+2\delta}(\mathbb{F}(I)_t).$$

and

$$\phi_\delta \circ \psi_\delta(\mathbb{F}(J)_t) = \phi_\delta(\mathbb{F}(I)_{(t+\delta)}) = \mathbb{F}(J)_{(t+2\delta)} = \theta_{t \leq t+2\delta}(\mathbb{F}(J)_t).$$

Therefore,  $\phi_\delta$  and  $\psi_\delta$  are a pair of  $\delta$ -interleaving morphisms.  $\square$

Once we are able to build a  $\delta$ -interleaving between two  $\delta$ -matched interval modules, we will use the Structure Theorem for persistence modules to generalize the construction for arbitrary persistence modules. This will be useful to prove the first inequality needed to prove Theorem 3.0.1.

**Proposition 3.1.2** (Theorem 3.0.1, [14]). *Given two persistence modules  $V$ ,  $W$ , if there is a  $\delta$ -matching between their barcodes, then there is a  $\delta$ -interleaving morphism between them.*

*Proof.* Suppose that  $\rho: \text{Bar}(V) \rightarrow \text{Bar}(W)$  is a  $\delta$ -matching between the barcodes of  $V$  and  $W$ . By the Structure Theorem 2.1.6,  $V$  and  $W$  decompose in a finite direct sum of interval modules

$$V \cong \bigoplus_{I \in \text{Bar}(V)} \mathbb{F}(I), \quad W \cong \bigoplus_{J \in \text{Bar}(W)} \mathbb{F}(J).$$

We can express  $V = V_Y \oplus V_N$ ,  $W = W_Y \oplus W_N$  denoting

$$\begin{aligned} V_Y &\cong \bigoplus_{I \in \text{coim } \rho} \mathbb{F}(I), & V_N &\cong \bigoplus_{I \in \text{Bar}(V) \setminus \text{coim } \rho} \mathbb{F}(I), \\ W_Y &\cong \bigoplus_{J \in \text{im } \rho} \mathbb{F}(J), & W_N &\cong \bigoplus_{J \in \text{Bar}(W) \setminus \text{im } \rho} \mathbb{F}(J). \end{aligned}$$

The  $V_Y, W_Y$  modules separate the “yes, matched” intervals, from the  $V_N, W_N$  “not matched” intervals. For every interval  $I \in \text{Bar}(V_Y)$ ,  $I$  is  $\delta$ -matched to an interval

$J \in \text{Bar}(W_Y)$  by  $\rho(I) = J$ . Thus, by Lemma 3.1.1, for all pair  $I, J$  of matched intervals, there exist a pair of  $\delta$ -interleaved morphisms

$$\phi_\delta: \mathbb{F}(I) \rightarrow \mathbb{F}(J)_\delta \quad \text{and} \quad \psi_\delta: \mathbb{F}(J) \rightarrow \mathbb{F}(I)_\delta$$

which induce the pair of  $\delta$ -interleaved morphisms

$$\phi_\delta: V_Y \rightarrow W_{Y\delta} \quad \text{and} \quad \psi_\delta: W_Y \rightarrow V_{Y\delta}.$$

Not matched intervals are of length smaller than  $2\delta$ , therefore both,  $V_N$  and  $V_Y$  are  $\delta$ -interleaved with the empty set. We can now construct the  $\delta$ -interleaving morphism  $\phi: V \rightarrow W$  such as  $\phi|_{V_Y} = \phi_Y$  and  $\phi|_{V_N} = 0$ . In a similar way, we also construct  $\psi: W \rightarrow V$ .  $\square$

## 3.2 Second inequality

With Proposition 3.1.2 we have proven the first half of Stability Theorem 3.0.1. Now we need to prove that we can build a  $\delta$ -interleaving morphism from a  $\delta$ -matching. To verify this claim we need several previous lemmas that will lead us to prove Proposition 3.1.2.

First, we will introduce some notation. Let  $(V, \pi), (W, \theta)$  be two persistence modules and let  $I = (b, d]$  be an interval with  $d \in \mathbb{R} \cup \{+\infty\}$ . Denote the set of bars of  $\text{Bar}(V)$  that start before  $b$  and end exactly at  $d$  as

$$\text{Bar}_{I-}(V) := \{(a, d] \in \text{Bar}(V) : a \leq b\}.$$

Analogously, denote the set of bars that start at  $b$  and end after  $d$  as

$$\text{Bar}_{I+}(V) := \{(b, c] \in \text{Bar}(V) : c \geq d\}.$$

**Lemma 3.2.1** (Proposition 3.1.1, [14]). *Let  $I = (b, d]$  be an interval. It exists an injective morphism  $\iota: (V, \pi) \rightarrow (W, \theta)$ , then  $\#(\text{Bar}_{I-}(V)) \leq \#(\text{Bar}_{I-}(W))$ . Where  $\#(\cdot)$  denotes the cardinal operator.*

*Proof.* For  $b < s < d < r$ , denote the set of elements in  $V_d$  which come from all  $V_s$  and disappear in all  $V_r$  as

$$E_{I-} = \bigcap_{b < s < d} \text{im } \pi_{s \leq d} \cap \bigcap_{r > d} \ker \pi_{d \leq r} \subseteq V_d.$$

It holds that  $\dim E_{I-}(V) = \#(\text{Bar}_{I-}(V))$ . For every morphism  $p: (V, \pi) \rightarrow (W, \theta)$  the following diagram commutes

$$\begin{array}{ccc} V_s & \xrightarrow{\pi_{s \leq r}} & V_r \\ p_s \downarrow & & \downarrow p_r \\ W_s & \xrightarrow{\theta_{s \leq r}} & W_r \end{array}$$

This implies that

$$p_r(\operatorname{im} \pi_{s \leq r}) \subseteq \operatorname{im} \theta_{s \leq r}, \quad p_s(\ker \pi_{s \leq r}) \subseteq \ker \theta_{s \leq r}.$$

Taking  $r = d$ ,  $b < s < d$  in the first inclusion, and  $s = d$ ,  $r > d$  in the second, we have that

$$p_d(\operatorname{im} \pi_{s \leq d}) \subseteq \operatorname{im} \theta_{s \leq d}, \quad p_d(\ker \pi_{d \leq r}) \subseteq \ker \theta_{d \leq r},$$

and

$$p_d(E_{I-}(V)) \subseteq E_{I-}(W).$$

If we now take  $p$  as the injective morphism of the hypothesis,  $p = \iota$ , we get

$$\dim E_{I-}(V) \leq \dim E_{I-}(W).$$

□

**Lemma 3.2.2** (Exercise 3.1.3, [14]). *Let  $I = (b, d]$  be an interval. It exists a surjective morphism  $s: (V, \pi) \rightarrow (W, \theta)$ , then  $\#(\operatorname{Bar}_{I+}(V)) \geq \#(\operatorname{Bar}_{I+}(W))$ .*

*Proof.* Analogously to the proof of Lemma 3.2.1 we now define

$$E_{I+}(V) = \bigcap \operatorname{im} \pi_{d \leq r}.$$

Therefore  $\dim E_{I+}(V) = \#(\operatorname{Bar}_{I+}(V))$ , and recalling the diagram used for the previous proof, and using the fact that is commutative, we have that

$$p_r(\operatorname{im} \pi_{s \leq r}) \supseteq \operatorname{im} \theta_{s \leq r}.$$

Taking  $s = d$  we then have that

$$p_d(E_{I+}(V)) \supseteq E_{I+}(W).$$

And finally, taking the surjective morphism  $p = s$  we have that

$$\dim E_{I-}(V) \geq \dim E_{I-}(W).$$

□

To build our  $\delta$ -matching we first define two induced matchings, by an injection and by a surjection respectively. First, suppose that there exists an injection  $\iota: V \rightarrow W$ . For every  $c \in \mathbb{R} \cup \{\infty\}$ , sort the bars  $(a_i, c] \in \operatorname{Bar}(V)$ ,  $i \in \{1, \dots, k\}$  by decreasing length order,

$$(a_1, c] \supseteq (a_2, c] \supseteq \dots \supseteq (a_k, c], \text{ with } a_1 \leq a_2 \leq \dots \leq a_k.$$

Sort in the same manner the bars  $(b_j, c] \in \text{Bar}(V)$ ,  $j \in \{1, \dots, l\}$ ,

$$(b_1, c] \supseteq (b_2, c] \supseteq \dots \supseteq (b_k, c], \text{ with } b_1 \leq b_2 \leq \dots \leq b_k.$$

As there is an injection between  $V$  and  $W$ , Lemma 3.2.1 assures that the amount of bars in  $\text{Bar}(V)$  is lower than the amount in  $\text{Bar}(W)$ , i.e.,  $k \leq l$ . We define the *injective induced matching*  $\mu_{inj}: \text{Bar}(V) \rightarrow \text{Bar}(W)$  matching, for each  $c \in \mathbb{R} \cup \{\infty\}$ , the intervals from both lists by decreasing length.

**Lemma 3.2.3** (Proposition 3.1.5, [14]). *If there exists an injection  $\iota: (V, \pi) \rightarrow (W, \theta)$ , then the induced matching  $\mu_{inj}: \text{Bar}(V) \rightarrow \text{Bar}(W)$  satisfies:*

1.  $\text{coim } \mu_{inj} = \text{Bar}(V)$ ,
2.  $\mu_{inj}(a, c] = (b, c], \forall b \leq a, \forall (a, d] \in \text{Bar}(V)$ .

*Proof.* Applying Lemma 3.2.1 with the interval  $(a_k, c]$ , we have that for each  $c \in \mathbb{R} \cup \{\infty\}$ ,  $\# \text{Bar}_{(a_k, c]}(V) \leq \# \text{Bar}_{(a_k, c]}(W)$ , having  $k \leq l$  as we note earlier. This means that every bar in  $\text{Bar}(V)$  is matched to some bar in  $\text{Bar}(W)$ . Hence  $\text{coim } \mu_{inj} = \text{Bar}(V)$ . Moreover, as the matching is carried out in length descending order, for each  $i \in \{1, \dots, k\}$ ,  $\mu_{inj}(a_i, c] = (b_i, c]$ , and applying Lemma 3.2.1, now with the interval  $(a_i, c]$ , and making the same reasoning,  $a_i \leq b_i$ .  $\square$

Now we suppose that there exists a surjection  $\sigma: V \rightarrow W$ . For every  $a \in \mathbb{R}$ , sort the intervals  $(a, c_i] \in \text{Bar}(V)$ ,  $i \in \{1, \dots, k\}$  by decreasing length order as before,

$$(a, c_1] \supseteq (a, c_2] \supseteq \dots \supseteq (a, c_k], \text{ with } c_1 \geq c_2 \geq \dots \geq c_k,$$

and again in the same manner, sort the intervals  $(a, d_j] \in \text{Bar}(V)$ ,  $j \in \{1, \dots, l\}$ ,

$$(a, d_1] \supseteq (a, d_2] \supseteq \dots \supseteq (a, d_l], \text{ with } d_1 \geq d_2 \geq \dots \geq d_l.$$

We define the *surjective induced matching*  $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(W)$  matching, for each  $a \in \mathbb{R}$ , the intervals from both lists by decreasing length.

**Lemma 3.2.4** (Exercise 3.1.8, [14]). *If there exists a surjection  $s: (V, \pi) \rightarrow (W, \theta)$ , then the induced matching  $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(W)$  satisfies:*

1.  $\text{im } \mu_{sur} = \text{Bar}(W)$ ,
2.  $\mu_{sur}(a, c] = (a, d], \forall c \geq d, \forall (a, d] \in \text{Bar}(V)$ .

*Proof.* Using Lemma 3.2.2 with the interval  $(b, d_k]$  for each  $b \in \mathbb{R}$ , we get that, as there exists a surjection between the modules, now  $k \geq l$ . Therefore, every bar in  $\text{Bar}(W)$  is matched to some bar in  $\text{Bar}(V)$  and  $\text{im } \mu_{sur} = \text{Bar}(W)$ . In an analogue way to the previous lemma, as the intervals in both lists are matched in a decreasing manner, for every  $j \in \{1, \dots, l\}$ ,  $\mu_{sur}(a, c_j] = (a, d_j]$ , and if we now apply Lemma 3.2.2, we get that  $c_j \geq d_j$ .  $\square$

Hence, with the injective and the surjective induced matchings at hand, for a general morphism  $f$ , we can define the *induced matching*  $\mu(f): \text{Bar}(V) \rightarrow \text{Bar}(W)$ , as the composition  $\mu_{inj} \circ \mu_{sur}$ , defined as  $\text{im } \mu_{sur} = \text{Bar}(\text{im } f) = \text{coim } \mu_{inj}$ .

The following lemma verifies that, in fact, the mapping between persistence modules with its morphisms and barcodes with induced matchings (either the injective or the surjective versions) has functorial properties between the two categories.

**Lemma 3.2.5** (Claim 3.1.13, [14]). *Let  $U, V$  and  $W$  persistence diagrams and  $f, g, h$  morphisms between them defined as in the following diagram:*

$$\begin{array}{ccccc} U & \xrightarrow{f} & V & \xrightarrow{g} & W \\ & & \searrow & \nearrow & \\ & & h & & \end{array} .$$

*If all  $f, g, h$  are all injections, or all surjections, then the corresponding diagram formed by the barcodes of the modules, and their respective matchings commutes as well.*

$$\begin{array}{ccccc} \text{Bar}(U) & \xrightarrow{\mu_*(f)} & \text{Bar}(V) & \xrightarrow{\mu_*(g)} & \text{Bar}(W) \\ & & \searrow & \nearrow & \\ & & \mu_*(h) & & \end{array} .$$

Where  $\mu_*$  denotes  $\mu_{inj}$  or  $\mu_{sur}$  accordingly.

*Proof.* Let  $f, g, h$  injective morphisms, by the definition of the injective induced matching and Lemma 3.2.1 for any  $d \in \mathbb{R} \cup \{+\infty\}$ , there exist  $k \leq l \leq q$  such that the barcodes of  $U, V, W$  consist on the following bars:

$$\begin{aligned} \text{Bar}(U) &: (a_1, d] \supset \dots \supset (a_k, d] \\ \text{Bar}(V) &: (b_1, d] \supset \dots \supset (b_k, d] \supset \dots \supset (b_l, d] \\ \text{Bar}(W) &: (c_1, d] \supset \dots \supset (c_k, d] \supset \dots \supset (c_l, d] \supset \dots \supset (c_q, d]. \end{aligned}$$

Therefore, for any  $d$  the diagram commutes as

$$\mu_{inj}(f)(a_i, d] = (b_i, d], \quad \mu_{inj}(g)(b_i, d] = (c_i, d], \quad \mu_{inj}(h)(a_i, d] = (c_i, d]$$

for  $1 \leq i \leq k$ . If  $f, g, h$  were surjective morphisms, an analogue reasoning using the surjective induced matching definition and Lemma 3.2.2 completes the proof.  $\square$

Finally, we can claim the two main lemmas from which we will construct our desired  $\delta$ -matching.

**Lemma 3.2.6** (Lemma 3.2.1, [14]). *Let  $(V, \pi), (W, \theta)$  be  $\delta$ -interleaved persistence modules, with  $\delta$ -interleaving morphisms  $\phi: V \rightarrow W_\delta$  and  $\psi: W \rightarrow V_\delta$ . Let  $\phi: V \rightarrow \text{im } \phi$  be a surjection and  $\mu_{sur}: \text{Bar}(V) \rightarrow \text{Bar}(\text{im } \phi)$  the induced matching. Then*

1.  $\text{coim } \mu_{sur} \supseteq \text{Bar}(V)_{\geq 2\delta}$ ,
2.  $\text{im } \mu_{sur} = \text{Bar}(\text{im } \phi)$  and
3.  $\mu_{sur}(b, d] = (b, d']$ ,  $\forall (b, d] \in \text{coim } \mu_{sur}$ ,  $d' \in [d - 2\delta, d]$ .

*Proof.* 1. To check the first part, we observe that, in the following diagram, the three morphisms are surjective as  $\phi$  and  $\pi_{t \leq t+2\delta}$  are defined onto their images, and the diagram commutes,

$$\begin{array}{ccccc} V & \xrightarrow{\phi} & \text{im } \phi & \xrightarrow{\psi_\delta} & \text{im } \pi_{t \leq t+2\delta} \\ & \searrow & & \nearrow & \\ & & \pi_{t \leq t+2\delta} & & \end{array}.$$

Therefore, because of Lemma 3.2.5 the barcode diagram also commutes:

$$\begin{array}{ccccc} \text{Bar}(V) & \xrightarrow{\mu_{sur}(\phi)} & \text{Bar}(\text{im } \phi) & \xrightarrow{\mu_{sur}(\psi_\delta)} & \text{Bar}(\text{im } \pi_{t \leq t+2\delta}) \\ & \searrow & & \nearrow & \\ & & \mu_{sur}(\pi_{t \leq t+2\delta}) & & \end{array}.$$

By the definition of the surjective induced matching,

$$\text{coim } \mu_{sur}(\pi_{t \leq t+2\delta}) = \text{Bar}(V)_{\geq 2\delta}.$$

For each starting point  $a \in \mathbb{R}$ , we have that

$$\text{Bar}(\text{im } \pi_{t \leq t+2\delta}) = \{(a, b - 2\delta] : (a, b] \in \text{Bar}(V), b - a > 2\delta\}.$$

Sorting all bars of  $\text{Bar}(V)$  and of  $\text{Bar}(\text{im } \pi_{t \leq t+2\delta})$  in length-not-increasing order and matching the bars though the longest-first order, each bar  $(a, b] \in \text{Bar}(V)$  is matched with the bar  $(a, b - 2\delta] \in \text{Bar}(\text{im } \pi_{t \leq t+2\delta})$  while  $b - a > 2\delta$ . The smaller bars are not matched. Thus,

$$\text{coim } \mu_{sur}(\phi) \supseteq \text{coim } \mu_{sur}(\text{im } \pi_{t \leq t+2\delta}) = \text{Bar}(V)_{\geq 2\delta}.$$

2. The second part is just a reformulation of Lemma 3.2.1.



3. Let  $(b, d] \in \text{coim}$ . There are two cases:

On one hand, if  $d - b \leq 2\delta$ ,  $(b, d]$  is matched to  $(b, d']$  where  $d \geq d'$ , by definition of  $\mu_{sur}$ . Also,  $d' > b$  and, as in this case we have  $b \geq d - 2\delta$ , we have  $d' > d - 2\delta$ . Therefore,  $d' \in [d - 2\delta, d]$ .

On the other hand, if  $d - b > 2\delta$ ,  $(b, d]$  is matched to  $(b, d']$  by  $\mu_{sur}(\phi)$ , with  $(b, d'] \in W_{\leq 2\delta}$ . We can therefore use Lemma 3.2.4 to check that  $d' \geq d$ . In the same manner,  $(b, d']$  is matched to  $(b, d'']$  by  $\mu_{sur}(\psi)_\delta$  with  $d'' \geq d'$ . Finally, using the commutativity of the following diagram, we have that  $(b, d'') = (b, d - 2\delta]$ , making  $d' \in [d - 2\delta, d]$ .

$$\begin{array}{ccccc}
 \text{Bar}(V)_{\geq 2\delta} & & \text{Bar}(\text{im } \phi) & & \text{Bar}(\text{im } \pi_{t \leq t+2\delta}) \\
 \Psi & & \Psi & & \Psi \\
 (b, d] & \xrightarrow{\mu_{sur}(\phi)} & (b, d'] & \xrightarrow{\mu_{sur}(\psi)_\delta} & (b, d''] \\
 & & & & \parallel \\
 & \searrow \mu_{sur}(\pi_{t \leq t+2\delta}) & & & (b, d - 2\delta]
 \end{array}$$

□

**Lemma 3.2.7** (Proposition 3.2.2, [14]). *Let  $(V, \pi), (W, \theta)$  be  $\delta$ -interleaved persistence modules, with  $\delta$ -interleaving morphisms  $\phi: V \rightarrow W_\delta$  and  $\psi: W \rightarrow V_\delta$ . Let  $\phi: V \rightarrow \text{im } \phi$  be an injection and  $\mu_{inj}: \text{Bar}(\text{im } \phi) \rightarrow \text{Bar}(W_\delta)$  the induced matching. Then*

1.  $\text{coim } \mu_{sur} = \text{Bar}(\text{im } \phi)$ ,
2.  $\text{im } \mu_{inj} \supseteq \text{Bar}(W_\delta)_{\geq 2\delta}$  and
3.  $\mu_{inj}(b, d'] = (b', d']$ ,  $(b, d'] \in \text{coim } \mu_{inj}$ ,  $b' \in [b - 2\delta, b]$ .

*Proof.* 1. Immediate using Lemma 3.2.3.

2. As  $\phi_\delta \circ \psi = \theta_{t \leq t+2\delta}$  the following diagram commutes:

$$\begin{array}{ccccc}
 W & \xrightarrow{\psi} & \text{im } \psi & \xrightarrow{\phi_\delta} & W_{2\delta} \\
 & \searrow \theta_{t \leq t+2\delta} & & & \nearrow
 \end{array}$$

This implies that  $\text{im } \theta_{t \leq t+2\delta} \subseteq \text{im } \phi_\delta \subseteq W_{2\delta}$ , so there are some injections  $j$  and  $i$  which make the following diagram commute as well:

$$\begin{array}{ccccc}
 \text{im } \theta_{t \leq t+2\delta} & \xrightarrow{j} & \text{im } \phi_\delta & \xrightarrow{i} & W_{2\delta} \\
 & \searrow \theta_{t \leq t+2\delta} & & & \nearrow
 \end{array}$$

As all morphisms in the diagram above are injections, we can use the functorial properties of Lemma 3.2.5 having a commutative diagram of the barcodes of each of the previous persistence modules:

$$\begin{array}{ccccc} \text{Bar}(\text{im } \theta_{t \leq t+2\delta}) & \xrightarrow{\mu_{inj}(j)} & \text{Bar}(\text{im } \phi_\delta) & \xrightarrow{\mu_{inj}(i)} & \text{Bar}(W_{2\delta}) \\ & & & \nearrow & \\ & & & \mu_{inj}(\theta_{t \leq t+2\delta}) & \end{array} .$$

We have that

$$\begin{aligned} \text{Bar}(\text{im } \theta_{t \leq t+2\delta}) &= \{(b, d - 2\delta) : (b, d] \in \text{Bar}(W), b < d - 2\delta\}, \\ \text{Bar}(W_{2\delta}) &= \{(b - 2\delta, d - 2\delta) : (b, d] \in \text{Bar}(W)\} \text{ and} \\ \mu_{inj}(\theta_{t \leq t+2\delta})((b, d - 2\delta)) &= (b - 2\delta, d - 2\delta) \end{aligned}$$

Therefore  $\text{im } \mu_{inj}(i) \supseteq \text{im } \mu_{inj}(\psi_{t \leq t+2\delta}) = \text{Bar}(W_{2\delta})_{2\delta}$ . Thus, undoing the shifts, the proof is complete.

3. Let  $(b, d] \in \text{Bar}(\text{im } f_\delta)$  such as for some  $b'$ ,  $\mu_{inj}(b, d] = (b', d] \in \text{Bar}(W)$ . Because of Lemma 3.2.3,  $b' \leq b$ . There are again two cases:

If  $d - b \leq 2\delta$ , then  $b' \geq d - 2\delta \geq b > b - 2\delta$  and  $b' \in [b - 2\delta, b]$ .

Else, if  $d - b > 2\delta$ , there exists an interval  $(b' + 2\delta, d] \in \text{Bar}(\text{im } \theta_{t \leq t+2\delta})$  such that

$$\mu_{inj}(\theta_{t \leq t+2\delta})(b' + 2\delta, d] = \mu_{inj}(i)(b, d] = (a, d].$$

Thus,  $b \leq b' + 2\delta$  and  $b' \in [b - 2\delta, b]$ .

□

At last, we can now prove the other part of the Stability theorem. For so, we will construct a  $\delta$ -matching out of a  $\delta$ -interleaving morphism.

**Proposition 3.2.8** (Theorem 3.0.2, [14]). *Given two persistence modules  $V$ ,  $W$ , with a  $\delta$ -interleaving morphism between them, then there is a  $\delta$ -matching between their barcodes.*

*Proof.* Let  $\mu(\phi) = \mu_{inj} \circ \mu_{sur}$  and let  $\Phi_\delta: \text{Bar}(W_\delta) \rightarrow \text{Bar}(W)$  be the *shift map* that carries each bar  $(a, b]$  into  $(a + \delta, b + \delta]$ . The composition  $\Phi_\delta \circ \mu(\phi)$  is a matching between

$\text{Bar}(V)$  and  $\text{Bar}(W)$ . Hence, using Lemma 3.2.7 and 3.2.6, we get the following diagram:

$$\begin{array}{ccccc}
 \text{Bar}(V) & & \text{Bar}(W)_{\geq 2\delta} & & \text{Bar}(W)_{\geq 2\delta} \\
 \cup \downarrow & & \cap \downarrow & & \cap \downarrow \\
 \text{Bar}(V)_{\geq 2\delta} & \xrightarrow{\mu_{sur}} & \text{Bar}(\text{im } f) & \xrightarrow{\mu_{inj}} & \text{im } \mu_{inj} & \xrightarrow{\Psi_\delta} & \text{Bar } B(W) \\
 \Psi \downarrow & & \Psi \downarrow & & \Psi \downarrow & & \Psi \downarrow \\
 (b, d] & \longmapsto & (b, d'] & \longmapsto & (b', d'] & \longmapsto & (b' + \delta, d' + \delta]
 \end{array}$$

The diagram shows that, by Lemma 3.2.6, a bar  $(b, d] \in \text{Bar}(V)_{\geq 2\delta}$  is sent to  $\mu_{sur}(b, d] = (b, d'] \in \text{Bar}(\text{im } \phi)$  with  $d' \in [d - 2\delta, d]$ . Then, by Lemma 3.2.6, it is sent to  $\mu_{sur}(b, d'] = (b', d']$  with  $b' \in [b - 2\delta, b]$ . At last, using the shift morphism  $\Phi_\delta$  it is carried to  $(b' + \delta, d' + \delta]$ .

This shows that any bar in  $\text{Bar}(V)_{\geq 2\delta}$  is matched. In the same manner it can be seen that any bar in  $\text{Bar}(W)_{\geq 2\delta}$  is matched. Thus, we have that

$$\begin{cases} d - 2\delta \leq d' \leq d \\ b - 2\delta \leq b' \leq b \end{cases} \Rightarrow \begin{cases} d - \delta \leq d' + \delta \leq d + \delta \\ b - \delta \leq b' + \delta \leq b + \delta \end{cases},$$

and therefore,  $\Phi_\delta \circ \mu(\phi)$  is a  $\delta$ -matching between  $\text{Bar}(V)$  and  $\text{Bar}(W)$ .  $\square$

The constructions made by Proposition 3.1.2 and Proposition 3.2.8 assure that given a  $\delta$ -interleaving morphism we can build a  $\delta$ -matching, and conversely, given a  $\delta$ -matching we can build a  $\delta$ -interleaving morphism. This means that if one of the two exists, it fixes a  $\delta$ . Both the interleaving distance and the bottleneck distance try to minimize this  $\delta$ , so once fixed for one of them, the other needs an smaller or equal  $\delta'$ . Thus, with each of the propositions we can prove one of the inequalities needed to reach the isomorphism between the space of persistence diagrams and the space of their barcodes.

**Theorem 3.0.1** (Interleaving Stability, Theorem 2.2.8, [14]). *There is an isometry between a persistence module with the interleaving distance and its barcode with the bottleneck distance. That is, given two persistence modules  $V$  and  $W$ , it holds that*

$$d_{\text{int}}(V, W) = d_{\text{bot}}(\text{Bar}(V), \text{Bar}(W)).$$

*Proof.* Suppose  $d_{\text{int}}(V, W) = \delta$ . Proposition 3.2.8 assures there exist a  $\delta$ -matching between  $\text{Bar}(V)$  and  $\text{Bar}(W)$ . As  $d_{\text{bot}}(V, W)$  is the infimum  $\delta$  for which exists a  $\delta$ -matching,  $d_{\text{bot}}(V, W) \leq d_{\text{int}}(V, W)$ . On the other hand, Proposition 3.1.2 leads, with the same reasoning, to  $d_{\text{int}}(V, W) \leq d_{\text{bot}}(V, W)$ . Thus, it has to be  $d_{\text{int}}(V, W) = d_{\text{bot}}(\text{Bar}(V), \text{Bar}(W))$ .  $\square$

## Chapter 4

# Hausdorff Stability Theorem

Persistence diagrams help summarize the information given by the homology groups of a filtration over a certain data set. They represent the birth and death of every feature in an easy to analice format scattering points over the upper half of the plane  $\mathbb{R}^2$  and its diagonal  $\Delta$ . Once we have computed the diagrams given by two datasets we can measure distances between them using the bottleneck distance, enabling us to decide wether two diagrams are close to each other. However, this would be kind of useless if the bottleneck distance were not stable. If minor differences in original data would cause great changes in the bottleneck distance between the corresponding persistence diagrams, then this data summary method would be as good as any other random method.

Fortunately for us, David Cohen-Steiner, Herbert Edelsbrunner and John Harer, proved in their 2005 paper that the bottleneck distance over persistence diagrams is indeed stable [8]. This means that, when comparing the bottleneck distance between the persistence diagrams formed by the pre images of two tame functions, see Definition 1.1.13, the first will be always lower or equal that the Lebesgue  $L_\infty$  norm between the two functions.

Along this chapter we will follow Edelsbrunner et al. paper [8] to prove Theorem 4.0.2. Recall from Definition 1.1.12 that homological critical values are the real numbers at where homology changes. Recall too the notation given in Definition 1.1.14. For a triangulable topological space  $X$ , let  $f: X \rightarrow \mathbb{R}$  and fix some  $k \in \mathbb{Z}$ , we denoted

$$F_x := H_k(f^{-1}(-\infty, x]),$$

and for every  $x \leq y$ , we denoted the induced inclusion map from  $X_x$  to  $X_y$  as

$$f_x^y: F_x \rightarrow F_y.$$

The persistent homology module associated to  $x$  and  $y$  was the subspace of  $X_y$  given by

$$F_x^y := \text{im } f_x^y.$$

The persistent Betti numbers were defined as

$$\beta_x^y := \dim(F_x^y).$$

We can define the multiplicity of each critical point as follows.

**Definition 4.0.1** (Multiplicity). Let  $f: X \rightarrow \mathbb{R}$  be tame, and  $(a_i)_{i=1,\dots,n}$  be its homological critical values. Take  $(b_i)_{i=1,\dots,n}$  be an interleaved sequence of non critical values such that  $b_{i-1} < a_i < b_i$  for all  $i = 1, \dots, n$ . Define  $b_{-1} = a_0 = -\infty$ ,  $b_{n+1} = a_{n+1} = \infty$ . The **multiplicity** of  $(a_i, a_j) \in D(f)$ , denoted  $\mu_i^j$  is

$$\mu_i^j := \beta_{b_{i-1}}^{b_j} - \beta_{b_i}^{b_j} + \beta_{b_i}^{b_{j-1}} - \beta_{b_{i-1}}^{b_{j-1}}.$$

The **total multiplicity** of a multiset  $A$ , denoted  $\#(A)$  is the sum of the multiplicities of every element in  $A$ .

Note that the total multiplicity of a multiset is the the generalized concept of cardinality of a normal set. While the cardinality of a set counts the number of elements in the set, the multiplicity of a multiset counts how many elements, different or not, are there in the multiset.

When a function  $f$  is tame, we can form a finite multiset of barcodes taking each pair of critical points  $(a_i, a_j)$  with their corresponding multiplicity  $\mu_i^j$ , for each  $0 \leq i < j \leq n+1$ . Hence we can form a persistence diagram  $D(f)$ . The main theorem of this chapter states that the distance of two of those persistence diagrams is never grater that the  $L^\infty$  norm between the functions that form them.

**Theorem 4.0.2** (Main Theorem, [8]). *Let  $X$  be a triangulable space, and  $f, g: X \rightarrow \mathbb{R}$  continuous tame functions. Then,*

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_\infty$$

## 4.1 Hausdorff Stability

Before approaching the proof of Theorem 4.0.2, this section shows how the claim of the theorem is true when the bottleneck distance is replaced by the Hausdorff distance. On Section 4.2 this assertion is used to give an upper limit of the bottleneck distance by Hausdorff distance, proving the main theorem.

We will denote the closed upper left quadrant of a point  $(x, y) \in \mathbb{R}^2$  as

$$Q_x^y := [-\infty, x] \times [y, \infty].$$

**Lemma 4.1.1** ( $k$ -Triangle Lemma, [8]). *Let  $f: X \rightarrow \mathbb{R}$  be a tame function,  $x < y \in \mathbb{R}$  be non critical values of  $f$ . Then the multiplicity  $\mu$  of the persistence diagram of  $f$  in the closed upper left quadrant is*

$$\mu = \#(D(f) \cap Q_x^y) = \beta_x^y.$$

*Proof.* Let  $x = b_i$ ,  $y = b_{j-1}$ .

$$\mu = \sum_{k \leq i \leq j \leq l} \mu_k^l = \sum_{k \leq i \leq j \leq l} \beta_{b_{k-1}}^{b_l} - \beta_{b_k}^{b_l} + \beta_{b_k}^{b_{l-1}} - \beta_{b_{k-1}}^{b_{l-1}} \quad (4.1)$$

$$= \beta_{b_{-1}}^{b_{n+1}} - \beta_{b_i}^{b_{n+1}} + \beta_{b_i}^{b_{j-1}} - \beta_{b_{j-1}}^{b_{j-1}} = \beta_{b_k}^{b_{l-1}} = \beta_x^y. \quad (4.2)$$

The first two equalities in (4.1) are just the definition of total multiplicity. In (4.2), note that every other term in the sum cancels. Then note that  $\beta_{b_{-1}}^{b_{n+1}} = \dim F_{-\infty}^\infty$ ,  $\beta_{b_i}^{b_{n+1}} = \dim F_x^\infty$  and  $\beta_{b_{j-1}}^{b_{j-1}} = \dim F_{-\infty}^y$ . All of them are the dimension of the trivial group, therefore, equal to 0. This leaves only one remaining term and completes the proof.  $\square$

Denote the **upper left quadrants**  $Q := Q_b^c = [-\infty, b] \times [c, \infty]$ ,  $Q_\varepsilon := Q_{b-\varepsilon}^{c+\varepsilon} = [-\infty, b-\varepsilon] \times [c+\varepsilon, \infty]$ .

**Lemma 4.1.2** (Quadrant Lemma, [8]). *Let  $f, g: X \rightarrow \mathbb{R}$  be two tame functions. With the notation above, the following inequality holds,*

$$\#(D(f), \cap Q_\varepsilon) \leq \#(D(g) \cap Q).$$

*Proof.* Let  $\varepsilon := \|f - g\|_\infty$ . Hence, considering the pre-image of the functions, we have the following inclusions

$$f^{-1}((-\infty, x]) \subseteq g^{-1}((-\infty, x + \varepsilon)), \quad (4.3)$$

$$g^{-1}((-\infty, x]) \subseteq f^{-1}((-\infty, x + \varepsilon)). \quad (4.4)$$

Name  $\varphi_x: F_x \rightarrow G_{x+\varepsilon}$  to the inclusion map induced by (4.3) and  $\psi_x: G_x \rightarrow F_{x+\varepsilon}$  to the inclusion map induced by (4.4). Let  $b < c \in \mathbb{R}$ . With the described maps, we can form commutative diagram (4.5) where we observe that

$$\text{im}(f_{c-\varepsilon}^{c+\varepsilon}) = F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c \circ g_b^c(G_b) = \psi_c(G_b^c).$$

$$\begin{array}{ccc} F_{b-\varepsilon} & \xrightarrow{f_{b-\varepsilon}^{c+\varepsilon}} & F_{c+\varepsilon} \\ \varphi_{b-\varepsilon} \downarrow & & \uparrow \psi_c \\ G_b & \xrightarrow{g_b^c} & G_c \end{array} \quad (4.5)$$

Last inclusion is enough for the requirements of this proof, never the less, we will make one more note that will be useful latter on, through the proof of Lemma 4.1.3. Fit the maps so that they describe commutative diagram (4.6), showing that

$$\psi_c(G_b^c) = \psi_c \circ g_b^c(G_b) = f_{b+\varepsilon}^{c+\varepsilon} \circ \psi_b(G_b) \subseteq F_{b+\varepsilon}^{c+\varepsilon}.$$

$$\begin{array}{ccc}
F_{b+\varepsilon} & \xrightarrow{f_{b+\varepsilon}^{c+\varepsilon}} & F_{c+\varepsilon} \\
\varphi_b \uparrow & & \uparrow \psi_c \\
G_b & \xrightarrow{g_b^c} & G_c
\end{array} \tag{4.6}$$

From both diagrams we finally obtain the inclusion chain

$$F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c(G_b^c) \subseteq F_{b+\varepsilon}^{c+\varepsilon}. \tag{4.7}$$

By Lemma 4.1.1, we are able to count the elements in the intersection of the diagrams with the upper left quadrants. Hence

$$\begin{aligned}
\#(D(f) \cap Q_\varepsilon) &= \beta_{b-\varepsilon}^{c+\varepsilon} = \dim F_{b-\varepsilon}^{c+\varepsilon}, \\
\#(D(g) \cap Q) &= \beta_b^c = \dim G_b^c.
\end{aligned}$$

As if one homology group is contain in an other, the dimension of the first must be lower or equal to the one of the second. Also, the dimension is invariant under inclusion maps. Thus, the first inclusion of (4.7) asserts that  $F_{c-\varepsilon}^{c+\varepsilon} \subseteq \psi_c(G_b^c)$  and therefore we have proven that  $\dim F_{c-\varepsilon}^{c+\varepsilon} \leq \dim G_b^c$ .  $\square$

Before next lemma we will introduce some new notation. Let  $f: X \rightarrow \mathbb{R}$  be a tame function. Let  $w < x < y < z \in \mathbb{R}$  be numbers different from critical values of  $f$ . Recall that  $F_x = H_k(f^{-1}(-\infty, x])$ ,  $f_x^y: F_x \rightarrow F_y$  and  $F_x^y = \dim f_x^y$ . We denote

$$f_x^{y,z} := f_y^z|_{F_x^y}, \quad F_x^{y,z} := \dim f_x^{y,z}.$$

Note, from linear algebra, that  $\dim F_x^{y,z} = \dim F_x^y - \dim F_x^z$ . Also note that  $F_w^y \subseteq F_x^y$ . Therefore,  $\ker F_w^y \subseteq \ker F_x^y$  and we can define the quotient

$$F_{w,x}^{y,z} := F_x^{y,z} / F_w^{y,z}.$$

Let  $a < b < c < d \in \mathbb{R}$ . Denote the **rectangles**  $R := [a, b] \times [c, d]$ ,  $R_\varepsilon := [a + \varepsilon, b - \varepsilon] \times [c + \varepsilon, d - \varepsilon]$ .


**Lemma 4.1.3** (Box Lemma, [8]). *With the notation above, the following inequality holds,*

$$\#(D(f) \cap R_\varepsilon) \leq \#(D(g) \cap R).$$

*Proof.* Note that we can asume that  $a + \varepsilon < b - \varepsilon$  and  $c + \varepsilon < d - \varepsilon$ . Otherwise the rectangle  $R_\varepsilon$  would not exist. Also note that

$$\begin{aligned}
\#(D(f) \cap R_\varepsilon) &= \dim F_{a+\varepsilon, b-\varepsilon}^{c+\varepsilon, d-\varepsilon} \\
\#(D(g) \cap R) &= \dim G_{a,b}^{c,d}
\end{aligned}$$

To make our proof we draw diagram 4.1. Lets analice every element of the diagram.



First of all, the middle upside arrows, are

$$u_2 = f_{a+\varepsilon}^{c+\varepsilon, d-\varepsilon}, \quad u_3 = f_{b-\varepsilon}^{c+\varepsilon, d-\varepsilon}.$$

Right arrows  $r_1, r_2, r_3, r_4$  represent the inclusions from its respective vector spaces to their destination. The objective is to define the respective quotients to define  $G_{a,b}^{c,d}$ . Recall the inclusion maps defined in the proof of Lemma 4.1.2,  $\varphi_x: F_x \rightarrow G_{x+\varepsilon}$  and  $\psi_x: G_x \rightarrow F_{x+\varepsilon}$ . We define

$$E_b^c := \psi_c^{-1}(F_{b-\varepsilon}^{c+\varepsilon, d-\varepsilon}) \cap G_b^c, \quad E_a^c := G_a^c \cap E_b^c.$$

We have then that the outer upside arrows are the respective restrictions

$$u_1 = g_a^{c,d}|_{E_a^c}, \quad u_4 = g_b^{c,d}|_{E_b^c}.$$

We denote

$$s_1 := \varphi_{d-\varepsilon}|_{F_{b-\varepsilon}^{d-\varepsilon}}, \quad s_2 := \psi_c|_{E_a^c}, \quad s_3 := \psi_c(G_b^c)|_{E_b^c}.$$

By the inclusions (4.7), we have that

$$\varphi_{d-\varepsilon}(F_{b-\varepsilon}^{d-\varepsilon}) \subseteq G_b^d, \quad \psi_c(G_a^c) \subseteq F_{a+\varepsilon}^{c+\varepsilon}, \quad F_{b-\varepsilon}^{c+\varepsilon} \subseteq \psi_c(G_b^c).$$

Note that by the manner we have define the diagram, it need to be that

$$\text{im}(s_3) = \ker(u_3), \quad \text{im}(s_1) \subseteq G_b^d.$$

Also, we can observe that  $u_4 = s_1 \circ u_3 \circ s_3$ . As  $u_3 \circ s_3 = 0$ , then  $E_b^c = \ker(u_4)$ . Also, as  $r_1 \circ u_1 = u_4 \circ r_4 = 0$  and  $r_1$  is the inclusion, then  $E_a^c = \ker(u_1)$ . Hence, we can write

$$E_b^c = E_b^{c,d} \subseteq G_b^{c,d}, \quad E_a^c = E_a^{c,d} \subseteq G_a^{c,d}.$$



As  $E_a^{c,d} = E_b^{c,d} \cap G_a^{c,d}$ , the following quotient inclusion holds

$$E_{a,b}^{c,d} = E_b^{c,d} / E_a^{c,d} \subseteq G_b^{c,d} / G_a^{c,d} = G_{a,b}^{c,d}.$$

Therefore, we have

$$\dim(E_{a,b}^{c,d}) \leq \dim(G_{a,b}^{c,d}).$$

Now note that

$$E_{a,b}^{c,d} = \ker(u_4) / \ker(u_1), \quad F_{a+\varepsilon, b+\varepsilon}^{c+\varepsilon, d+\varepsilon} = \ker(u_3) / \ker(u_2).$$

By construction  $s_3(\ker(u_4)) = \ker(u_3)$ . As for every  $x \in \ker(u_1)$ ,  $r_2 \circ u_2 \circ s_2(x) = u_3 \circ s_3 \circ r_4(x) = 0$ , and  $r_2$  is an injection, then  $s_3(\ker(u_1)) = s_2(\ker(u_1)) \subseteq \ker(u_2)$ , we get

$$\dim(F_{a+\varepsilon, b+\varepsilon}^{c+\varepsilon, d+\varepsilon}) \leq \dim(E_{a,b}^{c,d}).$$

Hence, the desired inequality is hold as we have seen that

$$\#(D(f) \cap R_\varepsilon) = \dim F_{a+\varepsilon, b-\varepsilon}^{c+\varepsilon, d-\varepsilon} \leq \dim(E_{a,b}^{c,d}) \leq \dim(G_{a,b}^{c,d}) = \#(D(g) \cap R).$$

□

**Theorem 4.1.4** (Hausdorff Stability).

$$d_H(D(f), D(g)) \leq \|f - g\|_\infty.$$

*Proof.* As a direct consequence of Lemma 4.1.3 if  $(x, y) \in D(f)$  then there must exist some point at  $D(g)$  at distance less than or equal to  $\varepsilon = \|f - g\|_\infty$  from  $(x, y)$  since the total multiplicity of  $D(g) \cap R_\varepsilon$  is at least one. □

## 4.2 Bottleneck Stability

In this section we are going to prove Theorem 4.0.2.

**Definition 4.2.1** (Very close tame functions). Let  $f, g: X \rightarrow \mathbb{R}$  be tame functions. We define

$$\delta_f = \min\{\|p - q\|_\infty : p \in D(f) \setminus \Delta, q \in D(f), p \neq q\}.$$

We say that  $g$  is **very close** to  $f$  if  $\|f - g\|_\infty < \delta_f/2$ .

**Lemma 4.2.2** (Easy Bijection Lemma, [8]). *Let  $f, g: X \rightarrow \mathbb{R}$  be tame functions, where  $g$  is very close to  $f$ . Then, following holds,*

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_{\infty}.$$

*Proof.* Let  $p := (a_i, a_j) \in D(f) - \Delta$  be a point in the diagram of  $f$  that is not in the diagonal, and let  $\mu := \beta_i^j$  denote its multiplicity. Let  $S_{\varepsilon}$  be the square of center  $p$  and radius  $\varepsilon = \|f - g\|_{\infty}$ . That is, the square of side  $2\varepsilon$ . By definition of the square  $S_{\varepsilon}$  we have that the number of points of its intersection with the diagram of  $g$  must be greater or equal than the multiplicity at  $p$ . Hence, by the Box Lemma 4.1.3 we have

$$\mu \leq \#(D(g) \cap S_{\varepsilon}) \leq \#(D(f) \cap S_{2\varepsilon}).$$

As  $g$  is very close to  $f$ , we have  $2\varepsilon \leq \delta_f$ . Hence  $p$  is the only point of  $D(f)$  in  $S_{\varepsilon}$ , and therefore the previous inequality is in fact an equivalence. If there was a point in the intersection which was not in  $\mu$  then it would be inside the square  $S_{\varepsilon}$  and meaning the distance  $\|f - g\|$  would be smaller. That is

$$\mu = \#(D(g) \cap S_{\varepsilon}).$$

Hence we can map every point in  $D(g) \cap S_{\varepsilon}$  with  $p$ . We can then repeat this process for every other  $p \in D(f) \setminus \Delta$ . After this process, every point of  $D(g)$  which have not been matched yet, must be at distance greater than  $\varepsilon$  from  $D(f) \setminus \Delta$ . By Theorem 4.1.4, every unmatched point must be at distance at most  $\varepsilon$  from the diagonal  $\Delta$ . Hence, if we map each of this points to  $\Delta$ , we have built a bijection between  $D(f)$  and  $D(g)$  that moves each point at most  $\varepsilon$ .  $\square$

**Definition 4.2.3.** Let  $\hat{f}, \hat{g}$  be two piecewise linear functions over a simplicial complex  $K$ . Let  $\lambda \in [0, 1]$ . A **convex combination** of  $\hat{f}$  and  $\hat{g}$  is a function of the form

$$h_{\lambda} := (1 - \lambda)\hat{f} + \lambda\hat{g}.$$

**Lemma 4.2.4** (Interpolation Lemma, [8]). *Let  $K$  be a simplicial complex. Take two piecewise linear functions  $\hat{f}, \hat{g}: K \rightarrow \mathbb{R}$ . Then, the following holds,*

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \|\hat{f} - \hat{g}\|_{\infty}.$$

*Proof.* Let  $c := \|\hat{f} - \hat{g}\|_{\infty}$ . For every  $\lambda \in [0, 1]$ , define  $\delta(\lambda) := \delta_{h_{\lambda}} > 0$ . Let  $J_{\lambda}$  denote open intervals around each  $\delta$  as follows, and consider the set  $C$  of all  $J_{\lambda}$  be

$$C := \left\{ J_{\lambda} := \left( \lambda - \frac{\delta(\lambda)}{4c}, \lambda + \frac{\delta(\lambda)}{4c} \right) \right\}.$$

The set  $C$  is an open cover of the interval  $[0, 1]$ . Let  $C'$  be the minimal subcover of  $C$ . As  $[0, 1]$  is compact, the subcover  $C'$  must be finite. Consider then  $\lambda_1 < \lambda_2 < \dots < \lambda_n$  the midpoints of the intervals in  $C'$ . As  $C'$  is minimal, then each intersection  $J_{\lambda_i} \cap J_{\lambda_{i+1}}$  is not empty. Hence

$$\lambda_i + \lambda_{i+1} \leq \frac{\delta(\lambda_i) + \delta(\lambda_{i+1})}{4c} \leq \frac{\max\{\delta(\lambda_i), \delta(\lambda_{i+1})\}}{2c}.$$

Therefore, by definition of  $c$  and each  $h_{\delta_i}$ , it holds

$$\|h_{\delta_i} - h_{\delta_{i+1}}\|_\infty = c(\lambda_{i+1} - \lambda_i) \leq \frac{\max\{\delta(\lambda_i), \delta(\lambda_{i+1})\}}{2}.$$

This implies that  $h_{\delta_i}$  is very close to  $h_{\delta_{i+1}}$  or viceversa. Then, by Lemma 4.2.2, for every  $1 \leq i \leq n-1$ ,

$$d_{\text{bot}}(D(h_{\lambda_i}), D(h_{\lambda_{i+1}})) \leq \|h_{\lambda_i} - h_{\lambda_{i+1}}\|_\infty. \quad (4.8)$$

Let  $\lambda_0 = 0$  and  $\lambda_{n+1} = 1$ . Then  $h_{\lambda_0} = \hat{f}$  is very close to  $h_{\lambda_1}$  and  $h_{\lambda_1} = \hat{g}$  is very close to  $h_{\lambda_n}$  and therefore (4.8) also holds for  $i = 0$  and  $i = n+1$ . Finally, using the triangle inequality we have

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \sum_{i=0}^n d_{\text{bot}}(D(h_{\lambda_i}), D(h_{\lambda_{i+1}})) \leq \sum_{i=0}^n \|h_{\lambda_i} - h_{\lambda_{i+1}}\|_\infty = \|\hat{f} - \hat{g}\|_\infty.$$

□

For the final proof lets recall what the star of a simplicial complex is.

**Definition 4.2.5** (Star of a simplicial complex). Let  $\sigma$  be a simplex in a simplicial complex  $L$ . The **star**  $\text{St}(\sigma)$  of  $\sigma$  is the set of simplices in  $L$  which contain  $\sigma$  as a face. The **star of a subset**  $K$  of  $L$ , denoted  $\text{St}(K)$ , is the union of the stars of each simplex of  $K$ .

**Theorem 4.0.2** (Main Theorem, [8]). *Let  $X$  be a triangulable space, and  $f, g: X \rightarrow \mathbb{R}$  continuous tame functions. Then,*

$$d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_\infty$$

*Proof.* As  $X$  is triangulable, there exists a finite simplicial complex  $L$  and a homeomorphism  $\Phi: L \rightarrow X$ . Hence a persistence diagram is invariant under this change of variables. That is,  $f \circ \Phi: L \rightarrow \mathbb{R}$  is tame and  $D(f \circ \Phi) = D(f)$ . Since  $f$  and  $g$  are continuous and

$L$  is compact, for every  $\delta > 0$  there exists a subdivision  $K$  of  $L$  such that for every  $u, v$  points of a common simplex  $\sigma \in K$ ,

$$\begin{aligned} |f \circ \Phi(u) - f \circ \Phi(v)| &\leq \delta, \\ |g \circ \Phi(u) - g \circ \Phi(v)| &\leq \delta. \end{aligned}$$

Let  $\hat{f}, \hat{g}: \text{St}(K) \rightarrow \mathbb{R}$  be the piecewise linear interpolations of  $f \circ \Phi$  and  $g \circ \Phi$  on  $K$ . By construction of  $K$  and the definition of the  $L_\infty$ -norm, these interpolations satisfy

$$\begin{aligned} \|\hat{f} - f \circ \Phi\|_\infty &\leq \delta, \\ \|\hat{g} - g \circ \Phi\|_\infty &\leq \delta. \end{aligned}$$

Hence, by Lemma 4.2.4 and the triangle inequality

$$d_{\text{bot}}(D(\hat{f}), D(\hat{g})) \leq \|\hat{f} - \hat{g}\|_\infty \leq \|f \circ \Phi - g \circ \Phi\|_\infty + 2\delta \leq \|f - g\|_\infty + 2\delta.$$

Now, we can take some  $\delta$  such that  $\delta \leq \delta_f/2$  so  $\hat{f}$  is very close to  $f$ . This allows us to use Lemma 4.2.2 to make a bijection that satisfies

$$d_{\text{bot}}(D(f), D(\hat{f})) \leq d_{\text{bot}}(D(f \circ \Phi), D(\hat{f})) \leq \delta.$$

Analogously, also assuring  $\delta < \delta_g$  we also have

$$d_{\text{bot}}(D(g), D(\hat{g})) \leq d_{\text{bot}}(D(g \circ \Phi), D(\hat{g})) \leq \delta,$$

and therefore, by triangle inequality again,

$$d_{\text{bot}}(D(f), D(g)) \leq d_{\text{bot}}(D(f), D(\hat{f})) + d_{\text{bot}}(D(\hat{f}), D(\hat{g})) + d_{\text{bot}}(D(\hat{g}), D(g)) \leq 4\delta.$$

As this holds for any  $\delta$  smaller than  $\delta_f$  and  $\delta_g$ , taking the limit when  $\delta$  tends to 0, we complete the proof.  $\square$

## Chapter 5

# Gromov-Hausdorff's Stability Theorem

Chapter 4 established the stability of persistence diagrams under small-amplitude perturbations of filtering functions. Specifically, for two tame functions defined on the same topological space, the bottleneck distance between their persistence diagrams is bounded by the uniform norm of the function difference.

In this chapter, we extend this stability framework to distinct metric spaces, enabling quantitative comparisons of real-world datasets (e.g., point clouds representing different shapes). We achieve this through two classic filtrations from computational geometry, the Vietoris-Rips and the Čech filtrations, to obtain the corresponding modules from finite metric spaces and then prove how the bottleneck distance between their persistence diagrams is stable respect the Gromov-Hausdorff (see Definition 1.5.3) distance between the metric spaces. This bound facilitates efficient clustering and differentiation of metric spaces with distinct topological features (e.g., non-isometric shapes), as validated by experiments on 3D models.

Section 5.1 details Vietoris-Rips and the Čech filtrations and sets the necessary lemmas to prove the stability Theorem 5.1.8. Section 5.2 generalizes the result comparing the distance between filtrations given by some functions with the generalized Gromov-Hausdorff distance from Definition 1.5.12.

Along all the chapter we follow the 2009 paper from Frédéric Chazal and David Cohen-Steiner, [5].

## 5.1 Gromov-Hausdorff stability

**Definition 5.1.1** (Vietoris-Rips filtration). Let  $(X, d)$  be a finite metric space and let  $\alpha > 0$ . The **Vietoris-Rips complex** associated with  $X$  of radius  $\alpha$ ,  $\mathcal{R}_\alpha(X, d)$ , is the simplicial complex whose 0-simplices are the elements of  $X$  and, for  $k \geq 1$ , its  $k$ -dimensional simplices are formed by every subset  $\{x_0, x_1, \dots, x_k\} \subseteq X$  such that  $d(x_i, x_j) \leq \alpha$  for every  $i, j = 1, \dots, k$ .

The family  $\mathcal{R}(X, d) := \{\mathcal{R}_\alpha(X, d)\}_{\alpha>0}$  is named **Vietoris-Rips filtration**.

Given a real function  $f: X \rightarrow \mathbb{R}$ , let  $X_\alpha := f^{-1}((-\infty, \alpha]) \subseteq X$  be the **pre-image of  $f$  delimited by  $\alpha$** . We define the **Vietoris-Rips filtration associated with  $f$**  as  $\mathcal{R}(X, d, f) := \{\mathcal{R}_\alpha(X_\alpha, d)\}_{\alpha>0}$ .

**Definition 5.1.2** (Čech filtration). Let  $(X, d)$  be a finite metric space and let  $\alpha > 0$ . The **Čech complex** associated with  $X$  of radius  $\alpha$ ,  $\check{\mathcal{C}}_\alpha(X, d)$ , is the simplicial complex whose 0-simplices are the elements of  $X$  and, for  $k \geq 1$ , its  $k$ -dimensional simplices are formed by every subset  $\{x_0, x_1, \dots, x_k\} \subseteq X$  such that there exists some  $x \in X$  such that  $d(x, x_i) \leq \alpha$  for all  $i = 1, \dots, k$ .

The family  $\check{\mathcal{C}}(X, d) := \{\check{\mathcal{C}}_\alpha(X, d)\}_{\alpha>0}$  is named **Čech filtration**.

Given a real function  $f: X \rightarrow \mathbb{R}$ , let  $X_\alpha := f^{-1}((-\infty, \alpha]) \subseteq X$ . We define the **Čech filtration associated with  $f$**  as  $\check{\mathcal{C}}(X, d, f) := \{\check{\mathcal{C}}_\alpha(X_\alpha, d)\}_{\alpha>0}$ .

To prove Theorem 5.1.8 we first give tree auxiliary lemmas that we will need. The first, just shows how to embed a finite metric space into the Euclidean space with the uniform norm.

**Lemma 5.1.3** (Exercise 3.5.4, [4]). *Any finite metric space of cardinality  $n$  can be isometrically embedded into  $(\mathbb{R}^n, \ell^\infty)$ .*

*Proof.* Let  $X$  be a compact metric space and let  $C(X)$  be the space of all continuous functions from  $X$  to  $\mathbb{R}$ . Let  $f, g \in C(X)$ . Recall the uniform distance given by

$$d_\infty(f, g) := \sup |f(x) - g(x)|.$$

First, we will check that the pair  $(C(X), d_\infty)$  is a metric space. Naturally nonnegativity holds and

$$d_\infty(f, f) = \sup |f(x) - f(x)| = 0.$$

Commutativity also holds as

$$d_\infty(f, g) = \sup |f(x) - g(x)| = \sup |g(x) - f(x)| = d_\infty(g, f).$$

Finally, if  $h \in C(X)$ , triangle inequality holds because

$$\begin{aligned} d_\infty(f, h) &= \sup |f(x) - h(x)| = \sup |f(x) + g(x) - g(x) - h(x)| \\ &\leq \sup |f(x) - g(x)| + \sup |g(x) - h(x)| = d_\infty(f, g) + d_\infty(g, h). \end{aligned}$$

Now we are going to verify that the map  $E: X \rightarrow C(X)$  defined by  $E(x) = d(x, \cdot)$  is an isometric embedding onto its image. Note that

$$d_\infty(d(x, \cdot), d(y, \cdot)) = \sup_z |d(x, z) - d(y, z)| \leq \sup_z |d(x, y)| = d(x, y).$$

On the other hand if we take  $z = y$  we then have

$$|d(x, y) - d(y, y)| = d(x, y),$$

and therefore

$$\sup_z |d(x, z) - d(y, z)| \geq d(x, y).$$

The proof of the lemma is just an analogous case taking  $C_n(X)$  as the set of continuous functions  $f: X \rightarrow \mathbb{R}^n$ , and, for every  $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$ ,  $\ell^\infty(x, y) = \max_i |x_i - y_i|$ .  $\square$

**Lemma 5.1.4** (Lemma VII, [9]). *Let  $X \subset \mathbb{R}^n$  and  $\alpha > 0$ . Then the  $\alpha$ -Čech and the  $\alpha$ -Rips complexes coincide when using the  $\ell^\infty$ -norm. That is*

$$\check{C}_\alpha(X, \ell^\infty) = R_\alpha(X, \ell^\infty).$$

*Proof.* ...  $\square$

**Definition 5.1.5** (Paracompact space). Let  $X$  be a topological space. It is said to be **paracompact** if for all covering  $\mathcal{U}$  of  $X$ , there exists  $\mathcal{V} \subseteq \mathcal{U}$  such that  $\mathcal{V}$  is a finite covering.

**Definition 5.1.6** (Good cover). Let  $S$  be a topological space and  $I$  a set of indexes. A **good cover** of  $S$  is a family  $\mathcal{U} = U_i$  of open subsets covering  $S$  such that for every finite subset  $J \subset I$ , the common intersection

$$\bigcap_{j \in J} U_j$$

is either empty or contractible.

**Lemma 5.1.7** ([6]). *Let  $S \subset S'$  be two paracompact spaces. Let  $\mathcal{U} = \{U_x\}_{x \in A}$ ,  $\mathcal{U}' = \{U'_x\}_{x \in A'}$  be two good covers of  $S$  and  $S'$  respectively, based on finite parameter sets  $A \subset A'$  such that  $U_x \subset U'_x$  for all  $x \in A$ . Then the homotopy equivalences  $\mathcal{N}\mathcal{U} \rightarrow S$  and  $\mathcal{N}\mathcal{U}' \rightarrow S'$  commute with the canonical inclusions  $S \rightarrow S'$  and  $\mathcal{N}\mathcal{U} \rightarrow \mathcal{N}\mathcal{U}'$  at homology level.*

**Theorem 5.1.8** (Theorem 3.1, [5]). *Let  $(X, d_X)$ ,  $(Y, d_Y)$  be finite metric spaces. Then, for any  $k \in \mathbb{N}$ ,*

$$d_{\text{bot}}(D_k(\mathcal{R}(X, d_X)), D_k(\mathcal{R}(Y, d_Y))) \leq d_{\text{GH}}((X, d_X), (Y, d_Y)).$$

*Proof.* Let  $\varepsilon = d_{\text{GH}}((X, d_X), (Y, d_Y))$ . As  $X$  and  $Y$  are finite, they are compact, and therefore the infimum when computing Gromov-Hausdorff distance using Definition 1.5.3 is in fact a minimum. That is, there exists a metric space  $(Z, d_Z)$  and two isometric embeddings  $\gamma_X: X \rightarrow Z$  and  $\gamma_Y: Y \rightarrow Z$  such that

$$d_{\text{H}}^Z(\gamma_X(X), \gamma_Y(Y)) = \varepsilon,$$

where  $d_{\text{H}}^Z$  denotes the Hausdorff distance respect the distance  $d_Z$ . Consider the subspace  $\gamma_X(X) \cup \gamma_Y(Y) \subseteq Z$  with the induced metric from  $Z$ . As both  $X$  and  $Y$  are finite, let

$$n := \#(X) + \#(Y).$$

Hence, by Lemma 5.1.3, there exists an isometric embedding

$$\gamma: (\gamma_X(X) \cup \gamma_Y(Y), d_Z) \rightarrow (\mathbb{R}^n, \ell^\infty).$$

Let  $d_{\text{H}}^\infty$  denote the Hausdorff distance respect the distance  $d_\infty$ . We then have

$$d_{\text{H}}^\infty(\gamma \circ \gamma_X(X), \gamma \circ \gamma_Y(Y)) = d_{\text{H}}^Z(\gamma_X(X), \gamma_Y(Y)) = \varepsilon.$$

Let  $\delta_X$  be the distance function from a point in  $\mathbb{R}^n$  to  $X$ , and analogously, let  $\delta_Y$  be the distance function to  $Y$ . In  $\ell^\infty$  norm, by how we defined  $\varepsilon$ , we have

$$\|\delta_X - \delta_Y\|_\infty = \max_{i=1, \dots, n} |\delta_{x_i} - \delta_{y_i}| \leq \varepsilon.$$

As distance functions are linear, both  $\delta_X$  and  $\delta_Y$  are lower envelopes of piecewise-linear functions and therefore they are piecewise-linear too. Hence, both  $\delta_X$  and  $\delta_Y$  are tame and continuous so by Theorem 4.0.2 we have

$$d_{\text{bot}}(D(\delta_X), D(\delta_Y)) \leq \|\delta_X - \delta_Y\|_\infty \leq \varepsilon.$$

Let  $\alpha \in \mathbb{R}$ . Define an off-set of radius  $\alpha$  around the image of the embedding of  $X$  into  $\mathbb{R}^n$  as

$$\gamma \circ \gamma_X(X)^\alpha := \bigcup_{x \in \gamma \circ \gamma_X(X)} B_\alpha^{\ell^\infty}(x),$$

where  $B_\alpha^{\ell^\infty}(x)$  denotes the ball of radius  $\alpha$  and center  $x$  using distance  $d_\infty$ . As balls in  $\ell^\infty$  are hypercubes, they are convex, and therefore their intersection is either empty or



contractible. By Lemma 5.1.7 know that  $\delta_X$  has the same persistence diagram as the Čech complex  $\check{\mathcal{C}}(\gamma \circ \gamma_X, \ell^\infty)$ . By Lemma 5.1.4, when using the  $\ell^\infty$ -norm, Čech and Rips complexes coincide and so do their filtrations. As  $\gamma \circ \gamma_X$  is an isometric embedding, we then have

$$\check{\mathcal{C}}(\gamma \circ \gamma_X, \ell^\infty) = \mathcal{R}(\gamma \circ \gamma_X, \ell^\infty) = \mathcal{R}(X, d_X).$$

Hence, the persistence diagram of  $\mathcal{R}(X, \ell^\infty)$  is the same as the persistence diagram of  $\gamma_X$ . The same is true taking  $Y$  and therefore we have

$$d_{\text{bot}}(D(\mathcal{R}(X, d_X)), D(\mathcal{R}(Y, d_Y))) = d_{\text{bot}}(D(\gamma_X), D(\gamma_Y)) \leq \varepsilon.$$

□

**Proposition 5.1.9.** *Let  $(X, d_X)$ ,  $(Y, d_Y)$  be finite metric spaces. Then, for any  $k \in \mathbb{N}$ , the bottleneck distance*

$$d_{\text{bot}}(D_k(\mathcal{R}(X, d_X)), D_k(\mathcal{R}(Y, d_Y))),$$

*is a tight lower bound of*

$$d_{\text{GH}}((X, d_X), (Y, d_Y)).$$

*That is, it is the largest possible lower bound.*

*Proof.* Its enough to find an example were both distances are equal. For so, take the two point spaces  $X = \{a, b\}$  with distance  $d_X(a, b) = 2$ , and  $Y = \{c, d\}$  with  $d_Y(c, d) = 2 + 2\varepsilon$ . Both spaces can be isometrically mapped into the real line  $\mathbb{R}$ , with  $X$  mapped to  $\{0, 2\}$  and  $Y$  mapped to  $\{-\varepsilon, 2 + \varepsilon\}$ . Hence  $d_{\text{GH}}(X, Y) \leq \varepsilon$ .

On the other hand, the 0-dimensional persistence diagram of the Rips filtration of  $(X, d_X)$  and  $(Y, d_Y)$  are

$$\begin{aligned} D_0(\mathcal{R}(X, d_X)) &= \{(0, \infty), (0, 1)\}, \\ D_0(\mathcal{R}(Y, d_Y)) &= \{(0, \infty), (0, 1 - \varepsilon)\}, \end{aligned}$$

and therefore,  $d_{\text{bot}}(D_k(\mathcal{R}(X, d_X)), D_k(\mathcal{R}(Y, d_Y))) = \varepsilon$ .

□

## 5.2 A generalization with endowed functions

The following theorem generalizes Theorem 5.1.8.

**Theorem 5.2.1** (Theorem 3.2, [5]). *Let  $(X, d_X)$ ,  $(Y, d_Y)$  be finite metric spaces endowed with the functions  $f: X \rightarrow \mathbb{R}$  and  $g: Y \rightarrow \mathbb{R}$ . Then*

$$d_{\text{bot}}(D_k(\mathcal{R}(X, d_X, f)), D_k(\mathcal{R}(Y, d_Y, g))) \leq d_{\text{GH}}^1((X, d_X, f), (Y, d_Y, g)).$$

*Proof.* We follow a similar procedure to the proof of Theorem 5.1.8. Start setting

$$\varepsilon := d_{\text{GH}}^1((X, d_X, f), (Y, d_Y, g)).$$

For every  $\alpha \in \mathbb{R}$ , recall the notation for the pre-images of  $f$  and  $g$  by  $\alpha$ ,

$$\begin{aligned} X_\alpha &:= f^{-1}((-\infty, \alpha]) \subseteq X, \\ Y_\alpha &:= g^{-1}((-\infty, \alpha]) \subseteq Y. \end{aligned}$$

As before, as  $X$  and  $Y$  are finite, the infimum in  $d_{\text{GH}}^1$  of Definition 1.5.12 is actually a minimum realized by some correspondance  $R \in (X \times Y)$ . Also, the disjoint union  $Z = X \cup Y$ , can be endowed with a metric  $d_Z$  and a pair of inclusions  $\gamma_X: X \rightarrow Z$ ,  $\gamma_Y: Y \rightarrow Z$  such that for every  $(x, y) \in R$ ,

$$\begin{aligned} d_Z(\gamma_X(X), \gamma_Y(Y)) &\leq \frac{1}{2} \text{dis}(R) \leq \varepsilon, \text{ and} \\ |f(x) - g(y)| &\leq \|f - g\|_{\ell^\infty} \leq \varepsilon. \end{aligned}$$

By Lemma 5.1.3,  $(\gamma_X(X) \cup \gamma_Y(Y), d_Z)$  can be isometrically embedded by some  $\gamma$  into  $(\mathbb{R}^n, \ell^\infty)$ , where

$$n := \#(X) + \#(Y).$$

Hence, for every  $(x, y) \in R$  we have

$$\|\gamma \circ \gamma_X(X) - \gamma \circ \gamma_Y(Y)\|_{\ell^\infty}.$$

Note that the filtrations given by the off-set defined in the proof of Theorem 5.1.8, can be seen as persistence modules  $V := \{\gamma \circ \gamma_X(X_\alpha)^\alpha\}_{\alpha > 0}$  and  $W := \{\gamma \circ \gamma_Y(Y_\alpha)^\alpha\}_{\alpha > 0}$  are  $\varepsilon$ -interleaved. That is, for all  $\alpha > 0$ ,

$$\gamma \circ \gamma_X(X_\alpha)^\alpha \subseteq \gamma \circ \gamma_Y(Y_\alpha)^{\alpha+\varepsilon} \subseteq \gamma \circ \gamma_X(X_\alpha)^{\alpha+2\varepsilon}.$$

This is because for every element  $p \in \gamma \circ \gamma_X(X_\alpha)^\alpha$ , there exists some  $x \in X$  such that

$$\|p - \gamma \circ \gamma_X(x)\|_{\ell^\infty} \leq \alpha.$$

Hence, taking some  $y \in Y$  such that  $(x, y) \in R$  we have that

$$\|\gamma \circ \gamma_X(x) - \gamma \circ \gamma_Y(y)\|_{\ell^\infty} \leq \varepsilon,$$

and as

$$g(y) \leq f(x) + \varepsilon \leq \alpha + \varepsilon,$$

we have that  $y \in Y_{\alpha+\varepsilon}$  and therefore

$$\|p - \gamma \circ \gamma_Y(y)\|_{\ell^\infty} \leq \alpha + \varepsilon,$$

and so  $p \in \gamma \circ \gamma_Y(Y_\alpha)^{\alpha+\varepsilon}$ . The second inclusion follows analogously.

Note that as we have two  $\varepsilon$ -interleaved modules we can use 3.0.1 to express the bottleneck distance as the interleaving distance. Thus

$$d_{\text{bot}}(D_k(V), D_k(W)) = d_{\text{int}}(V, W) \leq \epsilon.$$

Analogously to the previous proof, Lemma 5.1.7 tells this inequality is also valid taking the Čech filtrations, and Lemma 5.1.4 let us take the Rips filtrations, completing the proof.  $\square$

# Chapter 6

## Vectorizations' Stability Theorems

The study of stability in topological data analysis is crucial for ensuring that small perturbations in the input data lead to controlled changes in its topological representations. This chapter explores various vectorization methods—persistence landscapes, persistence images, and Euler curves—and their associated stability theorems. By establishing bounds on how these vectorizations change with respect to distances between persistence modules or diagrams, we gain a deeper understanding of their robustness in practical applications.

Section 6.1 reviews the results of the 2015 paper from Peter Bubenik [2]. Section 6.2 follows the 2020 paper Henry Adams et al. [1] and Section 6.3 presents the results from Paweł Dłotko and Davide Gurnari 2023 paper, [7].

### 6.1 Persistence landscapes

Persistence landscapes provide a functional summary of persistence diagrams, converting them into a sequence of real-valued functions that capture topological features in a more tractable form. This section introduces the definition of persistence landscapes, their key properties, and stability results that relate the landscape distance to the bottleneck and Wasserstein distances between persistence diagrams.

Recall from Definition 1.1.14 that given some persistence module  $(V, \pi)$  and a fixed  $k$  the  $k$ -th persistent Betti number associated to  $x \leq y \in \mathbb{R}$  is given by

$$\beta_x^y(V) = \dim(\text{im } \pi_{x \leq y}).$$

**Lemma 6.1.1** (Lemma 1, [2]). *Let  $(V, \pi)$  be a persistence module and let  $a \leq b \leq c \leq d \in \mathbb{R}$ . Then  $\beta_b^c \geq \beta_a^d$ .*

*Proof.* Since  $\pi_{a \leq d} = \pi_{c \leq d} \circ \pi_{b \leq c} \circ \pi_{a \leq b}$ , the dimension of the image of  $\pi_{a \leq d}$  must be smaller

than the one of just  $\pi_{b \leq c}$ . □

**Lemma 6.1.2** (Lemma 2, [2]). *Let  $0 \leq h_1 \leq h_2$  be real numbers. Then  $\beta_{t-h_1}^{t+h_1} \geq \beta_{t-h_2}^{t+h_2}$ .*

*Proof.* ... □

Let  $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$  denote the extended real numbers.

**Definition 6.1.3** (Persistence landscape). A **persistence landscape** is a function  $\lambda: \mathbb{N} \times \mathbb{R} \rightarrow \overline{\mathbb{R}}$ , defined as

$$\lambda(k, t) := \sup\{m \geq 0 \mid \beta^{t-m, t+m} \geq k\}.$$

Note that this function can also be seen as a sequence of function  $\lambda_k: \mathbb{R} \rightarrow \overline{\mathbb{R}}$ , where  $\lambda_k(t) = \lambda(k, t)$ .

**Definition 6.1.4** ( $K$ -Lipschitz). Let  $(X, d_X), (Y, d_Y)$  be two metric spaces and let  $K > 0$ . A  **$K$ -Lipschitz** map is a map  $f: (X, d_X) \rightarrow (Y, d_Y)$ , such that for every  $x_1, x_2 \in X$ ,

$$d_Y(f(x_1), f(x_2)) \leq K d_X(x_1, x_2).$$

**Lemma 6.1.5** (Lemma 3, [2]). *Let  $\lambda_k: \mathbb{R} \rightarrow \overline{\mathbb{R}}$  be an element of a persistence landscape. The following properties are verified.*

1.  $\lambda_k(t) \geq 0$ ,
2.  $\lambda_k(t) \geq \lambda_{k+1}(t)$ ,
3.  $\lambda_k$  is 1-Lipschitz, that is, for  $t, s \in \mathbb{R}$ ,  $|\lambda_k(t) - \lambda_k(s)| \leq |t - s|$ .

*Proof.* Properties 1. and 2. came directly from the definition. For 3., suppose  $\lambda_k(s) \leq \lambda_k(t)$ . If  $\lambda_k(t) \leq |t - s|$  then, of course,  $\lambda_k(t) - \lambda_k(s) \leq \lambda_k(t) \leq |t - s|$ . Else, if  $\lambda_k(t) > |t - s|$ , we can take some  $h \in (0, \lambda_k(t) - |t - s|)$  verifying

$$t - \lambda_k(t) < s - h < s + h < t + \lambda_k(t).$$

Hence, by Lemma 6.1.1 □

**Definition 6.1.6** ( $p$ -landscape distance). Let  $V$  and  $W$  be two persistence modules, and let  $\lambda$  and  $\lambda'$  its corresponding persistence landscapes. Let  $1 \leq p \leq \infty$ . The  **$p$ -landscape distance between persistence modules  $V$  and  $W$**  is defined as

$$\Lambda_p(V, W) := \|\lambda - \lambda'\|_p.$$

Similarly, if  $D$  and  $D'$  are two persistence diagrams, and are  $\lambda$  and  $\lambda'$  its corresponding persistence landscapes, The  **$p$ -landscape distance between persistence diagrams**  $V$  and  $W$  is defined as

$$\Lambda_p(D, D') := \|\lambda - \lambda'\|_p.$$

**Theorem 6.1.7** (Theorem 17, [2]). *Let  $(V, \pi)$  and  $(W, \theta)$  two persistence modules. Then their  $\infty$ -landscape distance between them is a lower bound of their interleaving distance. That is*

$$\Lambda_\infty(V, W) \leq d_{\text{int}}(V, W)$$

*Proof.* Suppose  $V$  and  $W$  are  $\delta$ -interleaved, recall definition 1.2.6. Then, considering  $\pi_{t-m \leq t+m}$  and  $\pi_{t-m+\delta \leq t+m-\delta}$ , we have that  $t-m \leq t-m+\delta \leq t+m-\delta \leq t+m$ , hence by Lemma 6.1.1 we have

$$\beta_{t-m+\delta}^{t+m-\delta}(W) \geq \beta_{t-m}^{t+m}(V).$$

Now if  $\delta$  and  $\delta'$  are the corresponding persistence landscapes of  $V$  and  $W$ , by Definition 6.1.3, for every  $k \leq 1$ ,  $\lambda'(k, t) \geq \lambda(k, t) - \delta$ . Hence, it follows

$$\|\lambda - \lambda'\|_\infty \leq \delta.$$

□

**Theorem 6.1.8** (Theorem 12, [2]). *Consider the persistence modules  $V = F_x$ ,  $W = G_x$  given by the maps  $f, g: X \rightarrow \mathbb{R}$ . Then*

$$\Lambda_\infty(V, W) \leq \|f - g\|_\infty$$

*Proof.* The proofs follows from applying Theorem 6.1.7, Theorem 3 and Theorem 4.0.2, in such order. Hence

$$\Lambda_\infty(V, W) \leq d_{\text{int}}(V, W) = d_{\text{bot}}(D(f), D(g)) \leq \|f - g\|_\infty.$$

□

**Theorem 6.1.9** (Theorem 13, [2]). *Let  $D$  and  $D'$  be two persistence diagrams, then*

$$\Lambda_\infty(D, D') \leq d_{\text{bot}}(D, D').$$

*Proof.* ...

□

## 6.2 Persistence images

Persistence images offer an alternative vectorization by transforming persistence diagrams into finite-dimensional representations through kernel density estimation. Here, we define persistence surfaces and images, analyze their stability under the 1-Wasserstein distance, and derive bounds that ensure their reliability in machine learning and statistical applications.

**Definition 6.2.1** (Persistence surface). Let  $D$  be a persistence diagram. Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be the linear transformation  $T(x, y) = (x, y - x)$ . Fix a nonnegative weighting function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  that is zero along the horizontal axis, continuous and piecewise differentiable. Fix a differentiable probability distribution  $\phi_u: \mathbb{R}^2 \rightarrow \mathbb{R}$ , with mean  $u \in \mathbb{R}^2$ . The **persistence surface** associated to  $D$ , by  $f$  and  $\phi_u$  is a function  $\rho_D: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined as

$$\rho_D(z) := \sum_{u \in T(D)} f(u) \phi_u(z).$$

**Definition 6.2.2** (Persistence image). Let  $D$  be a persistence diagram with an associated persistence surface  $\rho_D$ . The **persistence image** of  $D$  by  $\rho_D$  is the collection  $\rho$  of **pixels**

$$I(\rho_D)_p := \iint_p \rho_D dy dx.$$

Let  $h: \mathbb{R}^2 \rightarrow \mathbb{R}$  be a differentiable function. We will denote the maximal norm of the gradient vector of  $h$  as

$$|\nabla h| = \sup_{z \in \mathbb{R}^2} \|\nabla h(z)\|_2.$$

Hence, by the fundamental theorem of calculus for line integrals, for every  $u, v \in \mathbb{R}^2$ , we have

$$|h(u) - h(v)| \leq |\nabla h| \|u - v\|_2.$$

Now, for a differentiable probability distribution  $\phi_u: \mathbb{R}^2 \rightarrow \mathbb{R}$ , with mean  $u \in \mathbb{R}^2$  we can denote  $|\nabla \phi_u|$  as  $|\nabla \phi|$  and  $\|\phi_u\|$  as  $\|\phi\|$  because both the maximal directional derivative and the uniform norm of a fixed differentiable probability function are invariable under translation.

**Lemma 6.2.3** (Lemma 3, [1]). Let  $u, v \in \mathbb{R}^2$ . Fix a nonnegative weighting function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  that is zero along the horizontal axis, continuous and piecewise differentiable. Fix a differentiable probability distribution  $\phi_u: \mathbb{R}^2 \rightarrow \mathbb{R}$ , with mean  $u \in \mathbb{R}^2$ . The following inequality asserts.

$$\|f(u)\phi_u - f(v)\phi_v\|_\infty \leq (\|f\|_\infty |\nabla \phi| + \|\phi\|_\infty |\nabla f|) \|u - v\|_2.$$

*Proof.* For any  $z \in \mathbb{R}^2$  we have

$$|\phi_u(z) - \phi_v(z)| = |\phi_u(z) - \phi_u(z + u - v)| \leq |\nabla \phi| \|u - v\|_2.$$

Hence,

$$\|\phi_u - \phi_v\| \leq \|\nabla\phi\| \|u - v\|_2.$$

Applying this, we can now develop

$$\begin{aligned} |f(u)\phi_u(z) - f(v)\phi_v(z)| &= |f(u)(\phi_u(z) - \phi_v(z)) + (f(u) - f(v))\phi_v(z)| \\ &\leq \|f\|_\infty \|\phi_u(z) - \phi_v(z)\| + \|\phi\|_\infty |f(u) - f(v)| \\ &\leq \|f\|_\infty \|\nabla\phi\| \|u - v\|_2 + \|\phi\|_\infty \|\nabla f\| \|u - v\|_2 \\ &= (\|f\|_\infty \|\nabla\phi\| + \|\phi\|_\infty \|\nabla f\|) \|u - v\|_2. \end{aligned}$$

□

Persistence surfaces are stable with respect to the 1-Wasserstein distance (see 1.4.7).

**Theorem 6.2.4** (Theorem 4, [1]). *Let  $D, D'$  be two persistent diagrams and  $\rho_D, \rho_{D'}$  two persistence surfaces associated to each diagram respectively. Then*

$$\|\rho_B - \rho_{B'}\|_\infty \leq \sqrt{10}(\|f\|_\infty \|\nabla\phi\| + \|\phi\|_\infty \|\nabla f\|) \omega_1(D, D').$$

*Proof.* Since  $D$  and  $D'$  consist of finitely many points, there exists a matching  $\gamma$  that achieves the infimum in the Wasserstein distance. Then

$$\begin{aligned} \|\rho_B - \rho_{B'}\|_\infty &= \left\| \sum_{u \in T(B)} f(u)\phi_u - \sum_{u \in T(B)} f(\gamma(u))\phi_{\gamma(u)} \right\|_\infty \\ &\leq \sum_{u \in T(B)} \|f(u)\phi_u - f(\gamma(u))\phi_{\gamma(u)}\|_\infty, \end{aligned}$$

We now apply Lemma 6.2.3 and then note that  $\|\cdot\|_2 \leq \sqrt{2}\|\cdot\|_\infty$  in  $\mathbb{R}^2$ . and that  $\|T(\cdot)\|_2 \leq \sqrt{5}\|\cdot\|_\infty$ ,

$$\begin{aligned} \sum_{u \in T(B)} \|f(u)\phi_u - f(\gamma(u))\phi_{\gamma(u)}\|_\infty &\leq \sqrt{2}(\|f\|_\infty \|\nabla\phi\| + \|\phi\|_\infty \|\nabla f\|) \sum_{u \in T(B)} \|u - \gamma(u)\|_\infty \\ &\leq \sqrt{10}(\|f\|_\infty \|\nabla\phi\| + \|\phi\|_\infty \|\nabla f\|) \sum_{u \in B} \|u - \gamma(u)\|_\infty \\ &= \sqrt{10}(\|f\|_\infty \|\nabla\phi\| + \|\phi\|_\infty \|\nabla f\|) W_1(B, B'). \end{aligned}$$

□

Persistence images are stable with respect to the 1-Wasserstein distance.



**Theorem 6.2.5** (Theorem 5, [1]). *Let  $A$  be the maximum area of any pixel in the image,  $A'$  the total area of the image, and  $n$  the number of pixels in the image. Then*

$$\begin{aligned}\|I(\rho_B) - I(\rho_{B'})\|_\infty &\leq \sqrt{10}A(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(D, D'), \\ \|I(\rho_B) - I(\rho_{B'})\|_1 &\leq \sqrt{10}A'(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(D, D'), \\ \|I(\rho_B) - I(\rho_{B'})\|_2 &\leq \sqrt{10n}A(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(D, D').\end{aligned}$$

*Proof.* Note for any pixel  $p$  with area  $A(p)$  we have

$$\begin{aligned}|I(\rho_B)_p - I(\rho_{B'})_p| &= \left| \iint_p \rho_B dydz - \iint_p \rho_{B'} dydx \right| \\ &= \left| \iint_p \rho_B - \rho_{B'} dydx \right| \\ &\leq A(p)\|\rho_B - \rho_{B'}\|_\infty \\ &\leq \sqrt{10}A(p)(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(B, B'),\end{aligned}$$

where the last inequality comes by applying Theorem 6.2.4. Hence we have

$$\begin{aligned}\|I(\rho_B) - I(\rho_{B'})\|_\infty &\leq \sqrt{10}A(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(B, B') \\ \|I(\rho_B) - I(\rho_{B'})\|_1 &\leq \sqrt{10}A'(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(B, B') \\ \|I(\rho_B) - I(\rho_{B'})\|_2 &\leq \sqrt{n}\|I(\rho_B) - I(\rho_{B'})\|_\infty \\ &\leq \sqrt{10n}A(\|f\|_\infty|\nabla\phi| + \|\phi\|_\infty|\nabla f|)\omega_1(B, B').\end{aligned}$$

□

## 6.3 Euler curves

The Euler characteristic curve provides a concise topological signature by tracking the evolution of the Euler characteristic across a filtration. This section examines its definition, connection to persistence diagrams, and stability properties, demonstrating how it serves as a stable summary for comparing filtered cell complexes.

**Definition 6.3.1** (Euler characteristic). Let  $K$  be a simplicial complex, and let  $K^p$  be its  $p$ -skeleton. The **Euler characteristic** of  $K$  is the alternating sum of the number of cells in its dimension

$$\chi(K) := \sum_d (-1)^d \#(K^d).$$

**Definition 6.3.2.** Let  $K$  be a simplicial complex. Let  $f: K \rightarrow \mathbb{R}$  be a filtration function. The **Euler characteristic curve** is a function that assign an Euler characteristic  $\chi$  for each filtration level  $t \in \mathbb{R}$ .

$$\text{ECC}(K, t) := \chi(K_t),$$

where  $K_t f^{-1}(-\infty, t]$ .

**Proposition 6.3.3** (Proposition 2, [7]). *Let  $X, Y$  be two filtered cell complexes and let  $D(X), D(Y)$  be its respective persistence diagrams. Then,*

$$\|\text{ECC}(X, t) - \text{ECC}(Y, t)\|_1 \leq \sum_k 2\omega_1(D(X), D(Y)).$$

*Proof.* ...

□

# Bibliography

- [1] ADAMS, H., EMERSON, T., KIRBY, M., NEVILLE, R., PETERSON, C., AND SHIPMAN, P. Persistence images: A stable vector representation of persistent homology. *Journal of Machine Learning Research* (2017).
- [2] BUBENIK, P. Statistical topological data analysis using persistence landscapes. *Journal of Machine Learning Research* (2015).
- [3] BUBENIK, P., AND WAGNER, A. Embeddings of persistence diagrams into hilbert spaces. *Journal of Applied and Computational Topology* (2020).
- [4] BURAGO, D., BURAGO, Y., AND IVANOV, S. *A Course in Metric Geometry*. American Mathematical Society, 2001.
- [5] CHAZAL, F., COHEN-STEINER, D., GUIBAS, L. J., MÉMOLI, F., AND OUDOT, S. Y. Gromov-Hausdorff stable signatures for shapes using persistence. *Eurographics Symposium on Geometry Processing* (2009).
- [6] CHAZAL, F., AND OUDOT, S. Y. Towards persistence-based reconstruction in Euclidean spaces. *Institut National de Recherche en Informatique et en Automatique* (2008).
- [7] DŁOTKO, P., AND GURNARI, D. Euler characteristic curves and profiles: a stable shape invariant for big data problems. *Oxford GigaScience* (2023).
- [8] EDELSBRUNNER, H., COHEN-STEINER, D., AND HARER, J. Stability of persistence diagrams. *University of Chicago* (2005).
- [9] GHRIST, R., AND MUHAMMAD, A. Coverage and hole-detection in sensor networks via homology. *Fourth International Symposium on Information Processing in Sensor Networks* (2005).
- [10] HATCHER, A. *Algebraic Topology*. Allen Hatcher, 2001.
- [11] HUNGERFORD, T. W. *Algebra*. Springer, 2000.
- [12] MUNKRES, J. R. *Elements of Algebraic Topology*. CRC Press, 1984.
- [13] NANDA, V. Computational algebraic topology, lecture notes. *University of Oxford* (2020).

- [14] POLTEROVICH, L., ROSEN, D., SAMVELYAN, K., AND ZHANG, J. *Topological Persistence in Geometry and Analysis*. American Mathematical Society, 2020.
- [15] RUDIN, W. *Real and Complex Analysis*. McGraw-Hill, 1987.
- [16] WANG, K. G. The basic theory of persistent homology. *University of Chicago* (2012).