


PERSISTENT π_1

& LOWER-CENTRAL-SERIES BARCODES

GONZALO ORTEGA CARPINTERO

gonzalo.ortega-carpintero@city.ac.uk




A joint work with Ximena Fernández at


CITY
ST GEORGE'S
UNIVERSITY OF LONDON

YTM 2026 - Monday, 29th June

THE LIMITATIONS OF H_*




$$T = S_1 \times S_1$$




$$X = S^2 \vee S^1 \vee S^1$$

All computations will be made with coefficients in a field $\mathbb{K}$.

THE LIMITATIONS OF H_*



$$T = S_1 \times S_1$$



$$X = S^2 \vee S^1 \vee S^1$$

$$H_0(T) = \mathbb{K} = H_0(X)$$


$$H_1(T) = \mathbb{K} \oplus \mathbb{K} = H_1(X)$$

$$H_2(T) = \mathbb{K} = H_2(X)$$


$$H_n(T) = 0 = H_n(X),$$

$$n \geq 3$$

THE LIMITATIONS OF H_*



$$T = S_1 \times S_1$$




$$X = S^2 \vee S^1 \vee S^1$$


$$H_1(T) = \mathbb{K} \oplus \mathbb{K} = H_1(X)$$

$$\pi_1(T) = \mathbb{Z} \oplus \mathbb{Z} \neq \mathbb{Z} * \mathbb{Z} = \pi_1(X)$$


THE LIMITATIONS OF PERSISTENT H_*



THE LIMITATIONS OF PERSISTENT H_*





T_1	$\longrightarrow$	T_2	$\longrightarrow$	T_3
$\downarrow$		$\downarrow$		$\downarrow$
$H_1(T_1)$	$\longrightarrow$	$H_1(T_2)$	$\longrightarrow$	$H_1(T_3)$



X_1	$\longrightarrow$	X_2	$\longrightarrow$	X_3
$\downarrow$		$\downarrow$		$\downarrow$
$H_1(X_1)$	$\longrightarrow$	$H_1(X_2)$	$\longrightarrow$	$H_1(X_3)$



THE LIMITATIONS OF PERSISTENT H_*



T_1	$\longrightarrow$	T_2	$\longrightarrow$	T_3
$\downarrow H_1$		$\downarrow H_1$		$\downarrow H_1$
$\mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} \oplus \mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} \oplus \mathbb{Z}$



X_1	$\longrightarrow$	X_2	$\longrightarrow$	X_3
$\downarrow H_1$		$\downarrow H_1$		$\downarrow H_1$
$\mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} \oplus \mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} \oplus \mathbb{Z}$

THE LIMITATIONS OF PERSISTENT H_*



$$\begin{array}{ccccc}
 T_1 & \longrightarrow & T_2 & \longrightarrow & T_3 \\
 \downarrow H_1 & & \downarrow H_1 & & \downarrow H_1 \\
 \mathbb{Z} & \longrightarrow & \mathbb{Z} \oplus \mathbb{Z} & \longrightarrow & \mathbb{Z} \oplus \mathbb{Z}
 \end{array}$$

$$\begin{array}{ccccc}
 X_1 & \longrightarrow & X_2 & \longrightarrow & X_3 \\
 \downarrow H_1 & & \downarrow H_1 & & \downarrow H_1 \\
 \mathbb{Z} & \longrightarrow & \mathbb{Z} \oplus \mathbb{Z} & \longrightarrow & \mathbb{Z} \oplus \mathbb{Z}
 \end{array}$$



$$PH_1(\mathcal{T}) = PH_1(\mathcal{X})$$


All barcodes coincide.

THE LIMITATIONS OF PERSISTENT H_*




WHAT WOULD WE LIKE TO DO?

THE LIMITATIONS OF PERSISTENT H_*



$$\begin{array}{ccccc}
 T_1 & \longrightarrow & T_2 & \longrightarrow & T_3 \\
 \downarrow & & \downarrow & & \downarrow \\
 H_1(T_1) & \longrightarrow & H_1(T_2) & \longrightarrow & H_1(T_3)
 \end{array}$$



$$\begin{array}{ccccc}
 X_1 & \longrightarrow & X_2 & \longrightarrow & X_3 \\
 \downarrow & & \downarrow & & \downarrow \\
 H_1(X_1) & \longrightarrow & H_1(X_2) & \longrightarrow & H_1(X_3)
 \end{array}$$

PERSISTENT π_1

$$\begin{array}{ccccc}
 T_1 & \longrightarrow & T_2 & \longrightarrow & T_3 \\
 \downarrow & & \downarrow & & \downarrow \\
 \pi_1(T_1) & \longrightarrow & \pi_1(T_2) & \longrightarrow & \pi_1(T_3)
 \end{array}$$

$$\begin{array}{ccccc}
 X_1 & \longrightarrow & X_2 & \longrightarrow & X_3 \\
 \downarrow & & \downarrow & & \downarrow \\
 \pi_1(X_1) & \longrightarrow & \pi_1(X_2) & \longrightarrow & \pi_1(X_3)
 \end{array}$$

Convention: All spaces are path-connected and contain a common base point.

CAN π_1 BE ALGORITHMICALLY COMPUTED?

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

*(All CW complexes will be **finite** and **path-connected** in this presentation.)*

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*


*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.




T

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.




T

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.




T

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.



$$T \cong \langle e_2, e_4, e_5, e_7, e_8 \mid e_5, e_2^{-1}e_7, e_5^{-1}, e_7^{-1}e_4^{-1}e_2e_8, e_8^{-1}e_4 \rangle$$

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Naive computation: Using the **spanning tree** algorithm.

Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.

Let K be a regular CW complex and let M be an acyclic matching on the cells of K .

Theorem (Forman, 1995; Chari 2000): K is homotopy equivalent to a CW complex with exactly one cell of dimension p for each critical (ie., unmatched) cell of dimension p .

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.

Let K be a regular CW complex and let M be an acyclic matching on the cells of K with only one 0-dimensional critical cell.

Theorem (Fernández, 2024): A presentation $\mathcal{P} = \langle X \mid R \rangle$ of $\pi_1(K)$ with

$$\#X = \#1\text{-dim critical cells}$$

$$\#R = \#2\text{-dim critical cells}$$


can be computed with complexity $\mathcal{O}(N^2)$, where

$$N = \#\text{cells in the 2-skeleton of } K.$$

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.




T

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.


Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.



CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.


Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.



CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

*We can compute a **presentation** of π_1 of a CW complex.

Efficient computation: Using **discrete Morse theory** to obtain a reduced presentation.



$$T \cong \langle e_2, e_4 \mid e_2^{-1} e_4^{-1} e_2 e_4 \rangle$$

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN π_1 BE COMPUTED OVER A FILTRATION?

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED?

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

Let $K_\bullet$ be a path-connected based filtration of finite CW-complexes.

$$\begin{array}{ccccccc} K_1 & \longrightarrow & K_2 & \longrightarrow & \cdots & \longrightarrow & K_n \\ \downarrow & & \downarrow & & & & \downarrow \\ \pi_1(K_1) & \longrightarrow & \pi_1(K_2) & \longrightarrow & \cdots & \longrightarrow & \pi_1(K_n) \end{array}$$

Naive computation: Compute π_1 presentations **independently** over every element in the filtration.

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

Let $K_\bullet$ be a path-connected based filtration of finite CW-complexes.

$$\begin{array}{ccccccc} K_1 & \longrightarrow & K_2 & \longrightarrow & \cdots & \longrightarrow & K_n \\ \downarrow & & \downarrow & & & & \downarrow \\ \pi_1(K_1) & \longrightarrow & \pi_1(K_2) & \longrightarrow & \cdots & \longrightarrow & \pi_1(K_n) \end{array}$$

Naive computation: Compute π_1 presentations **independently** over every element in the filtration.

Efficient computation: Perform a **single computation** over the whole filtration.

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

Efficient computation: Perform a **single computation** over the whole filtration.

Theorem (Fernández, O., 2026): Let $M_\bullet$ be an acyclic matching with only one 0-dim critical cell over $K_\bullet$. Presentations $\mathcal{P}_i = \langle X_i \mid R_i \rangle$ of $\pi_1(K_i)$ with

$\#X_i = \#1\text{-dim critical cells of } K_i,$

$\#R_i = \#2\text{-dim critical cells of } K_i$

and morphisms $\psi_{i \rightarrow j} : \mathcal{P}_i \rightarrow \mathcal{P}_j$ such that the following commutes

$$\begin{array}{ccc} K_i & \hookrightarrow & K_j \\ \downarrow & & \downarrow \\ \mathcal{P}_i & \xrightarrow{\psi_{i \rightarrow j}} & \mathcal{P}_j \end{array}$$

can be computed in $\mathcal{O}(N^2)$, $N = \#$ cells in the 2-skeleton of K .

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

Efficient computation: Perform a **single computation** over the whole filtration.

SAGE Math implementation

```
def filtrated_fundamental_group(self, fil, verbose = True):
    S = set()
    R = []
    filtrated_fundamental_group = []
    maps = [[None] * (fil+1) for _ in range(len(self.cells_lists[1]))]


    for f in range(0, fil+1):

        for cell in self.filtration_unmatched_2_cells[f]:
            R.append(cell.boundary_cells)

        for cell in self.filtration_matched_1_cells_with_0_cells[f]:
            cell.equivalence = '1'
            maps[cell_count][f] = 1
```


CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*




CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***




CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***




CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*




CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***



CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***



CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***

$$P\pi_1(\mathcal{T}) = \langle t_1 \mid - \rangle \rightarrow \langle t_1, t_2 \mid - \rangle \rightarrow \langle t_1, t_2 \mid t_1^{-1}t_2^{-1}t_1t_2 \rangle,$$

$$P\pi_1(\mathcal{X}) = \langle x_1 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle.$$

CAN π_1 BE ALGORITHMICALLY COMPUTED? YES*

CAN PERSISTENT π_1 BE COMPUTED? YES*

$$P\pi_1(\mathcal{T}) = \langle t_1 \mid - \rangle \rightarrow \langle t_1, t_2 \mid - \rangle \rightarrow \langle t_1, t_2 \mid t_1^{-1}t_2^{-1}t_1t_2 \rangle,$$

$$P\pi_1(\mathcal{X}) = \langle x_1 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle.$$

CAN WE MAKE π_1 BARCODES AS IN H_* ?

CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***

$$P\pi_1(\mathcal{T}) = \langle t_1 \mid - \rangle \rightarrow \langle t_1, t_2 \mid - \rangle \rightarrow \langle t_1, t_2 \mid t_1^{-1}t_2^{-1}t_1t_2 \rangle,$$

$$P\pi_1(\mathcal{X}) = \langle x_1 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle.$$

CAN WE MAKE π_1 BARCODES AS IN H_* ? **NO***

CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***

$$P\pi_1(\mathcal{T}) = \langle t_1 \mid - \rangle \rightarrow \langle t_1, t_2 \mid - \rangle \rightarrow \langle t_1, t_2 \mid t_1^{-1}t_2^{-1}t_1t_2 \rangle,$$

$$P\pi_1(\mathcal{X}) = \langle x_1 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle.$$

CAN WE MAKE π_1 BARCODES AS IN H_* ? **NO***

- The groups $\pi_1(K_\alpha)$ in $K_\bullet \rightarrow P\pi_1(K)_\bullet$.

$$\pi_1(K_1) \longrightarrow \pi_1(K_2) \longrightarrow \cdots \longrightarrow \pi_1(K_n)$$

are generally **non-abelian**.

CAN π_1 BE ALGORITHMICALLY COMPUTED? **YES***

CAN PERSISTENT π_1 BE COMPUTED? **YES***

$$P\pi_1(\mathcal{T}) = \langle t_1 \mid - \rangle \rightarrow \langle t_1, t_2 \mid - \rangle \rightarrow \langle t_1, t_2 \mid t_1^{-1}t_2^{-1}t_1t_2 \rangle,$$

$$P\pi_1(\mathcal{X}) = \langle x_1 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle \rightarrow \langle x_1, x_2 \mid - \rangle.$$

CAN WE MAKE π_1 BARCODES AS IN H_* ? **NO***

- The groups $\pi_1(K_\alpha)$ in $K_\bullet \rightarrow P\pi_1(K)$.

$$\pi_1(K_1) \longrightarrow \pi_1(K_2) \longrightarrow \cdots \longrightarrow \pi_1(K_n)$$

are generally **non-abelian**.

- Finite presentations are **not group invariants** in general.
- Ordinary barcode decompositions do not apply.

IDEA:

LOOK FOR AN ABELIAN INVARIANT OF π_1

LOWER CENTRAL QUOTIENTS

LOWER CENTRAL QUOTIENTS

Definition: The **lower central series** of a group G is the descending series of normal subgroups

$$G = \gamma_1(G) \triangleright \gamma_2(G) \triangleright \cdots \triangleright \gamma_n(G) \triangleright \cdots$$

where

$$\gamma_{n+1}(G) = [\gamma_n(G), G] = \langle [h, g] : h \in \gamma_n(G), g \in G \rangle.$$

LOWER CENTRAL QUOTIENTS

Definition: The **lower central series** of a group G is the descending series of normal subgroups

$$G = \gamma_1(G) \supseteq \gamma_2(G) \supseteq \cdots \supseteq \gamma_n(G) \supseteq \cdots$$

where

$$\gamma_{n+1}(G) = [\gamma_n(G), G] = \langle [h, g] : h \in \gamma_n(G), g \in G \rangle.$$

Definition: We define the n^{th} -**lower central quotient** of G as

$$lcq_n(G) := \frac{\gamma_n(G)}{\gamma_{n+1}(G)}.$$

LOWER CENTRAL QUOTIENTS

Example: Let $G = \pi_1(K)$.

LOWER CENTRAL QUOTIENTS

Example: Let $G = \pi_1(K)$.

$$lcq_1(G) = \frac{G}{[G, G]} \cong H_1(K),$$

$$lcq_2(G) = \frac{[G, G]}{[[G, G], G]},$$

$$lcq_3(G) = \frac{[[G, G], G]}{[[[G, G], G], G]},$$

...

LOWER CENTRAL QUOTIENTS

Example (Wedge of circles): Let $K = \bigvee_n S^1$, $G = \pi_1 (K) = F_n$.

$$lcq_1(F_n) \cong \mathbb{Z}^n,$$

$$lcq_2(F_n) \cong \mathbb{Z}^{\binom{n}{2}},$$

$$lcq_3(F_n) \cong \mathbb{Z}^{n \binom{n}{2} - \binom{n}{3}}.$$

LOWER CENTRAL QUOTIENTS

Example (Wedge of circles): Let $K = \bigvee_n S^1$, $G = \pi_1(K) = F_n$.

$$lcq_1(F_n) \cong \mathbb{Z}^n,$$

$$lcq_2(F_n) \cong \mathbb{Z}^{\binom{n}{2}},$$

$$lcq_3(F_n) \cong \mathbb{Z}^{n\binom{n}{2} - \binom{n}{3}}.$$

For every n , $lcq_n(G)$ *is abelian*.

LOWER CENTRAL QUOTIENTS

For every n , $lcq_n(G)$ is abelian.

For every n , $lcq_n(G)$ can be algorithmically computed from a presentation of G . (Sims, 1994)

LOWER CENTRAL QUOTIENTS

For every n , $lcq_n(G)$ is abelian.

For every n , $lcq_n(G)$ can be algorithmically computed from a presentation of G . (Sims, 1994)

The complexity of the computation of $lcq_n(G)$ depends on the number of generators and relations of the presentation of G we are using.

PERSISTENT LOWER CENTRAL QUOTIENTS

PERSISTENT LOWER CENTRAL QUOTIENTS

Definition: Let $\mathbb{K}$ a fixed field. We define the $\mathbb{K}$ -**tensor** n^{th} -**lower central quotient** as

$$lcq_n^{\mathbb{K}} := lcq_n \otimes_{\mathbb{Z}} \mathbb{K}$$

PERSISTENT LOWER CENTRAL QUOTIENTS

$$\begin{array}{ccccccc}
 K_1 & \longrightarrow & K_2 & \longrightarrow & \cdots \longrightarrow & & K_n \\
 \downarrow & & \downarrow & & & & \downarrow \\
 \pi_1(K_1) & \longrightarrow & \pi_1(K_2) & \longrightarrow & \cdots \longrightarrow & & \pi_1(K_n) \\
 \downarrow \xi_1 & & \downarrow \xi_2 & & & & \downarrow \xi_n \\
 lcq_n^{\mathbb{K}}(\pi_1(K_1)) & \longrightarrow & lcq_n^{\mathbb{K}}(\pi_1(K_2)) & \longrightarrow & \cdots \longrightarrow & & lcq_n^{\mathbb{K}}(\pi_1(K_n))
 \end{array}$$

Here we obtain a functor

$$\xi_\alpha : \mathbf{Grp} \rightarrow \mathbf{Vect}_{\mathbb{K}}$$

$$G \mapsto \frac{\gamma_n(G)}{\gamma_{n+1}(G)} \otimes_{\mathbb{Z}} \mathbb{K}$$

PERSISTENT LOWER CENTRAL QUOTIENTS

Every $lcq_n(\pi_1(K_\alpha))$ is abelian so

$$lcq_n^{\mathbb{K}}(\pi_1(K_1)) \longrightarrow lcq_n^{\mathbb{K}}(\pi_1(K_2)) \longrightarrow \cdots \longrightarrow lcq_n^{\mathbb{K}}(\pi_1(K_n))$$

is a **persistent module** over the field $\mathbb{K}$ (ie. a graded $\mathbb{K}[t]$ -module).

PERSISTENT LOWER CENTRAL QUOTIENTS



Every $lcq_n(\pi_1(K_\alpha))$ is abelian so

$$lcq_n^{\mathbb{K}}(\pi_1(K_1)) \longrightarrow lcq_n^{\mathbb{K}}(\pi_1(K_2)) \longrightarrow \cdots \longrightarrow lcq_n^{\mathbb{K}}(\pi_1(K_n))$$

is a **persistent module** over the field $\mathbb{K}$ (ie. a graded $\mathbb{K}[t]$ -module).



*We can compute **barcodes**. (Skraba & Vejdemo-Johansson, 2013)*

LOWER CENTRAL QUOTIENTS




T_1	$\longrightarrow$	T_2	$\longrightarrow$	T_3
$\downarrow \pi_1$		$\downarrow \pi_1$		$\downarrow \pi_1$
$\mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} * \mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} \oplus \mathbb{Z}$
$\downarrow lcq_2^{\mathbb{K}}$		$\downarrow lcq_2^{\mathbb{K}}$		$\downarrow lcq_2^{\mathbb{K}}$
0	$\longrightarrow$	$\mathbb{K}$	$\longrightarrow$	0

X_1	$\longrightarrow$	X_2	$\longrightarrow$	X_3
$\downarrow \pi_1$		$\downarrow \pi_1$		$\downarrow \pi_1$
$\mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} * \mathbb{Z}$	$\longrightarrow$	$\mathbb{Z} * \mathbb{Z}$
$\downarrow lcq_2^{\mathbb{K}}$		$\downarrow lcq_2^{\mathbb{K}}$		$\downarrow lcq_2^{\mathbb{K}}$
0	$\longrightarrow$	$\mathbb{K}$	$\longrightarrow$	$\mathbb{K}$




LOWER CENTRAL QUOTIENTS

Example: lcq_n barcodes of links.




Example: lcq_n barcodes of links. (π_1)



$$\pi_1(\mathbb{R}^3 \setminus U_1) = \langle u_1, u_2, u_3 \mid - \rangle$$

$$\pi_1(\mathbb{R}^3 \setminus U_2) = \langle u_1, u_2 \mid - \rangle$$


$$\pi_1(\mathbb{R}^3 \setminus U_3) = \langle u_1 \mid - \rangle$$



$$\pi_1(\mathbb{R}^3 \setminus L_1) = \langle l_1, l_2, l_3 \mid [l_1, l_2], [l_1 l_3, l_2] \rangle$$

$$\pi_1(\mathbb{R}^3 \setminus L_2) = \langle l_1, l_2 \mid [l_1, l_2] \rangle$$

$$\pi_1(\mathbb{R}^3 \setminus L_3) = \langle l_1 \mid - \rangle$$





$$\pi_1(\mathbb{R}^3 \setminus R_1) = \langle r_1, r_2, r_3 \mid r_3 r_2^{-1} r_3^{-1} r_1^{-1} r_3 r_2 r_3^{-1} r_2^{-1} r_1 r_2, r_3 r_2 r_3^{-1} r_1 r_3 r_1^{-1} r_2^{-1} r_1 r_3^{-1} r_1^{-1} \rangle$$


$$\pi_1(\mathbb{R}^3 \setminus R_2) = \langle r_1, r_2 \mid - \rangle$$

$$\pi_1(\mathbb{R}^3 \setminus R_3) = \langle r_1 \mid - \rangle$$

Example: lcq_n barcodes of links. $(lcq_1^{\mathbb{K}})$









$\mathbb{R}^3 \setminus U_1$	$\longrightarrow$	$\mathbb{R}^3 \setminus U_2$	$\longrightarrow$	$\mathbb{R}^3 \setminus U_3$	$\mathbb{R}^3 \setminus L_1$	$\longrightarrow$	$\mathbb{R}^3 \setminus L_2$	$\longrightarrow$	$\mathbb{R}^3 \setminus L_3$
$\downarrow H_1$		$\downarrow H_1$		$\downarrow H_1$	$\downarrow H_1$		$\downarrow H_1$		$\downarrow H_1$
$\mathbb{K}^3$	$\longrightarrow$	$\mathbb{K}^2$	$\longrightarrow$	$\mathbb{K}$	$\mathbb{K}^3$	$\longrightarrow$	$\mathbb{K}^2$	$\longrightarrow$	$\mathbb{K}$




$\mathbb{R}^3 \setminus R_1$	$\longrightarrow$	$\mathbb{R}^3 \setminus R_2$	$\longrightarrow$	$\mathbb{R}^3 \setminus R_3$
$\downarrow H_1$		$\downarrow H_1$		$\downarrow H_1$
$\mathbb{K}^3$	$\longrightarrow$	$\mathbb{K}^2$	$\longrightarrow$	$\mathbb{K}$

Example: lcq_n barcodes of links. $(lcq_1^{\mathbb{K}})$




Example: lcq_n barcodes of links. $(lcq_2^{\mathbb{K}})$



$\mathbb{R}^3 \setminus U_1 \longrightarrow \mathbb{R}^3 \setminus U_2 \longrightarrow \mathbb{R}^3 \setminus U_3$
 $\mathbb{R}^3 \setminus L_1 \longrightarrow \mathbb{R}^3 \setminus L_2 \longrightarrow \mathbb{R}^3 \setminus L_3$

$\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$

$\mathbb{K}^3 \longrightarrow \mathbb{K} \longrightarrow 0$
 $\mathbb{K} \longrightarrow 0 \longrightarrow 0$









$\mathbb{R}^3 \setminus R_1 \longrightarrow \mathbb{R}^3 \setminus R_2 \longrightarrow \mathbb{R}^3 \setminus R_3$


$\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$
 $\downarrow lcq_2^{\mathbb{K}}$

$\mathbb{K}^3 \longrightarrow \mathbb{K} \longrightarrow 0$

Example: lcq_n barcodes of links. $(lcq_2^{\mathbb{K}})$



Example: lcq_n barcodes of links. $(lcq_3^{\mathbb{K}})$



$\mathbb{R}^3 \setminus U_1$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus U_2$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus U_3$


$\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$

$\mathbb{K}^8$
 $\longrightarrow$
 $\mathbb{K}^2$
 $\longrightarrow$
 0

$\mathbb{R}^3 \setminus L_1$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus L_2$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus L_3$

$\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$

$\mathbb{K}^2$
 $\longrightarrow$
 0
 $\longrightarrow$
 0









$\mathbb{R}^3 \setminus R_1$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus R_2$
 $\longrightarrow$
 $\mathbb{R}^3 \setminus R_3$

$\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$
 $\downarrow lcq_3^{\mathbb{K}}$

$\mathbb{K}^6$
 $\longrightarrow$
 $\mathbb{K}^2$
 $\longrightarrow$
 0

Example: lcq_n barcodes of links. $(lcq_3^{\mathbb{K}})$



GAP CODE

```
1 LoadPackage("nq");
2
3 LcsQuotient := function(group, i)
4   local nq, lcs;
5   nq := NilpotentQuotient(group, i+1);
6   lcs := LowerCentralSeries(nq);
7   if i < Length(lcs) then
8     return AbelianInvariants(lcs[i] / lcs[i+1]);
9   fi;
10  return TrivialGroup();
11 end;
```

Using GAP 4 NQ Package (Nickel, 2007).

SUMMARY

- PH_* has limitations to distinguish certain kinds of filtrations.

SUMMARY

- PH_* **has limitations** to distinguish certain kinds of filtrations.
- We have design an **efficient algorithm to compute** $P\pi_1$ presentations.

SUMMARY

- PH_* has limitations to distinguish certain kinds of filtrations.
- We have design an efficient algorithm to compute $P\pi_1$ presentations.
- We found a way of computing $P\pi_1$ barcodes using lcq_n .

ONGOING AND FUTURE WORK

- Look for a computationally **efficient algorithm to compute** $lcq_n(G)$.

ONGOING AND FUTURE WORK

- Look for a computationally **efficient algorithm to compute** $lcq_n(G)$.
- Apply the methods to **metric spaces** and **real data**.

ONGOING AND FUTURE WORK

- Look for a computationally **efficient algorithm to compute** $lcq_n(G)$.
- Apply the methods to **metric spaces** and **real data**.
- Explore for **abelian invariants** beyond lcq_n .

REFERENCES

- **Robin Forman**. Morse theory for cell complexes. *Advances in Mathathematics*, 1998.
- **Ximena Fernández**. Morse theory for group presentations. *Transactions of the American Mathematical Society*, 2024.
- **Charles C. Sims**. Computation with finitely presented groups. *Cambridge University Press*, 1994.
- **Primoz Skraba** and **Mikael Vejdemo-Johansson**. Persistence modules: Algebra and algorithms. *Mathematics of Computation*, 2013.
- **Werner Nickel**. A gap 4 package computing nilpotent factor groups of finitely presented groups. *GAP 4 Package*, 2007.