

Chapter 6

Brief introduction on how to find
zeros of real functions

Solving explicitly the unconstrained problem

Remark. Let $f : D \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a \mathbb{C}^1 function. We have proven that to solve the problem

$$\min_{x \in \mathbb{R}^n} f(x)$$


it is necessary to find out points (vectors) x^* such that $\nabla f(x^*) = 0$.

Remark. A possible strategy for doing so is to start at a given vector $x_0 \in D$ and construct a sequence x_k such that

$$x_k = \min_{\alpha \in \mathbb{R}} f(x_{k-1} + \alpha p_k), \quad p_k \in \mathbb{R}^n.$$

Consequence. It is worthy to first do a quick overview for the one-dimensional problem of finding zeroes of functions.

Example



$$f(x) = x^3 - 1.35 x^2 + 0.49 x - 0.05$$

Strategy

Let $f : \mathcal{I} \subset \mathbb{R} \rightarrow \mathbb{R}$.

- 1 Location: Where the zeros are?
- 2 Separation (or uniqueness): Determine a domain (interval) with a unique zero.
- 3 Approximation (root-finding algorithms): Construct a sequence x_k in the domain above such that

$$x_k \rightarrow x^* \quad \text{with} \quad f(x^*) = 0.$$

Location: Bolzano's Theorem

Theorem (Bolzano). Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function satisfying $f(a)f(b) < 0$. Then, there exists $x^* \in (a, b)$ such that $f(x^*) = 0$.

Proof.

- (0) Let $[a_0, b_0] = [a, b]$ and let $n = 0$.
- (1) Compute $c_{n+1} = (a_n + b_n)/2$.
- (2) One (and only one) below can happen.
 - (2.a) $f(c_{n+1}) = 0$. Then $x^* = c_{n+1}$.
 - (2.b) $f(a_n)f(c_{n+1}) < 0$. Then $[a_{n+1}, b_{n+1}] = [a_n, c_{n+1}]$;
 - (2.c) $f(c_{n+1})f(b_n) < 0$. Then $[a_{n+1}, b_{n+1}] = [c_{n+1}, b_n]$.
- (3) Do $n = n + 1$ and move to step (1).

Location: Bolzano's Theorem

Proof. To conclude the theorem we argue as follows.

- (i) If the process stops, i.e., (2.a), we have found x^* .
- (ii) Otherwise we have constructed an infinite sequence of nested intervals

$$[a, b] = [a_0, b_0] \supset [a_1, b_1] \supset \dots \supset [a_n, b_n] \supset \dots,$$

(iii) Clearly $\ell([a_n, b_n]) = \frac{|b_0 - a_0|}{2^n} \rightarrow 0$

(iv) Let

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = x^*.$$

(v) Since $f(a_n)f(b_n) < 0$ for all $n \geq 0$, we have

$$0 \geq \lim_{n \rightarrow \infty} f(a_n)f(b_n) = (f(x^*))^2 \quad (\Rightarrow f(x^*) = 0).$$

Uniqueness: Rolle's Theorem


Theorem (Rolle). Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function in $[a, b]$ and derivable in (a, b) . Suppose $f(a) = f(b)$. Then there exists $\zeta^* \in (a, b)$ such that $f'(\zeta^*) = 0$.

Corollary. Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function in $[a, b]$ and derivable in (a, b) . Assume that $f(a)f(b) < 0$ and $f'(x) \neq 0$ for all $x \in (a, b)$. Then there exists a **unique** $x^* \in (a, b)$ such that $f(x^*) = 0$.

Proof. From Bolzano's Theorem it is clear that there exists $x^* \in (a, b)$ such that $f(x^*) = 0$. Assume there exists $y^* \neq x^*$ such that $f(y^*) = 0$. W.l.o.g. take $y^* > x^*$. Then $f|_{[x^*, y^]}$ is in the hypothesis of Rolle's Theorem and there exists $\zeta^* \in (x^*, y^*) \subset (a, b)$ such that $f'(\zeta^*) = 0$, a contradiction.


Root-finding algorithm: Bisection method

Assume there is a unique zero x^* of f in $[a, b]$. From [Bolzano's Theorem](#) we can construct a sequence $\{c_n\}_{n \geq 0}$ such that $c_n \rightarrow x^*$. The method always converge. We stop to process at $k = k_0$ with $|f(c_{k_0})| < \varepsilon$ for a given $\varepsilon > 0$.



The stop condition might be incorrect.

Bisection method



Bisection method

Computing $\alpha_1 = 0.99639033$ with $\varepsilon < 10^{-8}$.

| n | a_n | b_n | c_{n+1} | $f(c_{n+1})$ |
|-----|-------------|-------------|-------------|--------------|
| 0 | 0.950000000 | 1.050000000 | 1.000000000 | -1.3e-03 |
| 1 | 0.950000000 | 1.000000000 | 0.975000000 | 8.0e-03 |
| 2 | 0.975000000 | 1.000000000 | 0.987500000 | 3.2e-03 |
| 3 | 0.987500000 | 1.000000000 | 0.993750000 | 9.5e-04 |
| 4 | 0.993750000 | 1.000000000 | 0.996875000 | -1.7e-04 |
| 5 | 0.993750000 | 0.996875000 | 0.995312500 | 3.9e-04 |
| 10 | 0.996386719 | 0.996484375 | 0.996435547 | -1.6e-05 |
| 15 | 0.996389771 | 0.996392822 | 0.996391296 | -3.5e-07 |
| 20 | 0.996390247 | 0.996390343 | 0.996390295 | 1.2e-08 |
| 24 | 0.996390325 | 0.996390331 | 0.996390328 | 1.1e-09 |

Newton's method (The analysis)

Problem. Improve the efficiency of the bisection method under the extra hypothesis that f is derivable.


Newton's method. Instead of solving $f(x) = 0$ we argue as follows.

- (a) Assume $x_0 \approx x^*$ where x^* is a solution of $f(x) = 0$.
- (b) Consider the linear function which better approximate f near x_0 ; that is, $L(f, x_0) = f(x_0) + f'(x_0)(x - x_0)$.
- (c) Then consider $L(f, x_0) = 0$.
- (d) Set

$$x_1 := x_0 - \frac{f(x_0)}{f'(x_0)} \quad \text{and} \quad x_{n+1} := x_n - \frac{f(x_n)}{f'(x_n)}$$

- (e) We have $x_n \rightarrow x^*$ as $n \rightarrow \infty$.

Newton's method (The geometry)



Newton's method (Remarks)


- We have assumed $x_0 \approx x^*$.
- Morally, we are assuming $f'(x) \neq 0$ near x^* .
- **Stop criteria.** We might use either $|x_{n+1} - x_n| < \varepsilon$ or $|f(x_{n+1})| < \delta$, on $\varepsilon > 0, \delta > 0$. Both criteria might have problems.
- **Convergence.** The speed of convergence is much better, locally, than the bisection method.
- The **idea** can be generalized to higher dimension.

Newton's method (Example)

Computing α_1 , a zero of $f(x) = \exp(x - 0.5) - 2x + 0.35$.

| n | x_n | $f(x_n)$ | $f'(x_n)$ | $ x_n - x_{n-1} $ |
|-----|----------------|------------------|-------------------|-------------------|
| 0 | 0.950000000000 | 1.8312185490e-02 | -4.3168781451e-01 | |
| 1 | 0.99241997313 | 1.4312185890e-03 | -3.6372883516e-01 | 0.4e-01 |
| 2 | 0.99635482363 | 1.2683863782e-05 | -3.5727766888e-01 | 0.4e-02 |
| 3 | 0.99639032504 | 1.0352153579e-09 | -3.5721934888e-01 | 0.4e-04 |
| 4 | 0.99639032794 | 1.1102230246e-16 | -3.5721934412e-01 | 0.3e-08 |

Newton's method (Global approach)



Newton's method as a fixed point method

Newton's method can be viewed as a fixed point method. Indeed we have found a function g such that

$$f(x^*) = 0 \iff g(x^*) = x^*.$$


In other words our unknown value $x = x^*$ instead of a zero of f becomes a fixed point of g . Precisely

$$g(x) := N_f(x) = x - \frac{f(x)}{f'(x)}.$$


Remark. Newton's method is just a particular case of the fixed point theory to find out zeros of functions. The advantage (fixed points instead of zeros) is that, under certain conditions, the map g gives a natural path to create $x_n \rightarrow x^*$.

Fixed point method (example)


Exercise. Suppose we want to compute $x - \cos(x) = 0$. Then we might consider to iterate $g(x) = \cos(x)$, starting with a value close to a solution.



$$f(x) = \cos(x) - x$$



$$g(x) = \cos(x)$$



$$N_f(x) = x + \tan^{-1}(x)$$

Fixed point theory not always work

Exercise. We want to find out the unique zero of

$$f(x) = x - \exp(-x) = 0 \quad (x^* = 0.567143)$$

- As a fixed point of $g_1(x) = \exp(-x)$.
- As a fixed point of $g_2(x) = -\log(x)$.
- As a fixed point of $g_3(x) := N_f(x) = x - \frac{x - \exp(-x)}{1 + \exp(x)}$

$$g_2(x) = -\log(x) \rightarrow$$

Not convergent

| n | x_n | n | x_n |
|-----|----------|-----|-----------|
| 0 | 0.55 | 5 | 0.895394 |
| 1 | 0.597837 | 6 | 0.110492 |
| 2 | 0.514437 | 7 | 2.202816 |
| 3 | 0.664682 | 8 | -0.789737 |
| 4 | 0.408447 | | |

Fixed point Theorem

Theorem (fixed point). Let $g : [a, b] \longrightarrow [a, b]$ a continuous function. Suppose g is derivable in (a, b) and it satisfies

$$|g'(x)| \leq k < 1 \quad \forall x \in (a, b) .$$

Then, for all $x_0 \in (a, b)$ we have

$$x_n := g(x_{n-1}) \longrightarrow x^*$$

with $g(x^*) = x^*$. Moreover the following inequalities hold

$$|x_n - \alpha| \leq \frac{k^n}{1 - k} |x_0 - x_1| \quad \text{and} \quad |x_n - \alpha| \leq \frac{k}{1 - k} |x_n - x_{n-1}| .$$

Truncate conditions

- The blue inequality

$$|x_n - \alpha| \leq \frac{k^n}{1 - k} |x_0 - x_1|$$

is a **priori** estimate of the number of iterates.

- The red inequality


$$|x_n - \alpha| \leq \frac{k}{1 - k} |x_n - x_{n-1}|$$

is a **key information** to decide the stop condition.


Fixed point Theorem (example)

Exercise: We are calculating the unique zero of $f(x) = x - e^{-x}$. We use (see above) $g(x) = \exp(-x)$.

- $g : [0.2, 1] \mapsto [0.2, 1]$.
- $|g'(x)| < k = 0.82$ for all $x \in [0.2, 1]$.
- **Fixed point Theorem** The function g has a unique fixed point $x^* \in [0.2, 1]$ (i.e., $x = x^*$ is the unique zero of f in $[0.2, 1]$).



$$f(x) = x - e^{-x}$$



$$g(x) = e^{-x}$$

Fixed point Theorem (example)

The *a priori* bound is

$$\frac{0.82^n}{0.18} |0.5 - 0.60653| < 10^{-8} \quad \Rightarrow \quad n > 90.$$

| n | x_n | $ x_n - x_{n-1} $ | $\frac{k}{1-k} x_n - x_{n-1} $ |
|-----|----------------|-------------------|---------------------------------|
| 0 | 5.00000000e-01 | | |
| 1 | 6.06530660e-01 | 1.0653066e-01 | 4.8530633e-1 |
| 2 | 5.45239212e-01 | 6.12914478e-02 | 2.792166e-1 |
| 3 | 5.79703095e-01 | 3.44638830e-02 | 1.570021334e-1 |
| 10 | 5.66907213e-01 | 6.52421327e-04 | 2.9721416e-3 |
| 20 | 5.67142478e-01 | 2.24611113e-06 | 1.02323e-5 |
| 30 | 5.67143288e-01 | 7.73319819e-09 | 3.5229e-8 |

Remark. Observe that $n \approx 30$ its enough. This is so because of the bounds of the derivative.

Fixed point Theorem (example)

Remark. We want to find out the unique zero of

$$f(x) = x - \exp(-x) = 0 \quad (x^* = 0.567143).$$

- As a fixed point of $g_1(x) = \exp(-x)$.
- As a fixed point of $g_2(x) = -\log(x)$.
- As a fixed point of $g_3(x) := N_f(x) = x - \frac{x - \exp(-x)}{1 + \exp(x)}$

Remark. We just noticed above that the method fails. Observe that $|g_2'(x^*)| > 1$.

Order of convergence

Definition. Assume the above notation. Let $(x_n)_n \rightarrow x^*$ with $g(x^*) = x^*$. Denote by $(\varepsilon_n)_n := x_n - x^*$. We say that the (fixed point) iterative method has **order of convergence** $m > 0$ if

$$\lim_{n \rightarrow \infty} \frac{|\varepsilon_{n+1}|}{|\varepsilon_n|^m} = C > 0.$$

Remark/Exercise.

- If $0 < |g'(x^*)| < 1$ then $m = 1$ (**linear convergence**).
- If $g'(x^*) = \dots = g^{(k-1)}(x^*) = 0$ and $g^{(k)}(x^*) \neq 0$ then $m = k$.
- **Newton's method** $g = N_f$. If $f'(\alpha) \neq 0$ (simple zeros) then $m \geq 2$ (**quadràtic convergence**)