

Efficient Monte-Carlo sampling of metastable systems using non-local collective variable updates

Christoph Schönle,^{1,2} Davide Carbone,² Marylou Gabrié,² Tony Lelièvre,³ and Gabriel Stoltz³

¹⁾ *CMAP, CNRS, École polytechnique, Institut Polytechnique de Paris, 91120 Palaiseau, France*

²⁾ *Laboratoire de Physique de l'École normale supérieure ENS, Université PSL, CNRS, Sorbonne Université, Université de Paris, 75005 Paris, France*

³⁾ *CERMICS, Ecole des Ponts, Institut Polytechnique de Paris, Marne-la-Vallée, France & MATERIALS project-team, Inria Paris, France*

(Dated: December 19, 2025)

Monte-Carlo simulations are widely used to simulate complex molecular systems, but standard approaches suffer from metastability. Lately, the use of non-local proposal updates in a collective-variable (CV) space has been proposed in several works. Here, we generalize these approaches and explicitly spell out an algorithm for non-linear CVs and underdamped Langevin dynamics. We prove reversibility of the resulting scheme and demonstrate its performance on several numerical examples, observing a substantial performance increase compared to methods based on overdamped Langevin dynamics as considered previously. Advances in generative machine-learning-based proposal samplers now enable efficient sampling in CV spaces of intermediate dimensionality (tens to hundreds of variables), and our results extend their applicability toward more realistic molecular systems.

I. INTRODUCTION

Monte-Carlo simulations promise to give access to the equilibrium properties of physical systems even when the numerical simulation of their relaxation dynamics is prohibitively expensive. In practice, however, standard Markov Chain Monte Carlo methods (MCMC) rely on physically-inspired local moves and struggle to equilibrate across metastable states, as does molecular dynamics.

Enhanced sampling algorithms go beyond the naive Markov Chain Monte Carlo approach to tackle metastability, see Ref. 1 for a review in the context of molecular dynamics. A first class of methods originated from the observation that the metastability is removed at sufficiently high temperatures, which led to the development of algorithms like parallel tempering, annealed importance sampling, and sequential Monte Carlo^{2–6}. While known to work on a wide range of problems, they spend a substantial amount of computation time at temperatures not of direct interest and generally require a careful choice of parameters (e.g. for parallel tempering the temperature ladder and the frequency of exchange attempts).

Another important category of algorithms makes use of a collective variable (CV), a lower-dimensional representation of the system that is supposed to capture the metastability of the system. A ‘good’ CV that resolves the metastability can be used for biasing simulations to bridge modes, using enhanced sampling algorithms such as umbrella sampling⁷ and related histogram methods^{8,9}, thermodynamic integration¹⁰ or free energy adaptive biasing methods^{11–14}. However, identifying a suitable CV in low dimension is by no means trivial and thus poses the main limitation to this class of methods.

In recent years, another approach has emerged, which is exploiting generative machine learning models. Among

the various generative models, normalizing flows¹⁵ are particularly well-suited for the purposes of MCMC sampling since their tractable probability density allows to employ them as a non-local proposal density of physical configurations, where any imperfections in training are corrected for via a Metropolis-Hastings accept/reject scheme or self-normalized importance sampling^{16,17}. Thanks to the impressive flexibility of neural networks, proofs of concept have been made in different domains ranging from molecular systems to field theories in quantum chromodynamics^{16,18–21}. However, the limiting factor to scale these approaches appears to be the ability of the generative model to be precise enough either when the dimension of the problem becomes of the order of a few thousands or when the distribution has singularities^{17,22–24}. Other recent approaches make use for example of transformers²⁵ and diffusion models^{26,27}, but these do not give easy access to the underlying probability density and are therefore less suitable for MCMC sampling than normalizing flows.

A promising approach is therefore to combine the strength of CV-enhanced sampling and generative models (normalizing flows) with a CV of ‘intermediate’ dimension (from a dozen to a few hundred degrees of freedom). Within this dimensionality, identifying a CV such that it resolves the metastability will be much easier than finding a one to three-dimensional representation required by most classical CV-based approaches. Conversely, a normalizing flow in an ‘intermediate’ dimension can be trained to much higher accuracy than if one attempted to learn the distribution of the entire system. Since the overall goal is still sampling full configurations of the system, the question arises how to employ such a proposal sampler in CV space to obtain an unbiased sampler for the full distribution. This has been addressed in multiple works^{28–32}, in particular under the names of “Hybrid

Nonequilibrium Molecular Dynamics” (HNMD)^{30,31} and “Nonequilibrium candidate Monte Carlo” (NCMC)²⁹ in the context of molecular dynamics, but also in a Bayesian setting³³. The central idea is always to drive the system from one value of the CV to another, a procedure referred to as steering in the following, and then accept or reject the resulting proposal in the full-space. A similar procedure has also been used in the simulation of glassy system with shear deformations³⁴. In a previous article²³, we described such an algorithm with driven dynamics in continuous time and identified various time-discretizations for which we proved reversibility with respect to the equilibrium measure. Notably, we also related this to the Jarzynski and the Jarzynski–Crooks equalities^{35,36}, clarifying the algorithm design. In a parallel work³⁷, one version of these algorithms was applied to a toy molecular system, using a normalizing flow as the CV sampler and including an adaptive protocol to train it alongside the MCMC procedure.

However, both these works considered only overdamped Langevin dynamics for the steering and required the CV to be a subset of the Euclidean coordinates giving the conformation of the system. In the present article, we generalize these works by presenting an algorithm for a non-linear collective variable and using general underdamped Langevin dynamics. While in spirit this algorithm already fits into the paradigm of some of the previous works like NCMC and HNMD, we present here an algorithm that is readily implemented and treat all technical details arising from the non-linear CV and general Langevin dynamics. We explicitly prove that the associated Markov chains are reversible with respect to the target measure and present its application to several example systems of increasing complexity. In all of them, we see that the deterministic (Hamiltonian) steering dynamics exhibit the best performance, with an improvement of up to two orders of magnitude compared to overdamped dynamics.

The main part of the article divides into two sections: We explain all the theoretical aspects of the problem and the algorithm in Section II and go on to demonstrate its properties through several numerical examples in Section III. Within Section II, the sampling problem is introduced in Section II A, followed by a detailed explanation of the algorithm in Section II B. We state our main theoretical result on the reversibility of the algorithm in Section II C and conclude in Section II D with a normalized version of the algorithm and comments on the parameterization. For the numerical studies in Section III, we present results on four different model systems of increasing complexity in Sections III A to III D. We conclude the article and give an outlook on future directions in Section IV

II. SAMPLING WITH A NON-LINEAR COLLECTIVE VARIABLE

A. Target distribution and collective variable

We begin this section by outlining the sampling problem at hand. Our goal is to sample from the Boltzmann–Gibbs probability measure

$$\nu(dq) = Z_q^{-1} e^{-\beta V(q)} dq \quad (\text{II.1})$$

over positions $q \in \mathbb{R}^d$ of dimension d , with the energy function $V : \mathbb{R}^d \rightarrow \mathbb{R}$, the inverse temperature $\beta = (k_B T)^{-1} > 0$ and the normalizing constant Z_q . Let us assume that a collective variable (CV) $\xi : \mathbb{R}^d \rightarrow \mathbb{R}^\ell$ with $\ell < d$ has been identified and we have a ‘good’ sampler ρ to propose moves in CV space. As mentioned in the introduction, the question of how to use these two ingredients to construct an unbiased MCMC sampler in the full space $\mathbb{R}^d$ has previously been addressed in literature^{28–32}, but it is not obvious how to employ these schemes if the map ξ is non-linear. Notably, the geometry of the level sets $\Sigma(z)$ defined as

$$\Sigma(z) = \left\{ q \in \mathbb{R}^d \mid \xi(q) = z \right\} \quad (\text{II.2})$$

must be taken into consideration. We present an algorithm which explicitly treats the case of a non-linear CV in the following section. Much of the notation throughout this paper is based on Refs. 38 and 39 and we refer the reader to those for an in-depth introduction on the mathematical concepts used here.

Even though we are only interested in sampling $\nu(dq)$, it will turn out to be helpful to introduce also an auxiliary momentum variable p and define the Boltzmann–Gibbs measure in phase space (q, p) by

$$\mu(dq dp) = Z^{-1} e^{-\beta H(q,p)} dq dp \quad (\text{II.3})$$

where Z is the normalizing constant, $H(q, p) = \frac{1}{2} p^\top M^{-1} p + V(q)$ is the Hamiltonian, and $M \in \mathbb{R}^{d \times d}$ is a symmetric positive definite mass matrix.

B. The algorithm

To explain the algorithm, we first present its overall structure in Section II B 1, followed by the necessary ingredients and considerations in Sections II B 2 to II B 4.

1. Overall structure

The algorithm consists in iterating over $n \in \mathbb{N}$ the following steps: Starting from a state Q_n with collective variable value $Z_n := \xi(Q_n)$, generate Q_{n+1} as follows:

Step 1: Propose a move in the collective variable space:

$$\tilde{Z}_{n+1} \sim \rho(Z_n, z') dz',$$

where $\rho(z, z') \in \mathbb{R}^\ell \times \mathbb{R}^\ell \rightarrow \mathbb{R}_+$ denotes a Markov kernel with probability density $\rho(z, z')$ to jump to z' starting at z .

Step 2: Generate a new $\tilde{Q}_{n+1} \in \mathbb{R}^d$ that satisfies $\xi(\tilde{Q}_{n+1}) = \tilde{Z}_{n+1}$ by steering the system in CV space. To this end, (i) specify a schedule to constrain the positions and velocities in CV space, (ii) initialize the position $q^0 = Q_n$ and sample an initial momentum p^0 and (iii) use this to follow underdamped Langevin dynamics in the RATTLE discretization (all of this to be specified below). These dynamics yield a final state $\tilde{Q}_{n+1}$ and a quantity $\mathcal{W}_{n+1}$ that is identified as a work.

Step 3: Draw a random variable U_{n+1} with uniform law over $[0, 1]$ and

- If $U_{n+1} \leq \exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})}$,
accept the proposal: $Q_{n+1} = \tilde{Q}_{n+1}$.
- Otherwise, reject the proposal:
 $Q_{n+1} = Q_n$.

We explain the various ingredients required for **Step 2** in the next three subsections. The idea to sample a system by following some dynamics followed by an accept/reject step that is governed by a work is closely related to the Jarzynski and the Jarzynski–Crooks equalities^{35,36}.

Remark II.1. In all of the numerical examples in this work, we will actually use an independent sampler in the CV space, i.e. $\rho(z, z')$ is only a function of z' .

2. Schedule

To specify the constrained dynamics under which the system evolves from the value of the CV at the previous iteration Z_n to the newly proposed value $\tilde{Z}_{n+1}$, choose first the number of intermediate steps K_T associated with this jump. With the fixed timestep Δt , this sets the physical transition time $T = K_T \Delta t$. In practice, we will choose K_T as a function of the Euclidean distance $\Delta z = |\tilde{Z}_{n+1} - Z_n|$ of the proposed jump.

Introduce a schedule $(z^{(Z_n, \tilde{Z}_{n+1})}(t_k))_{0 \leq k \leq K_T}$ with $t_k = k \Delta t$ whose endpoints are

$$z^{(Z_n, \tilde{Z}_{n+1})}(t_0) = Z_n \quad (\text{II.4})$$

$$z^{(Z_n, \tilde{Z}_{n+1})}(t_{K_T}) = \tilde{Z}_{n+1} \quad (\text{II.5})$$

as well as a velocity schedule $(v_z^{(Z_n, \tilde{Z}_{n+1})}(t_k))_{0 \leq k \leq K_T}$ with zero initial and final velocity,

$$v_z^{(Z_n, \tilde{Z}_{n+1})}(t_0) = v_z^{(Z_n, \tilde{Z}_{n+1})}(t_{K_T}) = 0. \quad (\text{II.6})$$

We require that the schedules for position and velocity satisfy the following reversibility property with the switching of the endpoints: for any $(Z, \tilde{Z})$, we need that

$$z^{(Z, \tilde{Z})}(t_k) = z^{(\tilde{Z}, Z)}(t_{K_T-k}) \quad (\text{II.7})$$

$$v_z^{(Z, \tilde{Z})}(t_k) = -v_z^{(\tilde{Z}, Z)}(t_{K_T-k}), \quad (\text{II.8})$$

for all $k \in \{0, \dots, K_T\}$. Note that the requirement (II.8) precludes the simplest choice for the velocity of $v_z(t_k) = (z(t_{k+1}) - z(t_k))/\Delta t$. A simple choice is instead a schedule that connects the endpoints on a straight line with a continuously differentiable schedule function $f : [0, 1] \rightarrow [0, 1]$:

$$z^{(Z, \tilde{Z})}(t_k) = Z + (\tilde{Z} - Z)f(k/K_T) \quad (\text{II.9})$$

$$v_z^{(Z, \tilde{Z})}(t_k) = (\tilde{Z} - Z) \frac{\dot{f}(k/K_T)}{K_T \Delta t}. \quad (\text{II.10})$$

The conditions on the schedule translate to conditions on f as $f(0) = 0$, $f(1) = 1$ and $\dot{f}(0) = \dot{f}(1) = 0$, which are fulfilled for example by

$$f(\tau) = (1 - \cos(\tau\pi))/2. \quad (\text{II.11})$$

See Theorem B.4 in Section B 2 a for further comments on the velocity schedule. The assumptions (II.6) and (II.8) play a crucial role in the proof of the reversibility of the Markov chain $(Q_n)_{n \geq 0}$ and we demonstrate in Section H 2 how their violation can introduce a bias.

Remark II.2. Even though this is not explicitly indicated, the total time T can also depend on $(Z, \tilde{Z})$ in which case one requires $T^{(Z, \tilde{Z})} = T^{(\tilde{Z}, Z)}$.

3. Initialization of the momentum

For a given position q and momentum p , the velocity in ξ -direction is defined as

$$v_\xi(q, p) = \nabla \xi(q)^\top M^{-1} p \in \mathbb{R}^\ell \quad (\text{II.12})$$

where $\nabla \xi(q) = (\nabla \xi_1(q), \dots, \nabla \xi_\ell(q)) \in \mathbb{R}^{d \times \ell}$. The initial momentum that is sampled in **Step 2** of the algorithm needs to have zero velocity v_ξ to agree with condition (II.6) of the velocity schedule. The cotangent space of zero velocity associated with a position $q \in \Sigma(z)$ is

$$T_q^* \Sigma(z) = \{p \in \mathbb{R}^d \mid v_\xi(q, p) = 0\}, \quad (\text{II.13})$$

and the orthogonal projection on $T_q^* \Sigma(z)$ (for the scalar product induced by M^{-1}) is given by:

$$P_M(q) = \text{Id}_d - \nabla \xi(q) G_M^{-1}(q) \nabla \xi(q)^\top M^{-1}. \quad (\text{II.14})$$

Here, G_M is the Gram tensor defined as

$$G_M(q) = \nabla \xi(q)^\top M^{-1} \nabla \xi(q). \quad (\text{II.15})$$

Throughout this paper, we assume that G_M is invertible on the submanifolds $\Sigma(z)$ (for all z). It is easy to check that, for a given $q \in \Sigma(z)$, one has $P_M(q)p \in T_q^*\Sigma(z)$ for any p and that $P_M(q)^2 = P_M(q)$ and $M^{-1}P_M(q) = P_M(q)^\top M^{-1}$ and thus P_M is indeed the orthogonal projection, as stated above.

We initialize the momentum p^0 by sampling an unconstrained momentum $\tilde{p}^0$ from a centered normal distribution with covariance matrix $\frac{1}{\beta}M$ and projecting it onto the required subspace:

$$p^0 = P_M(q^0)\tilde{p}^0.$$

4. RATTLE dynamics and work

To relax the entire configuration q along the chosen CV and velocity schedules, for reasons related to the geometry of the level sets $\Sigma(z)$ (see also the reversibility proof in Section B 3), we will follow constrained dynamics with a modified potential energy $\tilde{V}(q) = V(q) + V_{\text{fix}}(q)$ with the so-called Fixman term⁴⁰

$$V_{\text{fix}}(q) = \frac{1}{2\beta} \log(\det G_M(q)). \quad (\text{II.16})$$

Starting from (q^0, p^0) , a discretized path $((q^k, p^k)_{0 \leq k \leq K_T})$ is generated by iterating on $0 \leq k \leq K_T - 1$ a version of the RATTLE dynamics with constraints found in literature³⁹ (for the sake of readability, we drop the superscript indicating the endpoints of the schedules $z(t)$ and $v_z(t)$ in this subsection). The scheme writes as follows:

$$p^{k+1/4} = p^k - \frac{\Delta t}{4} \gamma_P(q^k) M^{-1} (p^{k+1/4} + p^k) + \sqrt{\frac{\Delta t}{2}} \sigma_P(q^k) \mathcal{G}^k \quad (\text{II.17})$$

$$\begin{cases} p^{k+1/2} = p^{k+1/4} - \frac{\Delta t}{2} \nabla \tilde{V}(q^k) + \nabla \xi(q^k) \lambda^{k+1/2} \\ q^{k+1} = q^k + \Delta t M^{-1} p^{k+1/2} \\ \xi(q^{k+1}) = z(t_{k+1}) \quad (C_q) \\ p^{k+3/4} = p^{k+1/2} - \frac{\Delta t}{2} \nabla \tilde{V}(q^{k+1}) + \nabla \xi(q^{k+1}) \lambda^{k+3/4} \\ \nabla \xi(q^{k+1})^T M^{-1} p^{k+3/4} = v_z(t_{k+1}) \quad (C_p) \end{cases} \quad (\text{II.18})$$

$$p^{k+1} = p^{k+3/4} - \frac{\Delta t}{4} \gamma_P(q^{k+1}) M^{-1} (p^{k+3/4} + p^{k+1}) + \sqrt{\frac{\Delta t}{2}} \sigma_P(q^{k+1}) \mathcal{G}^{k+1/2} \quad (\text{II.19})$$

where $(\mathcal{G}^k)$ and $(\mathcal{G}^{k+1/2})$ are independent sequences of i.i.d. centered Gaussian random vectors with covariance matrix Id_d . The Lagrange multiplier $\lambda^{k+1/2}$ (resp.

$\lambda^{k+3/4}$) is chosen so that the constraint (C_q) (resp. (C_p)) is satisfied. For $\lambda^{k+3/4}$, this leads to a linear equation with an analytical solution, whereas the solution for $\lambda^{k+1/2}$ will in general be obtained numerically using Newton's methods. For further precisions on this point see Theorem B.3 in the appendix. Note that there is an equivalent formulation of (II.17) and (II.19) that is more suitable for numerical implementation, which we provide in Section A for completeness. The matrices $\gamma_P(q^k) = P_M(q) \gamma P_M(q)^\top$ and $\sigma_P(q^k) = P_M(q) \sigma$ satisfy the fluctuation-dissipation identity

$$\forall q \in \mathbb{R}^d, \quad \sigma_P(q) \sigma_P(q)^\top = \frac{2}{\beta} \gamma_P(q), \quad (\text{II.20})$$

ensured by defining the scalar diffusion parameter σ in terms of the scalar damping parameter $\gamma > 0$ as

$$\sigma^2 = \frac{2\gamma}{\beta}. \quad (\text{II.21})$$

We stick here for simplicity to scalar damping and diffusion parameters, but generalization to matrices is straightforward.

Assuming that the position and velocity schedules are compatible in the continuous-time limit $\Delta t \rightarrow 0$, i.e. $v_z(t) = \dot{z}(t)$, Eqs. (II.17) to (II.19) are discretizations of Langevin dynamics with constraints $(\xi(q^k), v_\xi(q^k, p^k)) = (z(t_k), v_z(t_k))$ see Ref. 39.

These dynamics yield a proposal move in the full configuration space $\tilde{Q}_{n+1} = q^{K_T}$ as well as a work $\mathcal{W}_{n+1} := \mathcal{W}^{K_T}$, which we use to compute the acceptance probability in **Step 3**. This work is defined as

$$\begin{cases} \mathcal{W}^0 = 0 \\ \mathcal{W}^{k+1} = \mathcal{W}^k + \tilde{H}(q^{k+1}, p^{k+3/4}) - \tilde{H}(q^k, p^{k+1/4}), \end{cases} \quad (\text{II.22})$$

for $k = 0 \dots K_T - 1$, where $\tilde{H}$ denotes the Hamiltonian that is modified by the Fixman term (II.16), namely $\tilde{H}(q, p) = \tilde{V}(q) + \frac{1}{2} p^\top M^{-1} p$.

For $\gamma = 0$, the RATTLE dynamics (II.17)-(II.19) become deterministic (driven Hamiltonian dynamics) and the only remaining source of randomness in **Step 2** is the initialization of the momentum p^0 . Furthermore, since in this case $p^{k+1/4} = p^k$ and $p^{k+3/4} = p^{k+1}$, the work simplifies to a telescopic sum and just depends on the endpoints of the path:

$$\mathcal{W}^{K_T} = \tilde{H}(q^{K_T}, p^{K_T}) - \tilde{H}(q^0, p^0).$$

Remark II.3. When the Gram tensor $G_M(q)$ does not depend on q (an example is given in Section III C), the Fixman term just adds an overall shift to the energy V and need not be considered. In this case, the requirement on the initial and final velocity can also be relaxed to $v_z(t_0) = v_z(t_{K_T})$ and can be non-zero, assuming that the initial momentum p^0 is still sampled from the appropriate subspace (see Theorems B.2 and B.10). This is

true in particular for the case of a linear CV, which, after a change of basis, can always be seen as representing a subset of ℓ degrees of freedom. This is the standard setting treated in previous works^{28–32}, where some degrees of freedom are driven through a non-equilibrium protocol.

The problem of constructing a steered path can be understood as transporting samples from the constrained target measure at the initial CV value Z_n to the final one $\tilde{Z}_{n+1}$. Our method is based on a stochastic differential equation (with the auxiliary variable of the momentum), but other dynamics are possible, notably deterministic transformations³⁰ or a combination of stochastic and deterministic steps, where the underlying SDE is modified by a deterministic ‘escorting’ term⁴¹. The latter can also be understood as an instance of a stochastic normalizing flow⁴². These approaches are not easy to adapt to our context of a nonlinear CV, however, since they require to transport samples supported on the submanifolds Σ_z .

C. Reversibility of the algorithm

Our main theoretical result is the following theorem:

Theorem II.4. *The MCMC algorithm presented in Section II B is reversible with respect to the target measure $\nu(dq)$.*

We prove this statement in Section B. The basic ingredient of the proof is to introduce a backward version of the dynamics (associated to the forward dynamics) and use a version of the Jarzynski–Crooks equality³⁶.

As a consequence of Theorem II.4, the MCMC algorithm admits ν as an invariant measure and, up to checking irreducibility which has to be done on a case-by-case basis, empirical averages over trajectories $(Q_n)_{n \geq 0}$ converge to averages with respect to ν .

Remark II.5. In Section H 2, we numerically demonstrate how the omission of the Fixman term or an inappropriate steering schedule can lead to a bias in the simulations, underlining that these two ingredients are indeed necessary for an unbiased algorithm.

D. A normalized version of the algorithm

The overall algorithm depends on many parameters: Δt , γ , σ , M , and the choice of the schedule $z(t)$ and $v_z(t)$ and transition time T . In order to reduce the number of parameters, we will use the following normalized version in the numerical examples: Assuming that the mass matrix $M = M \text{Id}$ is isotropic, we show in Section C that the overall dynamics only depends on the two parameters

$$\alpha_1 = \Delta t \gamma / (4M), \quad \alpha_2 = \Delta t^2 / (\beta M).$$

as well as the schedule $z(t_k)$ and the normalized velocities $\tilde{v}_z(t_k) = \Delta t v_z(t_k)$. To fix these schedules, one can use the

version given by (II.9)-(II.10) with an appropriate function f and choose the number of intermediate steps for a proposed jump as $K_T = \text{ceil}(|\tilde{Z}_{n+1} - Z_n|/\tilde{v})$, with an effective normalized velocity $\tilde{v}$. With this, the algorithm is then fully governed by the three parameters α_1, α_2 and $\tilde{v}$.

For the parameter choice $\alpha_1 = 1$, the steered dynamics (II.17)-(II.19) reduce to overdamped Langevin dynamics (see Section D). If ξ is linear in addition, so that after a suitable change of basis the CV can be identified with the first ℓ degrees of freedom, the overall algorithm is equivalent to the ‘asymmetric’ algorithm presented in our earlier work⁴³ (up to a slight change of the definition of the steered path: here we first move the CV and then the remaining degrees of freedom). Therein, we also provided the continuous-time limit for this version of the algorithm together with the corresponding stochastic differential equation (SDE). For the choice of $\alpha_1 = 0$, the steered dynamics correspond to the Verlet scheme (Leapfrog dynamics) in the orthogonal degrees of freedom and are therefore a discretization of Hamiltonian dynamics with a time-dependent Hamiltonian. Note that in this case, the continuous-time limit of the dynamics $\Delta t \rightarrow 0$ simply corresponds to taking $\alpha_2 \rightarrow 0$ and the acceptance of the move in **Step 3** is governed only by the (normalized) physical transition time $T/\sqrt{\beta M} = K_T \Delta t / \sqrt{\beta M} = \text{ceil}(|\tilde{Z}_{n+1} - Z_n|/\tilde{v}) \sqrt{\alpha_2}$ (see Fig. 14 in the appendix for a numerical illustration).

III. NUMERICAL EXAMPLES

In this section, we apply the proposed CV-guided sampler to examples of increasing complexity, demonstrating in particular the advantage of deterministic underdamped steering dynamics compared to the previously considered overdamped version. We consider four model systems of increasing complexity: a theoretical model-system dubbed ‘Gaussian tunnel’ (Section III A), the one-dimensional ϕ^4 model from statistical physics (Section III B), a dimer in a solvent (Section III C), and finally a 9-bead polymer in a solvent (Section III D).

A. Gaussian tunnel

We first consider a simple model with a linear CV as already introduced in previous work⁴³. It is a toy model in $d = 20$ dimensions with a designated one-dimensional variable z serving as the CV and $d - 1$ additional coordinates $(x_i^\perp)_{1 \leq i \leq d-1}$. The distribution of z is given by a Gaussian mixture and the conditional distribution of the remaining coordinates is a Gaussian with a mean de-

pending on z . More precisely,

$$\begin{aligned}\nu(dz, dx^\perp) &= \nu_{\text{CV}}(dz) \nu_\perp(dx^\perp|z), \\ \nu_{\text{CV}}(dz) &= \frac{1}{\sqrt{2\pi}} \left(w e^{-z^2/2} + (1-w) e^{-(z-b)^2/2} \right) dz, \\ \nu_\perp(dx^\perp|z) &= (2\pi)^{-(d-1)/2} \det(\Sigma)^{-1/2} \\ &\times e^{\left(-\frac{1}{2}(x^\perp - \mu(z))^\top \Sigma^{-1}(x^\perp - \mu(z))\right)} dx^\perp.\end{aligned}\tag{III.1}$$

$$\tag{III.2}$$

Here, $\mu(z) = \frac{b}{2} \cos(z\pi/b)(1, 1, \dots, 1)^\top$ is the conditional mean and Σ a diagonal covariance matrix with $\Sigma_{ii} = \sigma_i^2$. The weight of the first mode is $w = 0.3$, while the distance between the modes is $b = 10$. The values of σ_i were chosen from an evenly spaced grid in the interval $[0.5, 5]$.

To apply the algorithm from Section II B, we use a Gaussian mixture model as the proposal sampler in CV space in **Step 1** which is the same as the target ν_{CV} apart from a different mode weight $w = 0.5$. We identify $V(q) = V(z, x) = -\log \nu(z, x)$ with $\beta = 1$, where, with a slight abuse of notation, we still denote by ν the density of the measure $\nu(dq)$. We use the normalized version of the algorithm and schedule that only depends on α_1 , α_2 and $\tilde{v}$, detailed in Section II D. Since no Fixman term needs to be considered here, we simply use a linear schedule function $f(\tau) = \tau$.

To evaluate the performance, we consider (i) the acceptance rate for a fixed jump across the metastable basins from $z = 0$ to $z = b$ normalized by the number of intermediate steps K_T (as a measure of computational cost) and (ii) the inverse mode jump cost, which is the number of accumulated steps (calls to the force) that is needed on average to see a mode switch between $z < b/2$ and $z > b/2$ when running the full algorithm.

Running a rough optimization through a grid search reveals that the optimum performance is obtained for $\alpha_1 = 0$, which corresponds to a deterministic steered dynamics (see Fig. 13 of Section H1). We compare the deterministic ($\alpha_1 = 0$) and the overdamped ($\alpha_1 = 1$) steered dynamics in Fig. 1 using the two performance measures, which show consistent results. The deterministic dynamics offer a large performance increase over the overdamped dynamics with a difference of around two orders of magnitude. In both cases, we observed that for too large values of α_2 (a too coarse time discretization), the dynamics become unstable and the trajectories diverge, leading to zero acceptance of the overall move. The optimal value for α_2 is then just below this threshold. Typical driven transition paths are shown for the optimum of the overdamped and the deterministic case in Fig. 2. To verify the unbiasedness of the algorithm at the optimum, we show the marginals of z and $x_0^\perp$ in Fig. 3. They indeed agree with the target measure and show that the algorithm corrects for the biased proposal distribution in CV space.

Since the Fixman term and condition Eq. (II.6) on the schedule need not be considered in this simple case of a




Figure 1. Performance of the algorithm for the Gaussian tunnel example in the special cases of deterministic ($\alpha_1 = 0$) and overdamped Langevin dynamics ($\alpha_1 = 1$). The top row shows the acceptance rate normalized by K_T for a fixed jump from $z = 0$ to $\tilde{z} = b$ for different values of K_T , averaged over 10,000 random initializations of $x^\perp \sim \nu_\perp(x^\perp|z)$. The second row shows the inverse mode jump cost, averaged over 20 chains and 10,000 iterations, as a function of the number of steps $K_T(b)$ for a jump of distance b , which is related to the inverse of the dimensionless velocity $\tilde{v}$ via $K_T(b) = \text{ceil}(b/\tilde{v})$. White space indicates that at least one chain never switched mode.




Figure 2. Typical driven transition path from one mode center to another, $z = 0$ to $z = b = 10$, for overdamped ($\alpha_1 = 1$) and deterministic ($\alpha_1 = 0$) dynamics for the optimal α_2 and $K_T = b/\tilde{v}$.

linear CV, we also show results for a modified version with a non-linear CV in Section H2. In this case, we indeed observe that a bias emerges when these ingredients are disregarded.




Figure 3. Gaussian Tunnel marginals for the optimum performance values $\alpha_1 = 0$, $\alpha_2 = 0.67$ and $\tilde{v} = 0.2$, corresponding to $K_T(b) = 50$. Results are obtained by running 8 independent chains for 200,000 iterations.

B. ϕ^4 Model

As a second example we consider a field model from statistical physics, the one-dimensional ϕ^4 model (also considered previously⁴³). It consists of N degrees of freedom $\{\phi_i\}_{i=1}^N$ on a one-dimensional lattice and can be seen as a continuous version of the Ising model for ferromagnetism in solids. The target distribution is the Boltzmann–Gibbs probability distribution $\nu(d\phi)$ with the energy (using the simplified version of the model considered in Ref. 19)

$$\beta V(\phi) = \frac{a\beta N}{2} \sum_{i=1}^{N+1} (\phi_i - \phi_{i-1})^2 + \frac{\beta}{4aN} \sum_{i=1}^N (1 - \phi_i^2)^2, \quad (\text{III.3})$$

with Dirichlet boundary conditions $\phi_0 = \phi_{N+1} = 0$. For sufficiently low temperatures $1/\beta$, this system has two well-separated modes characterized by positive and negative magnetization $\bar{\phi} = \sum_{i=1}^N \phi_i / N$ concentrated at $\pm \bar{\phi}^*$ for some $\bar{\phi}^*$ which depends on β . We place ourselves in this bimodal phase by setting $\beta = 20$ and $a = 0.1$. In many practical settings, one will be interested in systems with an external field h , which corresponds to adding a term $h \sum_{i=1}^N \phi_i$ to the potential $V(\phi)$. For the purpose of testing the algorithm, however, the case $h = 0$ is most illustrative since the two modes will have exactly the same weight due to the symmetry $V(\phi) = V(-\phi)$, and one can verify that the algorithm produces no bias with respect to the mode weights.

We use the magnetization $\bar{\phi}$ as the collective variable, which is a natural order parameter of the system. We run the algorithm on the system in a wavelet basis (see Section E for details) since this order parameter then naturally emerges as the first component of the system. To fix the proposal sampler in CV space, we first ran a short simulation using the Metropolis Adjusted Langevin Algorithm (MALA)^{44,45}, which, due to the local proposal updates of MALA, always stays within one mode. From this data, the mean and standard deviation of the observed

mode were then estimated as $\bar{\phi}^* = 0.79$ and $\sigma = 0.06$. Consequently, we used as the proposal sampler a Gaussian mixture model with two modes centered at $\pm \bar{\phi}^*$ and of width σ .

Since we are again using a linear CV here, we use the same linear positions schedule $z(t_k)$ and constant velocity schedule $v_z(t_k)$ which we also used for the Gaussian tunnel. We compare the performance of the sampling algorithm over a grid of parameters α_1 , α_2 and $\tilde{v}$, using the mode jump cost as a criterion (the average number of calls to the force between transitions between $\bar{\phi} > 0$ and $\bar{\phi} < 0$). Optimal performance is again observed for the deterministic dynamics at $\alpha_2 = 0$ (see Fig. 16 in Section H3). Notably, as can be seen in Fig. 4, there is a performance increase by around two orders of magnitude passing from the overdamped to the deterministic dynamics. This is consistent with the experiments for the Gaussian tunnel. Away from the optimum, we see a white diagonal where no transitions were observed, corresponding to a particularly ‘bad’ physical transition time $T/\sqrt{\beta M} = K_T \sqrt{\alpha_2}$. This might be a resonance effect due to the deterministic dynamics, but it is in any case far away from the parameter range of optimum performance.

For the optimal parameters, we also run a modified experiment where we deliberately bias the CV proposal sampler by putting a larger weight on the mode corresponding to negative magnetization. As we show in Fig. 5, this bias is accurately corrected and the obtained samples align perfectly with the target density.

Driving the system from one mode to another corresponds to creating a domain of opposite orientation from either end of the lattice and gradually increasing its size. We indeed observe in Fig. 6 that ‘good’ transitions with a high probability of acceptance follow one of these two transition paths. The existence of multiple transition paths does not pose a difficulty per se for the algorithm presented here: the acceptance of a proposal move is only determined by the work accumulated along the way and thus the intermediate bimodality is not a problem for accurate sampling of the two modes and their relative weights.

C. Dimer in a solvent

We now apply the algorithm on a model with a nonlinear collective variable^{38,46,47}. It consists of N particles of equal masses M in a two-dimensional box of side length L with periodic boundary conditions. Two of the particles form a dimer; the remaining particles are designated as solvent particles. The solvent particles interact between them and with the dimer particles via the Weeks–Chandler–Andersen (WCA) pair potential, which




Figure 4. Performance of the algorithm applied to the ϕ^4 model, estimated from 10 chains over 20,000 iterations. The x-axis represents the number of steps for a steered trajectory of over the distance $2\bar{\phi}^*$ between the two modes. White space indicates that at least one chain never switched mode.




Figure 5. Marginal distribution of the ϕ^4 model obtained from running 8 chains for 10^4 steps with $\alpha_1 = 0$, $\alpha_2 = 0.0014$ and $\tilde{v} = 1.2 \times 10^{-3}$ (the optimal parameters obtained from optimization in Fig. 4). In this case, a biased GMM proposal with weights 0.3 and 0.7 was used.

is a truncated Lennard-Jones potential:

$$V_{\text{WCA}}(r) = \begin{cases} 4\epsilon \left[\left(\frac{r^*}{r} \right)^{12} - \left(\frac{r^*}{r} \right)^6 \right] + \epsilon & \text{if } r \leq r_0, \\ 0 & \text{if } r > r_0. \end{cases}$$




Figure 6. Two possible transition paths in ϕ^4 model from positive to negative values of $\bar{\phi}$ with the same parameters as in Fig. 5




Figure 7. Typical configurations for the dimer in the compact and the stretched state.

Here, r is the distance between two particles, $r_0 = 2^{1/6}r^*$ and ϵ and r^* are positive constants. The two particles of the dimer interact via a double-well potential

$$V_D(r) = h \left[1 - \frac{(r - r_0 - w)^2}{w^2} \right]^2, \quad (\text{III.4})$$

with the two positive parameters h and w . Denoting the particle positions by $\{q_i\}_{i=1}^N$ with each $q_i \in \mathbb{R}^2$, the total potential energy is thus given by

$$V(q) = V_D(|q_1 - q_2|) + \sum_{3 \leq i < j \leq N} V_{\text{WCA}}(|q_i - q_j|) + \sum_{i=1,2} \sum_{3 \leq j \leq N} V_{\text{WCA}}(|q_i - q_j|). \quad (\text{III.5})$$

The minima of V_D are at $r = r_0$ and $r = r_0 + 2w$, giving rise to a compact state and stretched state with an energy barrier of height h . Two configurations for the compact and the stretched state are shown in Fig. 7.

As a reaction coordinate, we use the normalized bond length

$$\xi(q) = \frac{|q_1 - q_2| - r_0}{2w}, \quad (\text{III.6})$$

so that the compact and the stretched states correspond to $\xi(q) = 0$ and $\xi(q) = 1$, respectively. As explained in Section F 1, the Gram tensor is given by $G_M(q) = (2w^2M)^{-1}$. Since it does not depend on q , the Fixman term

$$V_{\text{fix}}(q) = -\frac{1}{2\beta} \log(2w^2M)$$

is therefore a constant that does not change the dynamics and need not be considered. The Lagrange multipliers for the constrained dynamics can be obtained analytically for this system and CV (see Section F 2). We make use of this since it is numerically beneficial, but in principle, one could equally use numerical solvers as explained in Section II B 4.

We ran numerical experiments for parameters $h = w = 2$, $r^* = \epsilon = 1$ and $M = 1$ with $N = 100$ particles and

a box of length $L = 15$, in line with previous works^{38,39}. For the CV sampler we use a Gaussian mixture model with modes of equal weight at $z = 0$ and $z = 1$ and width $\sigma_{\text{prop}} = 0.2$. We reject any unphysical proposals with $z < z_{\text{min}} = -\frac{r_0}{2w}$ and $z > z_{\text{max}} = (L/\sqrt{2} - r_0)/(2w)$ since the distance $|q_1 - q_2|$ is always positive and the periodic boundary conditions (torus coordinates) impose a maximum possible distance.

We compare the performance of the algorithm for different parameter values using a mode jump cost that counts the number of calls to the force required to go from a state with $z < 0.1$ to a state with $z > 0.9$ or vice versa, in line with the ‘mean residence duration’ used in Ref. 48.

Once again, we observe a large advantage in using the deterministic algorithm with $\alpha_1 = 0$, whose performance for different values of α_2 and $\tilde{v}$ are shown in Fig. 8, while we see almost no mode switches for the overdamped algorithm within the same computational budget (see Section H 4).

In Fig. 9, we show the free energy profile obtained by running the algorithm for the optimized parameters until convergence together with the free energy in the absence of solvent particles (see Section F 3 for its derivation). As one can see, the presence of the solvent particles increases the likelihood of the compact state (lowering its free energy in comparison with the stretched state). The result agrees almost perfectly with a result obtained from thermodynamic integration (TI) (see Ref. 39 for a theoretical introduction). In thermodynamic integration, the system is sampled under the constraint $\xi(q) = z_i$ for a sequential series of CV-values z_i on a grid to estimate the mean force. This demonstrates that the inaccuracy of the proposal potential has been corrected by the algorithm.

Note that this model was also considered by the authors of Ref. 29. Contrary to our approach, however, they chose the full dimer configuration as the CV, $\xi(q) = (q_1, q_2)^\top$, and used a proposal kernel that samples a dimer configuration $\tilde{Z}_{n+1}$ conditioned on the current configuration Z_n with a new distance $|q_1 - q_2|$ but the same overall rotation angle.

D. Polymer in a solvent

As a final example, we consider a toy molecular system that consists of a 9-bead polymer embedded in a three-dimensional box with 200 solvent particles, similar to a model introduced in Ref. 37. The polymer beads and the solvent particles all interact via a Lennard-Jones (LJ) potential. Within the polymer, the nearest and second-nearest neighbor distances (and hence the binding angle) of the polymer beads are softly constrained with a harmonic potential. In addition, a bi-modal potential is imposed on the end-to-end distance of the polymer, giving rise to two stable confirmations in an open and a closed state, see Fig. 10. Denoting by $\{q_i\}_{i=1,\dots,9}$ the polymer coordinates and by $\{q_i\}_{i=10,\dots,N}$ the solvent coordinates,




Figure 8. Performance comparison for the dimer model introduced in Section III C for the deterministic case $\alpha_1 = 0$, each experiment with 20 chains being run for 5,000 iterations. The optimal performance is observed at $\alpha_2 = 4.52 \times 10^{-4}$ and $1/\tilde{v} = 378$. White areas indicate that at least one chain never switched mode.




Figure 9. Free energy profile for the dimer model obtained from running the CV guided algorithm with the optimal parameters determined in Fig. 8 with 20 chains and 50,000 iterations, shown alongside results from thermodynamic integration on a grid of 50 values and 10^6 steps each. For comparison, we also show the free energy profile of the dimer without solvent particles, see Section F 3, and the energy of the proposal distribution. An energy shift is applied to all of these curves such that they coincide at $z = 1$. Error bars are estimated from the standard deviation over parallel MCMC chains of counts of the histogram bins.




Figure 10. Open and closed state of the polymer (red) and the solvent particles (blue), characterized by the end-to-end distance




Figure 11. Simulation of the polymer with a one-dimensional CV (end-to-end distance). Left panel: Rough grid optimization of parameters for the deterministic case $\alpha_1 = 0$, each simulation run with 20 parallel chains and 2,000 iterations. Right panel: Marginals for a run of 20 chains and 100,000 iterations for the optimized parameters $1/\bar{v} = 180$, $\alpha_2 = 2^{-4}$ (overall 8% acceptance), shown alongside the CV proposal distribution and reference samples obtained from a biased simulation that has been reweighted.

the total energy is given by

$$\begin{aligned}
 V(q) = & \sum_{i=1}^8 V_1(|q_i - q_{i+1}|) + \sum_{i=1}^7 V_2(|q_i - q_{i+2}|) \\
 & + \sum_{\substack{1 \leq i \leq 6 \\ i+3 \leq j \leq 9}} V_{\text{LJ}}(|q_i - q_j|) + V_e(|q_1 - q_9|) \\
 & + \sum_{\substack{1 \leq i \leq 9 \\ 10 \leq j \leq N}} V_{\text{LJ}}(|q_i - q_j|) \\
 & + \sum_{10 \leq i < j \leq N} V_{\text{LJ}}(|q_i - q_j|), \quad (\text{III.7})
 \end{aligned}$$

with a Lennard-Jones interaction V_{LJ} , harmonic potentials $V_\kappa(r) = \frac{k_\kappa}{2}(r - d_\kappa)^2$ for $\kappa \in \{1, 2\}$ and a double-well potential (acting on the end-to-end distance) $V_e(r) = \frac{k_e}{4} \left(\frac{r-d_e}{a} \right)^4 - k_e \left(\frac{r-d_e}{a} \right)^2$ with two local minima at $r = d_e \pm a\sqrt{2}$. Precise parameter values and implementation details are provided in Section G.

We test the algorithm for two different choices of CV: (i) the end-to-end distance, $\xi(q) = |q_1 - q_9|$, a




Figure 12. Simulation of the polymer with the entire polymer as the CV. Left panel: Rough grid optimization of parameters for the deterministic case $\alpha_1 = 0$, each simulation run with 20 parallel chains and 2,000 iterations. Right panel: Marginals for a run of 20 chains and 10,000 iterations for the optimized parameters $1/\bar{v} = 60$, $\alpha_2 = 5 \times 10^{-4}$ (99% acceptance rate), shown alongside the marginalized CV proposal distribution and reference samples obtained from a biased simulation that has been reweighted.

one-dimensional CV, and (ii) the entire polymer chain, $\xi(q) = (q_1, \dots, q_9)$, a 27-dimensional CV. For the latter, we define a proposal sampler with the help of a normalizing flow.

When running simple MALA dynamics on this system, mode transitions are only observed very rarely, necessitating the use of an enhanced sampling technique. To obtain a reference dataset to compare against, we ran biased dynamics (removing the potential energy V_e on the end-to-end distance) and unbiased them via reweighting.

An obvious choice of CV that resolves the bimodality is the end-to-end distance. Since it is one-dimensional, it is easy to construct a proposal distribution in the form of a Gaussian mixture. For simplicity, we extract the position of the modes and their width from short Langevin simulations on the full system (see Section G). A rough grid search to optimize the parameters of the algorithm is shown in Fig. 11, with the best achieved inverse mode jump cost at around 9×10^{-5} . In contrast, the overdamped algorithm with $\alpha_1 = 1$ does not show any transitions within the same parameter range and computational budget, giving an upper limit of the performance of at most 6×10^{-7} ; it is thus at least two orders of magnitude worse in performance.

Another possible choice of collective variable in this case is to use the positions of the beads of the entire polymer configuration. Here, a good proposal distribution requires a careful parameterization, for which we use a normalizing flow (see Section G). Since our primary interest is in the performance of the algorithm, we trained our proposal sampler on already existing reference data. In practice, one could use adaptive schemes in which the CV-sampler is progressively trained while running the MCMC sampler^{19,37}. Note that for this choice of CV, the steered dynamics in **Step 2** of the algorithm effectively only acts on the solvent particles, since the schedule al-

ready imposes the evolution of the entire polymer. Running the algorithm, we again observe that the deterministic dynamics ($\alpha_1 = 0$) gives good results with frequent mode switches while the overdamped version ($\alpha_1 = 1$) does not work at all. Strikingly, for the optimized parameters, we achieve an overall acceptance rate of 99% with this algorithm. Since the proposals are drawn independently, this corresponds almost to the optimum performance with maximum decorrelation within the polymer degrees of freedom. A rough grid search for the deterministic algorithm is shown in Fig. 12 together with the marginal distribution of the end-to-end distance. Comparing the optimal mode jump cost obtained with the two choices of CV, we see an advantage of around one order of magnitude in using the positions of all polymer beads over the end-to-end distance between first and last bead as a CV. This only accounts for the calls to the force, however, and not for the effort in learning the proposal sampler in CV space.

IV. CONCLUSION AND OUTLOOK

In this work, we present an MCMC algorithm with non-local proposals in a collective variable space. This CV space may be high-dimensional, provided a good proposal sampler can still be learned. We generalize related algorithms by considering the underdamped Langevin dynamics and a non-linear collective variable and we prove reversibility of the resulting scheme. We provide a detailed description of the algorithm so that it is readily implementable and we illustrate its performance on a number of examples, ranging from a toy low-dimensional example to molecular systems. We demonstrate a significant performance advantage in using Hamiltonian over overdamped steered Langevin dynamics.

We leave certain questions for future investigations, notably the performance of the algorithm with respect to different choices of CV, as well as the choice of schedule (path) in CV space and the training of the proposal sampler. Furthermore, it would be very interesting to apply this algorithm to more complex molecular systems such as the conformational equilibrium of biomolecules, which is something we leave for future work.

AUTHOR DECLARATIONS

Conflict of interest

The authors have no conflicts to disclose.

DATA AVAILABILITY

A running implementation of the algorithms and all code to reproduce results presented in this work are publicly available at

https://github.com/cschoenle/mcmc_nonlinvcv. Simulation data can be reproduced at reasonable computational cost.

ACKNOWLEDGMENTS

The authors thank warmly Alessandro Laio and Samuel Tamagnone for their help in setting up the last numerical example of the polymer. The authors remember fondly Samuel. This work was supported by Hi! Paris, a French government grant managed by the Agence Nationale de la Recherche under the France 2030 program, reference ANR-23-IACL-0005, and supported as well by PR[AI]RIE-PSAI, also a French government grant managed by the Agence Nationale de la Recherche under the France 2030 program, reference ANR-23-IACL-0008. The works of T.L. and G.S. are partially funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (project EMC2, grant agreement No 810367), and from the Agence Nationale de la Recherche through the grants ANR-19-CE40-0010-01 (QuAMProcs) and ANR-21-CE40-0006 (SINEQ). This work was partially performed using HPC resources from GENCI-IDRIS (AD011015234R1).

REFERENCES

- ¹J. Hénin, T. Lelièvre, M. Shirts, O. Valsson, and L. Delemotte, *Living Journal of Computational Molecular Science* **4**, 1583 (2022).
- ²R. Swendsen and J.-S. Wang, *Phys. Rev. Lett.* **57**, 2607 (1986).
- ³R. Neal, *Statistics and Computing* **11**, 125 (2001).
- ⁴P. Del Moral, A. Doucet, and A. Jasra, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **68**, 411 (2006).
- ⁵D. B. Woodard, S. C. Schmidler, and M. Huber, *The Annals of Applied Probability* **19**, 617 (2009).
- ⁶S. Syed, A. Bouchard-Côté, G. Deligiannidis, and A. Doucet, *Journal of the Royal Statistical Society Series B: Statistical Methodology* **84**, 321 (2022).
- ⁷G. Torrie and J. Valleau, *J. Comp. Phys.* **23**, 187 (1977).
- ⁸S. Kumar, D. Bouzida, R. H. Swendsen, P. A. Kollman, and J. M. Rosenberg, *J. Comput. Chem.* **13**, 1011 (1992).
- ⁹M. R. Shirts and J. D. Chodera, *J. Chem. Phys.* **124**, 124105 (2008).
- ¹⁰J. Kirkwood, *J. Chem. Phys.* **3**, 300 (1935).
- ¹¹E. Darve and A. Porohille, *J. Chem. Phys.* **115**, 9169 (2001).
- ¹²J. Hénin and C. Chipot, *J. Chem. Phys.* **121**, 2904 (2004).
- ¹³A. Laio and M. Parrinello, *PNAS* **99**, 12562 (2002).
- ¹⁴A. Barducci, G. Bussi, and M. Parrinello, *Phys. Rev. Lett.* **100**, 020603 (2008).
- ¹⁵G. Papamakarios, E. Nalisnick, D. J. Rezende, S. Mohamed, and B. Lakshminarayanan, *J. Mach. Learn. Res.* **22**, 1 (2021).
- ¹⁶F. Noé, S. Olsson, J. Köhler, and H. Wu, *Science* **365**, eaaw1147 (2019).
- ¹⁷L. Grenioux, A. Durmus, E. Moulines, and M. Gabrié, in *Proceedings of the 40th International Conference on Machine Learning* (PMLR, 2023) pp. 11698–11733.
- ¹⁸S.-H. Li and L. Wang, *Phys. Rev. Lett.* **121**, 260601 (2018).
- ¹⁹M. Gabrié, G. Rotskoff, and E. Vanden-Eijnden, *Proc. Natl. Acad. Sci. U.S.A* **119**, e2109420119 (2022).

- ²⁰M. Invernizzi, A. Krämer, C. Clementi, and F. Noé, *J. Phys. Chem. Lett.* **13**, 11643 (2022).
- ²¹R. Abbott, M. S. Albergo, A. Botev, D. Boyda, K. Cranmer, D. C. Hackett, G. Kanwar, A. G. D. G. Matthews, S. Racanière, A. Razavi, D. J. Rezende, F. Romero-López, P. E. Shanahan, and J. M. Urban, arXiv:2208.03832 (2022).
- ²²L. Del Debbio, J. Rossney, and M. Wilson, *Phys. Rev. D* **104**, 094507 (2021).
- ²³C. Schönle and M. Gabrié, in *NeurIPS 2023 AI for Science Workshop* (2023).
- ²⁴R. Abbott, M. S. Albergo, A. Botev, D. Boyda, K. Cranmer, D. C. Hackett, A. G. D. G. Matthews, S. Racanière, A. Razavi, D. J. Rezende, F. Romero-López, P. E. Shanahan, and J. M. Urban, arXiv:2211.07541 (2022).
- ²⁵P. Bera and J. Mondal, *Chem. Sci.* **16**, 8735 (2025).
- ²⁶S. Lewis, T. Hempel, J. Jiménez-Luna, M. Gastegger, Y. Xie, A. Y. K. Foong, V. G. Satorras, O. Abdin, B. S. Veeling, I. Zaporozhets, Y. Chen, S. Yang, A. E. Foster, A. Schneuing, J. Nigam, F. Barbero, V. Stimper, A. Campbell, J. Yim, M. Lienen, Y. Shi, S. Zheng, H. Schulz, U. Munir, R. Sordillo, R. Tomioka, C. Clementi, and F. Noé, *Science* **389**, eadv9817 (2025).
- ²⁷P. Bera and J. Mondal, *J. Chem. Phys.* **163**, 114110 (2025).
- ²⁸M. Athènes, *Phys. Rev. E* **66**, 046705 (2002).
- ²⁹J. Nilmeier, G. Crooks, D. Minh, and J. Chodera, *PNAS* **108**, E1009 (2011).
- ³⁰Y. Chen and B. Roux, *J. Chem. Theory Comput.* **11**, 3572 (2015).
- ³¹Y. Chen and B. Roux, *J. Chem. Phys.* **142**, 024101 (2015).
- ³²R. Neal, arXiv:math/0502099 (2005).
- ³³G. Karagiannis and C. Andrieu, *J. Comp. Graph. Stat.* **22**, 623 (2013).
- ³⁴P. Das, A. D. S. Parmar, and S. Sastry, *J. Chem. Phys.* **157**, 044501 (2022).
- ³⁵C. Jarzynski, *Phys. Rev. E* **56**, 5018 (1997).
- ³⁶G. Crooks, *Phys. Rev. E* **60**, 2721 (1999).
- ³⁷S. Tamagnone, A. Laio, and M. Gabrié, *J. Chem. Theory Comput.* **20**, 7796 (2024).
- ³⁸T. Lelièvre, M. Rousset, and G. Stoltz, *Free Energy Computations: A Mathematical Perspective* (Imperial College Press, 2010).
- ³⁹T. Lelièvre, M. Rousset, and G. Stoltz, *Math. Comp.* **81**, 2071 (2012).
- ⁴⁰M. Fixman, *J. Chem. Phys.* **69**, 1527 (1978).
- ⁴¹S. Vaikuntanathan and C. Jarzynski, *The Journal of Chemical Physics* **134**, 054107 (2011).
- ⁴²M. Caselle, E. Cellini, A. Nada, and M. Panero, *J. High Energ. Phys.* **2022**, 15 (2022), arXiv:2201.08862 [hep-lat].
- ⁴³C. Schönle, M. Gabrié, T. Lelièvre, and G. Stoltz, *J. Comp. Phys* **527**, 113806 (2025).
- ⁴⁴P. Rossky, J. Doll, and H. Friedman, *J. Chem. Phys.* **69**, 4628 (1978).
- ⁴⁵G. Roberts and R. Tweedie, *Bernoulli* **2**, 341 (1996).
- ⁴⁶J. E. Straub, M. Borkovec, and B. J. Berne, *J. Chem. Phys.* **89**, 4833 (1988).
- ⁴⁷C. Dellago, P. G. Bolhuis, and D. Chandler, *J. Chem. Phys.* **110**, 6617 (1999).
- ⁴⁸T. Lelièvre, M. Rousset, and G. Stoltz, *Numer. Math.* **143**, 379 (2019).
- ⁴⁹D. McDuff and D. Salamon, *Introduction to Symplectic Topology* (Oxford University Press, 2017).
- ⁵⁰E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*, Springer Series in Computational Mathematics, Vol. 31 (Springer-Verlag, 2006).
- ⁵¹B. J. Leimkuhler and R. D. Skeel, *Journal of Computational Physics* **112**, 117 (1994).
- ⁵²C. Chui, *An Introduction to Wavelets, Wavelet Analysis and its Applications*, Vol. 1 (Academic Press, Inc., Boston, MA, 1992).
- ⁵³S. Mallat, *A Wavelet Tour of Signal Processing: The Sparse Way* (Academic Press, 2008).
- ⁵⁴S. S. Schoenholz and E. D. Cubuk, in *Advances in Neural Information Processing Systems*, Vol. 33 (Curran Associates, Inc., 2020).
- ⁵⁵D. Ward, “FlowJAX: Distributions and normalizing flows in JAX,” (2025), v17.2.0.
- ⁵⁶C. Durkan, A. Bekasov, I. Murray, and G. Papamakarios, arXiv:1906.04032 (2019).

Appendix A: Numerical implementation of the evolution of momenta

For practical purposes, one can reformulate (II.17) (and (II.19) in the same fashion) in the following equivalent form that is more suitable to numerical implementation, see Sec. 5.4.2 in Ref. 39 for further explanations:

$$\begin{cases} p^{k+1/4} = p^k - \frac{\Delta t}{4} \gamma M^{-1} \left(p^k + p^{k+1/4} - 2 \nabla \xi G_M^{-1}(q^k) v_z(t_k) \right) \\ \quad + \sqrt{\frac{\Delta t}{2}} \sigma \mathcal{G}^k + \nabla \xi(q^k) \lambda^{k+1/4}, \\ \nabla \xi(q^k)^\top M^{-1} p^{k+1/4} = v_z(t_k), \end{cases} \quad (C_p) \quad (\text{A.1})$$

where the Lagrange multiplier $\lambda^{k+1/4}$ is associated with the constraint (C_p) . It can be determined analytically by solving

$$v_z(t_k) = \nabla \xi(q^k)^\top M^{-1} \left(\text{Id} + \frac{\Delta t}{4} \gamma M^{-1} \right)^{-1} \left(\left(\text{Id} - \frac{\Delta t}{4} \gamma M^{-1} \right) p^k + \frac{\Delta t}{2} \gamma M^{-1} \nabla \xi G_M^{-1}(q^k) v_z(t_k) + \sqrt{\frac{\Delta t}{2}} \sigma \mathcal{G}^k + \nabla \xi(q^k) \lambda^{k+1/4} \right),$$

assuming that $\nabla \xi(q^k)^\top M^{-1} \left(\text{Id} + \frac{\Delta t}{4} \gamma M^{-1} \right)^{-1} \nabla \xi(q^k)$ is invertible.

Appendix B: Proof of the reversibility of the algorithm

The objective of this section is to prove the reversibility of the algorithm introduced in Section II B (namely Theorem II.4). This is done in three steps. In Section B 1, we introduce the necessary material, in particular concerning measures on submanifolds associated with the collective variable ξ . Then, in Section B 2, we present fundamental properties of the steered dynamics (II.17)-(II.19), and in particular the Jazynski-Crooks equality. This equality is the cornerstone of the proof of reversibility which is finally provided in Section B 3.

1. Introducing notation

This subsection largely uses tools and results from Ref. 39. Let us introduce a number of quantities which we will need later on. A central quantity of interest is the free energy $F : \mathbb{R}^\ell \rightarrow \mathbb{R}$ associated with the target measure ν and the collective variable ξ , defined as (see (II.2) for the definition of $\Sigma(z)$)

$$F(z) = -\frac{1}{\beta} \log \int_{\Sigma(z)} Z_q^{-1} e^{-\beta V(q)} \delta_{\xi(q)-z}(\text{d}q). \quad (\text{B.1})$$

The phase space associated with constraints on the collective variable ξ and effective velocity v_ξ is denoted as

$$\Sigma_{\xi, v_\xi}(z, v_z) = \left\{ (q, p) \in \mathbb{R}^{2d} \mid \xi(q) = z, v_\xi(q, p) = v_z \right\}. \quad (\text{B.2})$$

For a fixed position q , we denote the affine space of constrained momenta as

$$\Sigma_{v_\xi(q, \cdot)}(v_z) = \{ p \in \mathbb{R}^d \mid v_\xi(q, p) = v_z \}. \quad (\text{B.3})$$

Note that $T_q^* \Sigma(z) = \Sigma_{v_\xi(q, \cdot)}(0)$, which was defined in (II.13). The Boltzmann–Gibbs probability distribution can also be defined on these constrained spaces as follows:

$$\begin{cases} \mu_{\Sigma_{\xi, v_\xi}(z, v_z)}(\text{d}q \text{d}p) := \frac{e^{-\beta H(q, p)}}{\mathcal{Z}_{z, v_z}} \sigma_{\Sigma_{\xi, v_\xi}(z, v_z)}(\text{d}q \text{d}p), \\ \mathcal{Z}_{z, v_z} := \int_{\Sigma_{\xi, v_\xi}(z, v_z)} e^{-\beta H} \text{d}\sigma_{\Sigma_{\xi, v_\xi}(z, v_z)}, \end{cases} \quad (\text{B.4})$$

where $\sigma_{\Sigma_{\xi, v_\xi}(z, v_z)}(\text{d}q \text{d}p)$ is the phase space Liouville measure on $\Sigma_{\xi, v_\xi}(z, v_z)$.

Using the Fixman term V_{fix} introduced in Eq. (II.16), it is possible to relate the free energy (B.1) to the normalisation of the restricted Boltzmann–Gibbs measure (B.4). More precisely, the modified potential $\tilde{V}(q) = V(q) + V_{\text{fix}}(q)$ gives rise to a new Hamiltonian denoted $\tilde{H}(p, q) = H(q, p) + V_{\text{fix}}(q)$ and an associated constrained measure $\tilde{\mu}_{\Sigma_\xi, v_\xi}$ with normalizing constant $\tilde{Z}_{z, v_z}$ (using the formulas in (B.4) with H replaced by $\tilde{H}$). We then obtain the following reformulation of free energy differences in terms of ratios of the normalizing constants $\tilde{Z}_{z, v_z}$.

Lemma B.1. For all $(z, v_z) \in \mathbb{R}^\ell \times \mathbb{R}^\ell$, it holds that

$$\tilde{Z}_{z, v_z} = (\det M)^{\frac{1}{2}} (2\pi\beta^{-1})^{\frac{d-\ell}{2}} \int_{\Sigma(z)} e^{-\beta V(q)} \exp\left(-\frac{\beta}{2} v_z^\top G_M(q)^{-1} v_z\right) \delta_{\xi(q)-z}(\mathrm{d}q).$$

In particular, for any $z_1, z_2 \in \mathbb{R}^\ell$,

$$\frac{\tilde{Z}_{z_1, 0}}{\tilde{Z}_{z_2, 0}} = e^{-\beta(F(z_1) - F(z_2))}. \quad (\text{B.5})$$

Proof. The proof relies on Ref. 39. From Sec. 2.3 therein, we restate the following two identities: By the co-area formula, one can show that

$$\delta_{\xi(q)-z}(\mathrm{d}q) = (\det M)^{-1/2} |\det G_M(q)|^{-1/2} \sigma_{\Sigma(z)}^M(\mathrm{d}q). \quad (\text{B.6})$$

Furthermore, the measure $\sigma_{\Sigma_\xi, v_\xi}(z, v_z)$ can be written as a product of two measures:

$$\sigma_{\Sigma_\xi, v_\xi}(z, v_z)(\mathrm{d}q \mathrm{d}p) = \sigma_{\Sigma_{v_\xi(q, \cdot)}(v_z)}^{M^{-1}}(\mathrm{d}p) \sigma_{\Sigma(z)}^M(\mathrm{d}q), \quad (\text{B.7})$$

where we denote by $\sigma_{\Sigma(z)}^M(\mathrm{d}q)$ the surface measure on $\Sigma(z)$ induced by the scalar product $\langle q, \tilde{q} \rangle_M = q^\top M \tilde{q}$ on $\mathbb{R}^d$, and, for a given $q \in \Sigma(z)$, by $\sigma_{\Sigma_{v_\xi(q, \cdot)}(v_z)}^{M^{-1}}(\mathrm{d}p)$ the surface measure on the affine space $\Sigma_{v_\xi(q, \cdot)}(v_z)$, induced by the scalar product $\langle p, \tilde{p} \rangle_{M^{-1}} = p^\top M^{-1} \tilde{p}$ on $\mathbb{R}^d$. For more precise definitions of these measures, we refer to Ref. 38, Sections 3.2.1 and 3.3.2 and the references therein.

Using the above identities, we write

$$\begin{aligned} \tilde{Z}_{z, v_z} &= \int_{\Sigma_\xi, v_\xi}(z, v_z) e^{-\beta \tilde{H}(q, p)} \sigma_{\Sigma_\xi, v_\xi}(z, v_z)(\mathrm{d}q \mathrm{d}p) \\ &= \int_{\Sigma_\xi, v_\xi}(z, v_z) e^{-\beta H(q, p)} (\det(G_M(q)))^{-1/2} \sigma_{\Sigma_{v_\xi(q, \cdot)}(v_z)}^{M^{-1}}(\mathrm{d}p) \sigma_{\Sigma(z)}^M(\mathrm{d}q) \\ &= \int_{\Sigma(z)} \int_{\Sigma_{v_\xi(q, \cdot)}(v_z)} \exp\left(-\frac{\beta}{2} p^\top M^{-1} p\right) \sigma_{\Sigma_{v_\xi(q, \cdot)}(v_z)}^{M^{-1}}(\mathrm{d}p) e^{-\beta V(q)} (\det(G_M(q)))^{-1/2} \sigma_{\Sigma(z)}^M(\mathrm{d}q) \\ &= \int_{\Sigma(z)} (2\pi\beta^{-1})^{\frac{d-\ell}{2}} \exp\left(-\frac{\beta}{2} v_z^\top G_M(q)^{-1} v_z\right) e^{-\beta V(q)} (\det(G_M(q)))^{-1/2} \sigma_{\Sigma(z)}^M(\mathrm{d}q) \\ &= (\det M)^{\frac{1}{2}} (2\pi\beta^{-1})^{\frac{d-\ell}{2}} \int_{\Sigma(z)} e^{-\beta V(q)} \exp\left(-\frac{\beta}{2} v_z^\top G_M(q)^{-1} v_z\right) \delta_{\xi(q)-z}(\mathrm{d}q). \end{aligned}$$

Equation (B.5) follows by taking $v_z = 0$. For the last but one equality in the series of equalities above, we use a computation from Sec. 5.1 in Ref. 39, namely, for fixed q ,

$$\begin{aligned} &\int_{\Sigma_{v_\xi(q, \cdot)}(v_z)} \exp\left(-\frac{\beta}{2} p^\top M^{-1} p\right) \sigma_{\Sigma_{v_\xi(q, \cdot)}(v_z)}^{M^{-1}}(\mathrm{d}p) \\ &= \exp\left(-\frac{\beta}{2} v_z^\top G_M(q)^{-1} v_z\right) \int_{\Sigma_{v_\xi(q, \cdot)}(0)} \exp\left(-\frac{\beta}{2} p^\top M^{-1} p\right) \sigma_{\Sigma_{v_\xi(q, \cdot)}(0)}^{M^{-1}}(\mathrm{d}p) \\ &= (2\pi\beta^{-1})^{\frac{d-\ell}{2}} \exp\left(-\frac{\beta}{2} v_z^\top G_M(q)^{-1} v_z\right). \end{aligned}$$

The first equality follows from a change of variable: the space $\Sigma_{v_\xi(q, \cdot)}(v_z)$ is composed of momenta fulfilling $0 = v_\xi(q, p) - v_z = \nabla \xi(q)^\top M^{-1} (p - \nabla \xi(q) G_M^{-1}(q) v_z)$, and therefore, when replacing p by $p - \nabla \xi(q) G_M^{-1}(q) v_z$, the resulting integral is taken over the space $\Sigma_{v_\xi(q, \cdot)}(0)$. $\square$

Remark B.2. When the Gram tensor $G_M(q)$ is independent of q , the introduction of the Fixman term becomes unnecessary and one furthermore has that

$$\frac{\tilde{\mathcal{Z}}_{z_1, v_z}}{\tilde{\mathcal{Z}}_{z_2, v_z}} = \frac{\mathcal{Z}_{z_1, v_z}}{\mathcal{Z}_{z_2, v_z}} = e^{-\beta(F(z_1) - F(z_2))} \quad (\text{B.8})$$

for any velocity $v_z \in \mathbb{R}^\ell$.

2. Properties of the steered dynamics

In this section, we state a number of properties of the RATTLE scheme introduced in (II.17)-(II.19). Since we follow dynamics for the energy modified by the Fixman term, in principle we would need to put a tilde on all associated measures and quantities, but we omit it here for the sake of readability and just consider the dynamics for a general potential V .

a. Forward steered dynamics

Let us make two remarks concerning the forward steered dynamics (II.17)-(II.19)

Remark B.3. In Eq. (II.18), the constraint (C_p) yields a unique well-defined Lagrange multiplier $\lambda^{k+3/4}$ which can be obtained by solving the linear equation

$$v_z(t_{k+1}) = \nabla \xi(q^{k+1})^T M^{-1} p^{k+1/2} - \frac{\Delta t}{2} \nabla \xi(q^{k+1})^T M^{-1} \nabla V(q^{k+1}) + \nabla \xi(q^{k+1})^T M^{-1} \nabla \xi(q^{k+1}) \lambda^{k+3/4},$$

since the Gram tensor $G_M(q^{k+1}) = \nabla \xi(q^{k+1})^T M^{-1} \nabla \xi(q^{k+1})$ is invertible by assumption. In contrast, the constraint (C_q) leads to a non-linear equation for $\lambda^{k+1/2}$, which is solved using Newton's method. In practice, we never encountered non-convergence of Newton's method in our numerical experiments. See however Ref. 48 and Sec. 3.2.1 of Ref. 39) for techniques to deal with this difficulty, using so-called reversibility checks.

Remark B.4. Notice that the velocity schedule $(v_z^{(Z_n, \tilde{Z}_{n+1})}(t_k))_{0 \leq n \leq K_T}$ can in principle be chosen independently of the position schedule $(z^{(Z_n, \tilde{Z}_{n+1})}(t_k))_{0 \leq k \leq K_T}$. It is only in the continuous-time limit that the schedule $z(t)$ directly imposes the velocity schedule as $v_z(t) = \dot{z}(t)$. Still, certain choices can be numerically beneficial, as explained in Ref. 39, Sec. 5.4.1. In (II.18), the unconstrained part of the update is given by

$$\tilde{q}^{k+1} = q^k + \Delta t M^{-1} p^{k+1/4} - \frac{\Delta t^2}{2} M^{-1} \nabla V(q^k),$$

which is followed by a projection with $q^{k+1} = \tilde{q}^{k+1} + \Delta t M^{-1} \nabla \xi(q^k) \lambda^{k+1/2}$. The standard choice $v_z(t_k) = \frac{z(t_{k+1}) - z(t_k)}{\Delta t}$ ensures that $\xi(\tilde{q}^{k+1}) = z(t_{k+1}) + \mathcal{O}(\Delta t^2)$. However, such a choice is not possible in our context since it would violate the reversibility property (II.8). One simple choice that fulfills the requirements is presented in Section II B 2. Another canonical choice with a meaningful limit in continuous time that satisfies the reversibility condition is

$$v_z(t_k) = \frac{z(t_{k+1}) - z(t_{k-1}))}{2\Delta t}, 0 \leq k \leq K_T,$$

requiring just the auxiliary extension of the position schedule to times t_{-1} and t_{K_T+1} . To fulfill the requirement that $v_z(t_0) = v_z(t_{K_T}) = 0$, this implies that $z(t_{-1}) = z(t_1)$ and $z(t_{K_T+1}) = z(t_{K_T-1})$.

b. Backward steered dynamics

For given position schedule $(z(t_k))_{0 \leq k \leq K_T}$ and velocity schedule $(v_z(t_k))_{0 \leq k \leq K_T}$, we define the following backward process. Starting from $(q^{b,0}, p^{b,0}) \in \Sigma_{\xi, v_\xi}(z(t_{K_T}), v_z(t_{K_T}))$, iterate the following procedure on $0 \leq k' \leq K_T - 1$:

$$p^{b,k'+1/4} = p^{b,k'} - \frac{\Delta t}{4} \gamma_P(q^{b,k'}) M^{-1}(p^{b,k'+1/4} + p^{b,k'}) + \sqrt{\frac{\Delta t}{2}} \sigma_P(q^{b,k'}) \mathcal{G}^{b,k'} \quad (\text{B.9})$$

$$\begin{cases} p^{b,k'+1/2} = p^{b,k'+1/4} + \frac{\Delta t}{2} \nabla V(q^{b,k'}) + \nabla \xi(q^{b,k'}) \lambda^{b,k'+1/2} \\ q^{b,k'+1} = q^{b,k'} - \Delta t M^{-1} p^{b,k'+1/2} \\ \xi(q^{b,k'+1}) = z(t_{K_T-k'-1}) \end{cases} \quad (C_q) \quad (\text{B.10})$$

$$\begin{cases} p^{b,k'+3/4} = p^{b,k'+1/2} + \frac{\Delta t}{2} \nabla V(q^{b,k'+1}) + \nabla \xi(q^{b,k'+1}) \lambda^{b,k'+3/4} \\ \nabla \xi(q^{b,k'+1})^T M^{-1} p^{b,k'+3/4} = v_z(t_{K_T-k'-1}) \end{cases} \quad (C_p) \quad (\text{B.11})$$

where $(\mathcal{G}^{b,k'})$ and $(\mathcal{G}^{b,k'+1/2})$ are independent sequences of i.i.d. centered Gaussian random vectors with covariance matrix Id_d and $\lambda^{b,k'+1/2}$ and $\lambda^{b,k'+3/4}$ are Lagrange multipliers such that the conditions (C_q) and (C_p) are fulfilled.

c. Relation between processes with opposite endpoints

Thanks to the reversibility property of the schedule (II.7)-(II.8), we can relate the law of the forward and backward steered dynamics with schedules with swapped endpoints one to another: With the right initialization, the solution of the backward dynamics for a schedule with endpoints $(z, \tilde{z})$ has the same law as the forward dynamics for a schedule with endpoints $(\tilde{z}, z)$ up to a momentum flip. We make the statement more precise in the following lemma.

Lemma B.5. Consider a forward process $((q^k, p^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T}, p^{K_T})$ for a schedule $(z^{(\tilde{z}, z)}(t_k))_{0 \leq k \leq K_T}, (v_z^{(\tilde{z}, z)}(t_k))_{0 \leq k \leq K_T}$ with initial conditions

$$(q^0, p^0) \sim \mu_{\Sigma_{\xi, v_\xi}(z^{(\tilde{z}, z)}(t_0), v_z^{(\tilde{z}, z)}(t_0))}(\text{d}q \text{d}p), \quad (\text{B.12})$$

and a backward process $((q^{b,k}, p^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T}, p^{b,K_T})$ for a schedule with swapped endpoints $(z^{(z, \tilde{z})}(t_k))_{0 \leq k \leq K_T}, (v_z^{(z, \tilde{z})}(t_k))_{0 \leq k \leq K_T}$ and initial conditions

$$(q^{b,0}, p^{b,0}) \sim \mu_{\Sigma_{\xi, v_\xi}(z^{(z, \tilde{z})}(t_{K_T}), v_z^{(z, \tilde{z})}(t_{K_T}))}(\text{d}q \text{d}p). \quad (\text{B.13})$$

Then, this backward process has the same law as the momentum flipped version of the forward process, $((q^k, -p^k, -p^{k+1/4}, -p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T}, -p^{K_T})$.

Proof. The initial distribution of the forward process satisfies $\mu_{\Sigma_{\xi, v_\xi}(z^{(\tilde{z}, z)}(t_0), v_z^{(\tilde{z}, z)}(t_0))} = \mu_{\Sigma_{\xi, v_\xi}(z^{(z, \tilde{z})}(t_{K_T}), -v_z^{(z, \tilde{z})}(t_{K_T}))}$ by the symmetry properties (II.7)-(II.8) of the schedule. Since

$$(q, p) \in \Sigma_{\xi, v_\xi}(z, -v_z) \Leftrightarrow (q, -p) \in \Sigma_{\xi, v_\xi}(z, v_z),$$

and from the definition of the constrained measure in (B.4) with the symmetry of the Hamiltonian $H(q, p) = H(q, -p)$, this implies that $(q^{b,0}, p^{b,0})$ and $(q^0, -p^0)$ are indeed drawn from the same distribution. From there, using again the symmetry of the schedule, one can easily verify that the equations describing the forward dynamics with endpoints $(\tilde{z}, z)$ for $(q^k, -p^k)$ are exactly the same as the equations describing the backward dynamics with endpoints $(z, \tilde{z})$ for $(q^{b,k}, p^{b,k})$. $\square$

d. Symplecticity and reversibility of the deterministic flows

Lemma B.6. Consider the numerical flow

$$\Phi^k : \begin{cases} \Sigma_{\xi, v_\xi}(z(t_k), v_z(t_k)) \rightarrow \Sigma_{\xi, v_\xi}(z(t_{k+1}), v_z(t_{k+1})) \\ (q^k, p^{k+1/4}) \mapsto (q^{k+1}, p^{k+3/4}) \end{cases} \quad (\text{B.14})$$

associated with the deterministic part of the forward dynamics (II.18) as well as the flow

$$\Phi^{b, k'} : \begin{cases} \Sigma_{\xi, v_\xi}(z(t_{K_T-k'}), v_z(t_{K_T-k'})) \rightarrow \Sigma_{\xi, v_\xi}(z(t_{K_T-k'-1}), v_z(t_{K_T-k'-1})) \\ (q^{b, k'}, p^{b, k'+1/4}) \mapsto (q^{b, k'+1}, p^{b, k'+3/4}) \end{cases} \quad (\text{B.15})$$

associated with the deterministic part of the backward dynamics (B.10).

The flow Φ^k is a symplectic map for all $k \in \{0, \dots, K_T - 1\}$, and the same holds true for $\Phi^{b, k'}$ for all $k' \in \{0, \dots, K_T - 1\}$.

To prove this statement, we show that the 2-form $dq \wedge dp$ is preserved, where $\wedge$ denotes the wedge-product (see Ref. 49 for a textbook introduction to symplectic structures). For the simplest version with a constant constraint, this is a standard result found for example in Refs. 50 and 51. We follow here the same arguments as in Ref. 51. The differentials obey:

$$dp^{k+1/2} = dp^{k+1/4} - \frac{\Delta t}{2} d(\nabla V(q^k)) + d(\nabla \xi(q^k) \lambda^{k+1/2}) \quad (\text{B.16})$$

$$dq^{k+1} = dq^k + \Delta t M^{-1} dp^{k+1/2} \quad (\text{B.17})$$

$$\nabla \xi(q^{k+1})^\top dq^{k+1} = 0 \quad (\text{B.18})$$

$$dp^{k+3/4} = dp^{k+1/2} - \frac{\Delta t}{2} d(\nabla V(q^{k+1})) + d(\nabla \xi(q^{k+1}) \lambda^{k+3/4}). \quad (\text{B.19})$$

Denoting the Hessian of V by $\nabla^2 V$, we have $d(\nabla V(q^k)) = \nabla^2 V(q^k) dq^k$. Taking the wedge product at the endpoints, we have

$$\begin{aligned} dq^{k+1} \wedge dp^{k+3/4} &= dq^{k+1} \wedge \left(dp^{k+1/2} - \frac{\Delta t}{2} d(\nabla V(q^{k+1})) + d(\nabla \xi(q^{k+1}) \lambda^{k+3/4}) \right) \\ &= dq^{k+1} \wedge dp^{k+1/2} - \frac{\Delta t}{2} dq^{k+1} \wedge \nabla^2 V(q^{k+1}) dq^{k+1} + dq^{k+1} \wedge d(\nabla \xi(q^{k+1}) \lambda^{k+3/4}) \end{aligned} \quad (\text{B.20})$$

As in the proof by Ref. 51, we make use of the following two lemmata.

Lemma B.7. Let du be an arbitrary differential in $\mathbb{R}^n$ and let A be any $n \times n$ real symmetric matrix. Then

$$du \wedge (Adu) = 0.$$

Proof. Writing this expression component-wise, we have

$$du \wedge (Adu) = \sum_{i,j=1}^n du_i \wedge (A_{ij} du_j) = \sum_{i,j=1}^n A_{ij} du_i \wedge du_j = - \sum_{i,j=1}^n A_{ji} du_j \wedge du_i = -du \wedge (Adu)$$

by the symmetry of A and the antisymmetry of the wedge-product, and thus $du \wedge (Adu) = 0$. $\square$

Lemma B.8. Let du and dv be arbitrary differentials in $\mathbb{R}^n$ and $\mathbb{R}^m$, respectively, and let B be any $n \times m$ real matrix, then

$$du \wedge (Bdv) = (B^\top du) \wedge dv.$$

Proof. Writing this expression component-wise, we have

$$du \wedge (Bdv) = \sum_{i=1}^n \sum_{j=1}^m du_i \wedge (B_{ij} dv_j) = \sum_{i=1}^n \sum_{j=1}^m (B_{ij} du_i) \wedge dv_j = \sum_{i=1}^n \sum_{j=1}^m (B_{ji}^\top du_i) \wedge dv_j = (B^\top du) \wedge dv,$$

which is the desired result. $\square$

Since the Hessian is symmetric, the second term in (B.20) vanishes by Theorem B.7. The third term in (B.20) also vanishes:

$$dq^{k+1} \wedge d(\nabla \xi(q^{k+1}) \lambda^{k+3/4}) = dq^{k+1} \wedge \nabla \xi(q^{k+1}) d\lambda^{k+3/4} + \sum_{i=1}^{\ell} \lambda_i^{k+3/4} dq^{k+1} \wedge (\nabla^2 \xi_i)(q^{k+1}) dq^{k+1} = 0,$$

where ξ_i is the i -th component of the CV. Indeed, thanks to Theorem B.8, we have $dq^{k+1} \wedge \nabla \xi(q^{k+1}) d\lambda^{k+3/4} = (\nabla \xi(q^{k+1}))^\top dq^{k+1} \wedge d\lambda^{k+3/4}$ which is zero due to (B.18). The final term equally vanishes since the Hessian is symmetric and one can thus apply Theorem B.7 to each term of the sum.

We therefore have that

$$dq^{k+1} \wedge dp^{k+3/4} = dq^{k+1} \wedge dp^{k+1/2}.$$

Using (B.17), we can continue (with the same reasoning as before)

$$\begin{aligned} dq^{k+1} \wedge dp^{k+1/2} &= dq^k \wedge dp^{k+1/2} + \Delta t M^{-1} dp^{k+1/2} \wedge dp^{k+1/2} \\ &= dq^k \wedge \left(dp^{k+1/4} - \frac{\Delta t}{2} d(\nabla V(q^k)) + d(\nabla \xi(q^k) \lambda^{k+1/2}) \right) \\ &= dq^k \wedge dp^{k+1/4} - \frac{\Delta t}{2} dq^k \wedge d(\nabla V(q^k)) + dq^k \wedge d(\nabla \xi(q^k) \lambda^{k+1/2}) \\ &= dq^k \wedge dp^{k+1/4} - \frac{\Delta t}{2} dq^k \wedge \nabla^2 V(q^k) dq^k + dq^k \wedge \left(\nabla \xi(q^k) d\lambda^{k+1/2} + \sum_{i=1}^{\ell} \lambda_i^{k+1/2} (\nabla^2 \xi_i)(q^k) dq^k \right) \\ &= dq^k \wedge dp^{k+1/4} - \frac{\Delta t}{2} dq^k \wedge \nabla^2 V(q^k) dq^k + (\nabla \xi(q^k))^\top dq^k \wedge d\lambda^{k+1/2} + \sum_{i=1}^{\ell} \lambda_i^{k+1/2} dq^k \wedge (\nabla^2 \xi_i)(q^k) dq^k \\ &= dq^k \wedge dp^{k+1/4}. \end{aligned}$$

Following exactly the same arguments, one can equally show that $dq^{b,k'} \wedge dp^{b,k'+1/4} = dq^{b,k'+1} \wedge dp^{b,k'+3/4}$. This concludes the proof of Theorem B.6.

In all of the following, we assume that the time-step Δt is sufficiently small so that, along the trajectory, the forward and backward flow associated with the deterministic parts (II.18) and (B.10) satisfy a reversibility property, see e.g. Sect. VII.1.4 of Ref. 50 and Sect. 5.4.3 of Ref. 39, as well as Lemma 2.9 in Ref. 48 where a precise setting is exhibited for which this is satisfied. More precisely, we assume that the following pointwise equality holds for the forward and backward flows Φ^k and $\Phi^{b,k'}$ introduced in (B.14) and (B.15). For all the configurations which are visited and all $k \in \{1, \dots, K_T\}$:

$$\Phi^{b,K_T-k} \circ \Phi^{k-1} = \text{Id}. \quad (\text{B.21})$$

e. Jarzynski–Crooks equality

We are now in position to state the Jarzynski–Crooks formula. We state it here in a slightly more general form than in Ref. 39, but otherwise follow exactly the presentation there.

Theorem B.9. *Consider the distribution $\mu_{\Sigma_\xi, v_\xi}(z(t), v_z(t))$ and its normalizing constant $\mathcal{Z}_{z(t), v_z(t)}$ defined in (B.4). Denote by $((q^k, p^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T}, p^{K_T})$ the solution of the forward discretized Langevin dynamics (II.17)–(II.19) with initial conditions distributed according to*

$$(q^0, p^0) \sim \mu_{\Sigma_\xi, v_\xi}(z(t_0), v_z(t_0)) (dq dp), \quad (\text{B.22})$$

and by $((q^{b,k}, p^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T}, p^{b,K_T})$ the solution of the discretized backward Langevin dynamics (B.9)-(B.11) with initial conditions distributed according to

$$(q^{b,0}, p^{b,0}) \sim \mu_{\Sigma_\xi, v_\xi}(z(t_{K_T}), v_z(t_{K_T})) (dq dp). \quad (\text{B.23})$$

Then, the following Jarzynski-Crooks identity holds: for any bounded discrete path functional $\varphi_{[0, K_T]}$,

$$\frac{\mathcal{Z}_{z(K_T), v_z(t_{K_T})}}{\mathcal{Z}_{z(t_0), v_z(t_0)}} = \frac{\mathbb{E} \left(\varphi_{[0, K_T]} \left((q^k, p^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T}, p^{K_T} \right) e^{-\beta \mathcal{W}^{K_T}} \right)}{\mathbb{E} \left(\varphi_{[0, K_T]}^r \left((q^{b,k}, p^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T}, p^{b,K_T} \right) \right)}, \quad (\text{B.24})$$

where $\mathcal{W}^{K_T}$ is computed according to (II.22) with $\tilde{H} = H$, and $(\cdot)^r$ denotes the composition with the operation of time reversal of paths:

$$\begin{aligned} \varphi_{[0, K_T]}^r & \left((q^{b,k}, p^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T}, p^{b,K_T} \right) \\ &= \varphi_{[0, K_T]} \left((q^{b, K_T-k}, p^{b, K_T-k}, p^{b, K_T-k-1/4}, p^{b, K_T-k-3/4})_{0 \leq k \leq K_T-1}, q^{b,0}, p^{b,0} \right). \end{aligned} \quad (\text{B.25})$$

Proof. The proof is the same as the proof of Ref. 39, Theorem 5.5. The only difference lies in the slightly different momentum constraint (C_p) for the forward and backward dynamics. Since we show in Theorem B.6 that the forward and backward deterministic flows are symplectic, the proof goes through in the same way. $\square$

3. Proof of reversibility

We are now in a position to prove Theorem II.4, namely that the algorithm presented in Section II B is reversible with respect to the target probability measure ν . Assume that Q_n is distributed according to ν . The aim is to prove that (Q_n, Q_{n+1}) has the same law as (Q_{n+1}, Q_n) . In order to study the law of (Q_n, Q_{n+1}) , let us consider a bounded measurable function $\varphi : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$. Recall that $Z_n = \xi(Q_n)$ and $\tilde{Z}_{n+1} = \xi(\tilde{Q}_{n+1})$. One has:

$$\begin{aligned} \mathbb{E}(\varphi(Q_n, Q_{n+1})) &= \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \mathbf{1}_{U_{n+1} \leq \exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})}} \right) \\ &+ \mathbb{E} \left(\varphi(Q_n, Q_n) \mathbf{1}_{U_{n+1} > \exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})}} \right) \\ &= \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right) \\ &+ \mathbb{E} \left(\varphi(Q_n, Q_n) \left[1 - 1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right), \end{aligned}$$

where here and in the following, $a \wedge b = \min(a, b)$. The expectation value is taken over $Q_n \sim \nu$ (which fixes Z_n), and all the random variables used to generate Q_{n+1} : sampling $\tilde{Z}_{n+1}$ from the density $\rho(Z_n, \cdot)$, following the steered dynamics to obtain $\tilde{Q}_{n+1}$ and accepting or rejecting the final move. The expectation value over Q_n can also be understood in two steps:

- First, sample $Z_n \sim \xi \# \nu$, where $(\xi \# \nu)(dz) = \exp(-\beta F(z)) dz$ denotes the image of the probability measure ν by ξ (see (B.1) for the definition of the free energy F).
- Second, sample a full state Q_n from $\nu(dq | \xi(q) = Z_n)$, where $\nu(dq | \xi(q) = z)$ denotes the family of conditional measures

$$\nu(dq | \xi(q) = z) = \frac{Z_q^{-1} e^{-\beta V(q)} \delta_{\xi(q)-z}(dq)}{Z(z)},$$

where $Z(z) = Z_q^{-1} \int_{\Sigma(z)} e^{-\beta V(q)} \delta_{\xi(q)-z}(dq) = \exp(-\beta F(z))$ is the normalizing constant.

By conditioning on the collective variables $(Z_n, \tilde{Z}_{n+1})$, one gets

$$\begin{aligned}
& \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right) \\
&= \int_{z, \tilde{z}} \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\xi(\tilde{Q}_{n+1}), \xi(Q_n))}{\rho(\xi(Q_n), \xi(\tilde{Q}_{n+1}))} \right) \right] \middle| (\xi(Q_n), \xi(\tilde{Q}_{n+1})) = (z, \tilde{z}) \right) \\
&\quad \times \exp(-\beta F(z)) \rho(z, \tilde{z}) \, dz \, d\tilde{z} \\
&= \int_{z, \tilde{z}} \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[\rho(\xi(Q_n), \xi(\tilde{Q}_{n+1})) \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \rho(\xi(\tilde{Q}_{n+1}), \xi(Q_n)) \right) \right] \middle| (\xi(Q_n), \xi(\tilde{Q}_{n+1})) = (z, \tilde{z}) \right) \\
&\quad \times \exp(-\beta F(z)) \, dz \, d\tilde{z}.
\end{aligned}$$

The conditional expectation value depends on the full steered path (and not just on the endpoints) via the work defined by (II.22), which we can rewrite here as

$$\mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) = \sum_{k=0}^{K_T-1} \left(\tilde{H}(q^{k+1}, p^{k+3/4}) - \tilde{H}(q^k, p^{k+1/4}) \right). \quad (\text{B.26})$$

Notice that the work of a time-reversed path is the negative of the work of the path without time-reversal:

$$\begin{aligned}
& \mathcal{W} \left((q^{K_T-k}, p^{K_T-k-1/4}, p^{K_T-k-3/4})_{0 \leq k \leq K_T-1}, q^0 \right) \\
&= \sum_{k=0}^{K_T-1} \left(\tilde{H}(q^{K_T-k-1}, p^{K_T-k-3/4}) - \tilde{H}(q^{K_T-k}, p^{K_T-k-1/4}) \right) \\
&= \sum_{k=0}^{K_T-1} \left(\tilde{H}(q^k, p^{k+1/4}) - \tilde{H}(q^{k+1}, p^{k+3/4}) \right) \\
&= -\mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right). \quad (\text{B.27})
\end{aligned}$$

Using the generalized Jarzynski–Crooks identity, one gets, introducing the notation $((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T})$ for the forward process and $((q^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T})$ for the backward process as in Theorem B.9,

$$\begin{aligned}
& \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[\rho(\xi(Q_n), \xi(\tilde{Q}_{n+1})) \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \rho(\xi(\tilde{Q}_{n+1}), \xi(Q_n)) \right) \right] \middle| (\xi(Q_n), \xi(\tilde{Q}_{n+1})) = (z, \tilde{z}) \right) \\
&= \mathbb{E} \left(\varphi(q^0, q^{K_T}) \left[\rho(\xi(q^0), \xi(q^{K_T})) \right. \right. \\
&\quad \left. \left. \wedge \left(\exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \rho(\xi(q^{K_T}), \xi(q^0)) \right) \right] \middle| (\xi(q^0), \xi(q^{K_T})) = (z, \tilde{z}) \right) \\
&= \mathbb{E} \left(\varphi(q^0, q^{K_T}) \left[\left(\exp \left(\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \rho(\xi(q^0), \xi(q^{K_T})) \right) \right. \right. \\
&\quad \left. \left. \wedge \rho(\xi(q^{K_T}), \xi(q^0)) \right] \exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \middle| (\xi(q^0), \xi(q^{K_T})) = (z, \tilde{z}) \right) \\
&\stackrel{(\text{B.24})}{=} \mathbb{E} \left(\varphi(q^{b,K_T}, q^{b,0}) \left[\left(\exp \left(\beta \mathcal{W} \left((q^{b,K_T-k}, p^{b,K_T-k-1/4}, p^{b,K_T-k-3/4})_{0 \leq k \leq K_T-1}, q^{b,0} \right) \right) \rho(\xi(q^{b,K_T}), \xi(q^{b,0})) \right) \right. \right. \\
&\quad \left. \left. \wedge \rho(\xi(q^{b,0}), \xi(q^{b,K_T})) \right] \middle| (\xi(q^{b,0}), \xi(q^{b,K_T})) = (\tilde{z}, z) \right) \frac{\tilde{\mathcal{Z}}(\tilde{z}, v_z(t_{K_T}))}{\tilde{\mathcal{Z}}(z, v_z(t_0))} \\
&\stackrel{(\text{B.5})}{=} \mathbb{E} \left(\varphi(q^{b,K_T}, q^{b,0}) \left[\left(\exp \left(\beta \mathcal{W} \left((q^{b,K_T-k}, p^{b,K_T-k-1/4}, p^{b,K_T-k-3/4})_{0 \leq k \leq K_T-1}, q^{b,0} \right) \right) \rho(\xi(q^{b,K_T}), \xi(q^{b,0})) \right) \right. \right. \\
&\quad \left. \left. \wedge \rho(\xi(q^{b,0}), \xi(q^{b,K_T})) \right] \middle| (\xi(q^{b,0}), \xi(q^{b,K_T})) = (\tilde{z}, z) \right) \exp(-\beta(F(\tilde{z}) - F(z))) \\
&\stackrel{(\text{B.27})}{=} \mathbb{E} \left(\varphi(q^{b,K_T}, q^{b,0}) \left[\left(\exp \left(-\beta \mathcal{W} \left((q^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T} \right) \right) \rho(\xi(q^{b,K_T}), \xi(q^{b,0})) \right) \right. \right. \\
&\quad \left. \left. \wedge \rho(\xi(q^{b,0}), \xi(q^{b,K_T})) \right] \middle| (\xi(q^{b,0}), \xi(q^{b,K_T})) = (\tilde{z}, z) \right) \exp(-\beta(F(\tilde{z}) - F(z))). \quad (\text{B.28})
\end{aligned}$$

In the last but one step, we used the assumption (II.6) that the initial and final velocity is zero, $v_z(t_0) = v_z(t_{K_T}) = 0$, and applied the identity (B.5). For the final step, we used the symmetry property (B.27) of the work function.

Remark B.10. As mentioned in Theorem B.2, when G_M does not depend on q , there is no Fixman term to be considered and (B.8) holds. One can use this equation instead of (B.5) in the computation above, and this in turn implies that reversibility can be proven only assuming that the initial and final velocities are equal i.e. $v_z(t_0) = v_z(t_{K_T})$, but not necessarily zero (see condition (II.6)).

We can use Theorem B.5 to rewrite the latter expectation value over a backward path as one over a forward path as

$$\begin{aligned}
& \mathbb{E} \left(\varphi(q^{b,K_T}, q^{b,0}) \left[\left(\exp \left(-\beta \mathcal{W} \left((q^{b,k}, p^{b,k+1/4}, p^{b,k+3/4})_{0 \leq k \leq K_T-1}, q^{b,K_T} \right) \right) \rho(\xi(q^{b,K_T}), \xi(q^{b,0})) \right) \right. \right. \\
& \quad \left. \left. \wedge \rho(\xi(q^{b,0}), \xi(q^{b,K_T})) \right) \middle| (\xi(q^{b,0}), \xi(q^{b,K_T})) = (\tilde{z}, z) \right) \exp(-\beta(F(\tilde{z}) - F(z))) \right) \\
&= \mathbb{E} \left(\varphi(q^{K_T}, q^0) \left[\left(\exp \left(-\beta \mathcal{W} \left((q^k, -p^{k+1/4}, -p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \rho(\xi(q^{K_T}), \xi(q^0)) \right) \right. \right. \\
& \quad \left. \left. \wedge \rho(\xi(q^0), \xi(q^{K_T})) \right) \middle| (\xi(q^0), \xi(q^{K_T})) = (\tilde{z}, z) \right) \exp(-\beta(F(\tilde{z}) - F(z))) \right) \\
&= \mathbb{E} \left(\varphi(q^{K_T}, q^0) \left[\left(\exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \rho(\xi(q^{K_T}), \xi(q^0)) \right) \right. \right. \\
& \quad \left. \left. \wedge \rho(\xi(q^0), \xi(q^{K_T})) \right) \middle| (\xi(q^0), \xi(q^{K_T})) = (\tilde{z}, z) \right) \exp(-\beta(F(\tilde{z}) - F(z))) \right). \tag{B.29}
\end{aligned}$$

In the last step, we used that the work function $\mathcal{W}$ is symmetric with respect to momentum reversal since $\tilde{H}(q, p) = \tilde{H}(q, -p)$. Using (B.28) and (B.29), we get (by just swapping the indices z and $\tilde{z}$ in the third equality)

$$\begin{aligned}
& \mathbb{E} \left(\varphi(Q_n, \tilde{Q}_{n+1}) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right) \\
&= \int_{z, \tilde{z}} \mathbb{E} \left(\varphi(q^{K_T}, q^0) \left[\left(\exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \rho(\xi(q^{K_T}), \xi(q^0)) \right) \right. \right. \\
& \quad \left. \left. \wedge \rho(\xi(q^0), \xi(q^{K_T})) \right) \middle| (\xi(q^0), \xi(q^{K_T})) = (\tilde{z}, z) \right) \exp(-\beta F(\tilde{z})) \, dz \, d\tilde{z} \right) \\
&= \int_{z, \tilde{z}} \mathbb{E} \left(\varphi(q^{K_T}, q^0) \left[1 \wedge \left(\exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \frac{\rho(\xi(q^{K_T}), \xi(q^0))}{\rho(\xi(q^0), \xi(q^{K_T}))} \right) \right] \right. \\
& \quad \left. \middle| (\xi(q^0), \xi(q^{K_T})) = (\tilde{z}, z) \right) \rho(\tilde{z}, z) \exp(-\beta F(\tilde{z})) \, dz \, d\tilde{z} \right) \\
&= \int_{z, \tilde{z}} \mathbb{E} \left(\varphi(q^{K_T}, q^0) \left[1 \wedge \left(\exp \left(-\beta \mathcal{W} \left((q^k, p^{k+1/4}, p^{k+3/4})_{0 \leq k \leq K_T-1}, q^{K_T} \right) \right) \frac{\rho(\xi(q^{K_T}), \xi(q^0))}{\rho(\xi(q^0), \xi(q^{K_T}))} \right) \right] \right. \\
& \quad \left. \middle| (\xi(q^0), \xi(q^{K_T})) = (z, \tilde{z}) \right) \rho(z, \tilde{z}) \exp(-\beta F(z)) \, dz \, d\tilde{z} \right) \\
&= \mathbb{E} \left(\varphi(\tilde{Q}_{n+1}, Q_n) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right).
\end{aligned}$$

Finally,

$$\begin{aligned}
\mathbb{E}(\varphi(Q_n, Q_{n+1})) &= \mathbb{E} \left(\varphi(\tilde{Q}_{n+1}, Q_n) \left[1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right) \\
&\quad + \mathbb{E} \left(\varphi(Q_n, Q_n) \left[1 - 1 \wedge \left(\exp(-\beta \mathcal{W}_{n+1}) \frac{\rho(\tilde{Z}_{n+1}, Z_n)}{\rho(Z_n, \tilde{Z}_{n+1})} \right) \right] \right) \\
&= \mathbb{E}(\varphi(Q_{n+1}, Q_n)),
\end{aligned}$$

which concludes the proof of Theorem II.4.

Appendix C: Normalization of the steered dynamics (II.17)-(II.19)

We assume in this section that the mass matrix $M = M\text{Id}$ is isotropic and furthermore write the velocity schedule as $v_z(t_k) = \tilde{v}_z(t_k)/\Delta t$, with a velocity $\tilde{v}_z(t_k)$ normalized by the timestep. The parameterization then simplifies

considerably. With the dimensionless momentum $\tilde{p} = \sqrt{\beta/M}p$, which is of identity covariance under the equilibrium distribution, the steered dynamics (II.17)-(II.19) on $(q, \tilde{p})$ read

$$\tilde{p}^{k+1/4} = \tilde{p}^k - \frac{\Delta t \gamma}{4M} P_M(q^k) P_M(q^k)^\top (\tilde{p}^{k+1/4} + \tilde{p}^k) + \sqrt{\frac{\Delta t \gamma}{2M}} P_M(q^k) \mathcal{G}^k \quad (\text{C.1})$$

$$\begin{cases} \tilde{p}^{k+1/2} = \tilde{p}^{k+1/4} - \frac{\Delta t}{2} \sqrt{\frac{1}{\beta M}} \nabla(\beta \tilde{V}(q^k)) + \nabla \xi(q^k) \tilde{\lambda}^{k+1/2} \\ q^{k+1} = q^k + \frac{\Delta t}{\sqrt{\beta M}} \tilde{p}^{k+1/2} \\ \xi(q^{k+1}) = z(t_{k+1}) \end{cases} \quad (\text{C}_q) \quad (\text{C.2})$$

$$\begin{cases} \tilde{p}^{k+3/4} = \tilde{p}^{k+1/2} - \frac{\Delta t}{2} \sqrt{\frac{1}{\beta M}} \nabla(\beta \tilde{V}(q^{k+1})) + \nabla \xi(q^{k+1}) \tilde{\lambda}^{k+3/4} \\ \nabla \xi(q^{k+1})^\top \tilde{p}^{k+3/4} = \frac{\sqrt{\beta M}}{\Delta t} \tilde{v}_z(t_{k+1}) \end{cases} \quad (\text{C}_p) \quad (\text{C.3})$$

$$\tilde{p}^{k+1} = \tilde{p}^{k+3/4} - \frac{\Delta t \gamma}{4M} P_M(q^{k+1}) P_M(q^{k+1})^\top (\tilde{p}^{k+3/4} + \tilde{p}^{k+1}) + \sqrt{\frac{\Delta t \gamma}{2M}} P_M(q^{k+1}) \mathcal{G}^{k+1/2},$$

with renormalized Lagrange multipliers $\tilde{\lambda}^{k+1/2}$ and $\tilde{\lambda}^{k+3/4}$ that ensure that the constraints (C_q) and (C_p) are satisfied. Here, we have deliberately moved β into the gradients of the potential since it is the quantity βV that defines the target distribution at equilibrium (note that the correction βV_{fix} is independent of β). With this, **Step 2** of the algorithm in Section II B 1 is fully defined by the schedule $z(t_k)$ and $\tilde{v}_z(t_k)$ as well as the two parameters $\alpha_1 = \Delta t \gamma / (4M)$ and $\alpha_2 = \Delta t^2 / (\beta M)$. Note that while the Fixman term in general depends on the mass M , under the assumption of the isotropic mass matrix, the gradient $\nabla V_{\text{fix}}(q)$ does not.

Appendix D: Simplification for a linear CV

To explicitly connect the presented algorithm to the previous one from Ref. 43, consider the simplest case where the reaction coordinate is just given by the first ℓ degrees of freedom, so that we can write $q = (q_{\text{CV}}, q_\perp)$ with $q_{\text{CV}} = \xi(q) = (q_1, \dots, q_\ell)$ and $q_\perp = (q_{\ell+1}, \dots, q_d)$. Similarly, one can then write the momenta as $p = (p_{\text{CV}}, p_\perp)$. Assume once again that $M = M \text{Id}_d$ (with a slight abuse of notation). Then the Gram tensor is $G_M(q) = M^{-1} \text{Id}_\ell$, and the projection operator for the momenta is just

$$P_M(q) = \begin{pmatrix} \mathbf{0}_{\ell \times \ell} & 0 \\ 0 & \mathbf{1}_{(d-\ell) \times (d-\ell)} \end{pmatrix} = P_M(q)^\top$$

setting the momenta p_{CV} associated with the CV to zero. With the scalar parameters γ and σ , one then has $\gamma_P(q) = P_M(q) \gamma P_M(q)^\top = \gamma P_M(q)$ and $\sigma_P(q) = P_M(q) \sigma$. Explicitly solving for the Lagrange multipliers, the resulting dynamics read (using the notation $\nabla_\perp V$ for the partial derivative of V with respect to $q_\perp$)

$$\begin{cases} p_{\text{CV}}^{k+1/4} = p_{\text{CV}}^k \\ p_\perp^{k+1/4} = p_\perp^k - \frac{\Delta t}{4M} \gamma (p_\perp^{k+1/4} + p_\perp^k) + \sqrt{\frac{\Delta t}{2}} \sigma \mathcal{G}_\perp^k \end{cases} \quad (\text{D.1})$$

$$\begin{cases} p_\perp^{k+1/2} = p_\perp^{k+1/4} - \frac{\Delta t}{2} \nabla_\perp V(q^k) \\ q_{\text{CV}}^{k+1} = z(t_{k+1}) \\ q_\perp^{k+1} = q_\perp^k + \frac{\Delta t}{M} p_\perp^{k+1/2} \\ p_{\text{CV}}^{k+3/4} = M v_z(t_{k+1}) \\ p_\perp^{k+3/4} = p_\perp^{k+1/2} - \frac{\Delta t}{2} \nabla_\perp V(q^{k+1}) \end{cases} \quad (\text{D.2})$$

$$\begin{cases} p_{\text{CV}}^{k+1} = p_{\text{CV}}^{k+3/4}, \\ p_\perp^{k+1} = p_\perp^{k+3/4} - \frac{\Delta t}{4M} \gamma (p_\perp^{k+3/4} + p_\perp^{k+1}) + \sqrt{\frac{\Delta t}{2}} \sigma \mathcal{G}_\perp^{k+1/2}. \end{cases} \quad (\text{D.3})$$

In the following we consider two limiting cases, the overdamped limit and the deterministic limit.

1. Overdamped limit

For the choice $\gamma = 2$ and $M = \frac{\Delta t}{2}$ (and therefore $\sigma = \frac{2}{\sqrt{\beta}}$), the forward dynamics (D.1)-(D.3) simplify to overdamped Langevin dynamics

$$\begin{cases} q_{\text{CV}}^{k+1} = z(t_{k+1}) \\ q_{\perp}^{k+1} = q_{\perp}^k + \sqrt{\frac{2\Delta t}{\beta}} \mathcal{G}_{\perp}^k - \Delta t \nabla_{\perp} V(q^k). \end{cases} \quad (\text{D.4})$$

In terms of the parameters of the normalized algorithm introduced in Section II D, this parameter choice corresponds to setting $\alpha_1 = 1$. Using (II.22), the associated work is calculated as

$$\mathcal{W}^{K_T} = V(q^{K_T}) - V(q^0) + \frac{1}{4\Delta t} \sum_{k=0}^{K_T-1} \left(\|q_{\perp}^k - q_{\perp}^{k+1} + \Delta t \nabla_{\perp} V(q^{k+1})\|_2^2 - \|q_{\perp}^{k+1} - q_{\perp}^k + \Delta t \nabla_{\perp} V(q^k)\|_2^2 \right).$$

For the overall accept/reject probability in the MCMC algorithm, this gives

$$\exp(-\beta \mathcal{W}^{K_T}) = \frac{e^{-\beta V(q^{K_T})} \exp\left(-\frac{\beta}{4\Delta t} \sum_{k=0}^{K_T-1} \|q_{\perp}^k - q_{\perp}^{k+1} + \Delta t \nabla_{\perp} V(q^{k+1})\|_2^2\right)}{e^{-\beta V(q^0)} \exp\left(-\frac{\beta}{4\Delta t} \sum_{k=0}^{K_T-1} \|q_{\perp}^{k+1} - q_{\perp}^k + \Delta t \nabla_{\perp} V(q^k)\|_2^2\right)}.$$

This algorithm is very closely related to the **asymmetric** algorithm from Ref. 43, with the only difference that forward and backward path have swapped their definitions.

2. Deterministic limit

In the deterministic limit, we have $\gamma = \sigma = 0$. The forward dynamics then simplify to

$$\begin{cases} q_{\text{CV}}^{k+1} = z(t_{k+1}) \\ q_{\perp}^{k+1} = q_{\perp}^k + \frac{\Delta t}{M} p_{\perp}^k - \frac{\Delta t^2}{2M} \nabla_{\perp} V(q^k) \\ p_{\text{CV}}^{k+1} = M v_z(t_{k+1}) \\ p_{\perp}^{k+1} = p_{\perp}^k - \frac{\Delta t}{2} (\nabla_{\perp} V(q^k) + \nabla_{\perp} V(q^{k+1})). \end{cases} \quad (\text{D.5})$$

This corresponds to the deterministic dynamics for the CV coordinate and Verlet integration for the orthogonal coordinates. The associated work is simply

$$\begin{aligned} \mathcal{W}^{K_T} &= \sum_{k=0}^{K_T-1} (H(q^{k+1}, p^{k+1}) - H(q^k, p^k)) \\ &= H(q^{K_T}, p^{K_T}) - H(q^0, p^0). \end{aligned}$$

For the overall MCMC algorithm defined earlier, this means that the proposal $\tilde{Q}_{n+1} = q^{K_T}$ is accepted with probability the minimum of 1 and

$$\frac{e^{-\beta H(q^{K_T}, p^{K_T})} \rho(\tilde{Z}_{n+1}, Z_n)}{e^{-\beta H(q^0, p^0)} \rho(Z_n, \tilde{Z}_{n+1})} = \frac{e^{-\beta V(q^{K_T})} e^{-\beta \|p^{K_T}\|^2/(2M)} \rho(\tilde{Z}_{n+1}, Z_n)}{e^{-\beta V(q^0)} e^{-\beta \|p^0\|^2/(2M)} \rho(Z_n, \tilde{Z}_{n+1})} = \frac{e^{-\beta V(q^{K_T})} e^{-\beta \|p_{\perp}^{K_T}\|^2/(2M)} \rho(\tilde{Z}_{n+1}, Z_n)}{e^{-\beta V(q^0)} e^{-\beta \|p_{\perp}^0\|^2/(2M)} \rho(Z_n, \tilde{Z}_{n+1})}$$

The last equality, where the norm of the momentum is only considered on the orthogonal degrees of freedom, comes from the assumption that the initial and final velocities in CV space are equal. This is the relaxed version of (II.6) when there is no Fixman term to be considered, see Theorem II.3.

Appendix E: Details of the wavelet basis for the ϕ^4 model

As mentioned in Section IIIB, we use a basis of Haar-wavelets to represent the state of the ϕ^4 model in our simulation. For an introduction to wavelet transformations we refer the reader to Refs. 52 and 53, for example. Here, we only give a very practical definition. For a one dimensional input signal $\{\phi_i\}_{i=1}^N$ of size $N = 2^n$, set $\varphi_i^{(0)} = \phi_i$ for $i \in \{1, \dots, N\}$ and repeatedly apply the following transformation n times for $j \in \{1, \dots, n-1\}$:

$$\varphi_i^{(j+1)} = \frac{\varphi_{2i+1}^{(j)} + \varphi_{2i}^{(j)}}{\sqrt{2}}, \quad \bar{\varphi}_i^{(j+1)} = \frac{\varphi_{2i+1}^{(j)} - \varphi_{2i}^{(j)}}{\sqrt{2}}$$

with $i \in \{1, \dots, 2^{n-j}\}$. This transformation constitutes a bijective linear map between the input $\{\phi_i\}_{i=1}^N$ and the wavelet fields $\varphi^{(n)}$ and $\{\bar{\varphi}^j\}_{j=1}^n$. The coarsest-scale wavelet field has a single component entry:

$$\varphi^{(n)} = \frac{1}{(\sqrt{2})^n} \sum_{i=1}^N \phi_i = \frac{1}{\sqrt{N}} \sum_{i=1}^N \phi_i.$$

For the ϕ^4 model, since $\varphi^{(n)} = \sqrt{N} \cdot \bar{\phi}$, this corresponds to the magnetization $\bar{\phi}$ up to a prefactor. In practice, we therefore run the dynamics (D.1)-(D.3) with $q_{\text{CV}} = \bar{\phi}$ and $q_{\perp}$ the remaining wavelet components $\{\bar{\varphi}^j\}_{j=1}^n$.

Appendix F: Details of the dimer model

1. Computation of the Fixman term

Recall that we use the reaction coordinate defined in (III.6):

$$\xi(q) = \frac{|q_1 - q_2| - r_0}{2w}.$$

We write the gradient of the reaction coordinate $\nabla \xi(q) \in \mathbb{R}^{d \times 1}$ in the form

$$\nabla \xi(q) = \frac{1}{2w} \begin{pmatrix} \frac{q_1 - q_2}{|q_1 - q_2|} \\ -\frac{q_1 - q_2}{|q_1 - q_2|} \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \frac{1}{2w} \begin{pmatrix} e_{12} \\ -e_{12} \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

with the normalized vector $e_{12} = \frac{q_1 - q_2}{|q_1 - q_2|} \in \mathbb{R}^2$. The Gram tensor is then given by

$$G_M(q) = \nabla \xi(q)^\top M^{-1} \nabla \xi(q) = \frac{1}{M} |\nabla \xi(q)|^2 = \frac{1}{2Mw^2},$$

leading to a Fixman term (defined in (II.16)) that is constant:

$$V_{\text{fix}}(q) = \frac{1}{2\beta} \log \det G_M(q) = -\frac{1}{2\beta} \log(2w^2 M).$$

2. Lagrange multiplier for position constraint

From (II.18), the enforcement of the position constraint can be rewritten as

$$\begin{cases} q^{k+1} = \tilde{q}^{k+1} + \Delta t M^{-1} \nabla \xi(q^k) \lambda^{k+1/2}, \\ \xi(q^{k+1}) = z(t_{k+1}), \end{cases} \quad (C_q)$$

with $\tilde{q}^{k+1} = q^k + \Delta t M^{-1} p^{k+1/4} - M^{-1} \frac{\Delta t^2}{2} \nabla \tilde{V}(q^k)$. Inserting the first equation into the second and using the definition (III.6) of ξ leads to a quadratic equation for $\lambda^{k+1/2}$ with the solution

$$\lambda_{\pm}^{k+1/2} = \frac{wM}{\Delta t} \left(-e_{12}^k \cdot (\tilde{q}_1^{k+1} - \tilde{q}_2^{k+1}) \pm \sqrt{\Delta} \right),$$

where $e_{12}^k = \frac{q_1^k - q_2^k}{|q_1^k - q_2^k|} \in \mathbb{R}^2$ is a normalized vector and the discriminant is given by

$$\Delta = [e_{12}^k \cdot (\tilde{q}_1^{k+1} - \tilde{q}_2^{k+1})]^2 - [|\tilde{q}_1^{k+1} - \tilde{q}_2^{k+1}|^2 - (2wz(t_{k+1}) + r_0)^2].$$

We choose the smallest value for $\lambda^{k+1/2}$, which corresponds to choosing $\lambda_+^{k+1/2}$ if $e_{12}^k \cdot (\tilde{q}_1^{k+1} - \tilde{q}_2^{k+1}) > 0$ and $\lambda_-^{k+1/2}$ in the opposite case. This choice is consistent with ensuring that the value of the Lagrange multiplier is of order Δt for small timesteps.

3. Free Dimer not interacting with solvent particles

Considering the dimer model and the collective variable introduced in Section III C, the free energy can be computed analytically for the special case when the solvent particles do not interact with the dimer. We have

$$\begin{aligned} F(z) &= -\frac{1}{\beta} \log \int e^{-\beta V(q)} \delta_{\xi(q)-z}(\mathrm{d}q) \\ &= -\frac{1}{\beta} \log \int e^{-\beta V(q)} |\nabla \xi(q)|^{-1} \sigma_{\Sigma(z)}^M(\mathrm{d}q) \\ &= -\frac{1}{\beta} \log \left(e^{-\beta V_D(2wz+r_0)} (2wz + r_0) \right) + \text{const.} \\ &= V_D(2wz + r_0) - \frac{1}{\beta} \log(2wz + r_0) + \text{const.} \end{aligned}$$

where we ignored constants that just shift the free energy. For the second equality, consider that if $q_1 = 0$ is fixed, the values of q_2 are constrained to a circle of radius $r = 2wz + r_0$, which introduces a Jacobian term.

Appendix G: Details of the polymer model

We simulate the polymer system introduced in Section III D at a temperature $1/\beta = 1.3$ using the jaxMD package⁵⁴. The system is contained in a three-dimensional box of length $L = 7.47$ with periodic boundaries, with $N_{\text{solvent}} = 200$ solvent particles, which yields a solvent density of $N_{\text{solvent}}/L^3 = 0.48$ particles per unit volume. The parameters of the polymer potentials are $d_1 = 1$, $d_2 = 1.63$ (to fix the bonding angle at 109°), $k_1 = k_2 = 500$, $d_e = 4.62$, $a_e = 0.8$ and $k_e = 9.1$, corresponding to an energy barrier of $7\beta^{-1}$. The Lennard-Jones potential is essentially given by

$$V_{\text{LJ}} = 4\epsilon \left[\left(\frac{r^*}{r} \right)^{12} - \left(\frac{r^*}{r} \right)^6 \right],$$

with $r^* = \epsilon = 1$, and thus very similar to the WCA potential used for the dimer. In practice, however, we use a modified version given by jaxMD that smoothly decays to zero between an onset radius $r_{\text{onset}} = 2.0$ and a cutoff radius $r_{\text{cutoff}} = 2.5$.

To implement **Step 1** of the algorithm in Section II B, we use an independent proposal sampler in CV space for both choices of CV considered here. For the first CV, i.e. the one-dimensional end-to-end distance, the proposal sampler is given by a Gaussian mixture model with two modes of equal weight and width $\sigma = 0.17$ at positions $z = 3.53$ and $z = 5.65$. For the second, 27-dimensional CV, i.e. the Cartesian coordinates of the polymer beads, we use a normalizing flow to propose distances between the polymer beads. These distances are the eight nearest neighbor and seven second-nearest neighbor distances as well as the six distances between the first bead and beads number 4 to 9, adding up to 21 distances in total. To complete such a proposed set of distances to a full proposal of polymer coordinates, one needs to additionally specify the global translation and rotation, as well as six sign values $\eta_i \in \{\pm 1\}$, $i \in \{1, \dots, 6\}$ (explained in the following). The global translation is fixed by always positioning the central

bead of the polymer at the center of the box (which we do throughout the simulation) and the global rotation is sampled uniformly. The necessity of the six sign values η_i can be understood from the triangulation process used to determine the Cartesian polymer bead positions. Assuming one has already placed three beads with the desired distances, there will be two possible solutions (if there is a solution) to place the fourth bead at the right distance to the previous three (two intersection points of three spheres), which corresponds to choosing the sign of the dihedral angle. For a given set of 21 distances as in our case, there are six such sign values to choose, and we sample them uniformly at random. We compute the Jacobian of this reconstruction using automatic differentiation to arrive at a probability measure in the CV space of polymer positions $z \in \mathbb{R}^{27}$. The flow will sometimes produce sets of distances for which no solution to the triangulation problem exists and these samples are then directly rejected. Since this restriction only changes the normalization of the flow proposal measure, it does not enter into the overall formula of the accept/reject probability in **Step 3** of the algorithm in Section II B. The flow is implemented using the FlowJax package⁵⁵, with one affine whitening layer and 20 blocks of RationalQuadraticSpline transformations⁵⁶ with 32 knots on the interval $[-4, 4]$ and internal neural networks of depth 2 and width 30 (see the reference for further explanations). It was trained on a large dataset of one million samples obtained from a long Langevin simulation over 200 epochs with a batch size of 40,000 and a learning rate of 10^{-3} using the Adam optimizer with otherwise default parameters.

Appendix H: Additional numerical results

1. Gaussian tunnel

To illustrate the optimal choice of parameters when running the algorithm on the example of the Gaussian tunnel from Section III A, we show the inverse mode jump cost (introduced in the main text) for different parameter values in Fig. 13. The optimal value of α_2 is roughly the same for all choices of α_1 , whereas the optimal number of intermediate steps K_T increases with α_1 . Clearly, the optimal performance is observed for the deterministic dynamics, namely $\alpha_1 = 0$.




Figure 13. Inverse mode jump cost when running the algorithm on the example of the Gaussian tunnel introduced in Section III A for different parameter values α_1, α_2 (introduced in Section II D). The number $K_T(b)$ represents the number of steps for a jump of distance b and is related to the inverse of the dimensionless velocity $\tilde{v}$ via $K_T(b) = \text{ceil}(b/\tilde{v})$. White space indicates that at least one chain never switched mode.

For the deterministic case, there is a simple continuous-time limit of $\Delta t \rightarrow 0$, in our parameterization corresponding to $\alpha_2 \rightarrow 0$, where the only relevant quantity determining the acceptance rate is the normalized transition time $T/\sqrt{\beta M}$. This is indeed what we observe in Fig. 14, where for small enough α_2 , the acceptance rate only depends on this physical time scale. For numerical performance, however, the relevant quantity is of course the acceptance rate normalized by the numerical cost.

2. Gaussian tunnel with a non-linear collective variable

We consider here a version of the Gaussian tunnel introduced in Section III A in dimension $d = 10$ with a non-linear collective variable. With the same underlying probability distribution, we consider the CV $\xi(z) = \tanh\left(\frac{z}{b}\right) \cdot \frac{b}{\tanh(1)}$. Using the push-forward of the probability density $\nu_{\text{CV}}(z)$ under this transformation as the proposal distribution (and thus exactly the marginal of the target distribution), we show in Fig. 15 how our algorithm with the schedule (II.11)




Figure 14. Acceptance rate for a fixed jump from $z = 0$ to $\tilde{z} = b$ for the Gaussian tunnel with the deterministic dynamics $\alpha_1 = 0$. The results shown here are the same as in the top left panel of Fig. 1 with a different choice of axes. The right panel shows all simulations with $\alpha_2 < 0.4$ combined in one plot.

leads to unbiased results, whereas a linear schedule that violates Eq. (II.6) or the omission of the Fixman term both lead to a bias. To test an inappropriate schedule, we simply use one with constant velocity, i.e. (II.9) and (II.10) with $f(\tau) = \tau$. The Fixman term is given by $V_{\text{fix}}(z, x^\perp) = \frac{1}{\beta} \log(\xi'(z))$ and just depends on z . We observe that upon omission of the Fixman term, the resulting distribution is indeed biased: its marginal along ξ is proportional to $\nu_{\text{CV}}(z) \cdot e^{\beta V_{\text{fix}}(z)}$.




Figure 15. Simulation of the Gaussian tunnel example introduced in Section III A (with smaller dimension $d = 10$) using steered moves in a non-linear CV, $\xi(z) = \tanh\left(\frac{z}{b}\right) \cdot \frac{b}{\tanh(1)}$. We show the results for an unbiased algorithm that follows the method specified in Section II, including the Fixman term (II.16) and using the cosine-schedule (II.11). For comparison, we also show results for modified algorithms where (i) a linear schedule with constant velocity was used (violating Eq. (II.6)) and (ii) the Fixman term was omitted. The proposal distribution in CV-space is exactly the target distribution, histograms are shown over 20 chains and 10,000 iterations with $\alpha_1 = 0$, $\alpha_2 = 0.67$ and inverse velocity $1/\tilde{v} = 2.5$, corresponding to $K = 25$ steps for a jump between the two mode centers.

3. ϕ^4 Model

To complement the results shown in Fig. 4 we ran the algorithm on the ϕ^4 model from Section IIIB for additional values of α_1 and show the associated mode-jump cost in Fig. 16.




Figure 16. Mode jump cost for the ϕ^4 model from Section IIIB for different parameters, estimated from running 20 MCMC chains over 10,000 iterations. The x -axis shows the number of steps for a steered schedule over the distance $2\bar{\phi}$ of the two modes

4. Dimer in a solvent

In addition to Fig. 8, we ran the algorithm on the dimer model for additional values of α_1 . The associated inverse mode-jump cost is shown in Fig. 17. Apart from the deterministic algorithm ($\alpha_1 = 0$), almost no mode switches were observed within the computational budget.




Figure 17. Mode jump cost for the Dimer model from Section IIIC for different parameters, estimated from running 20 MCMC chains over 5,000 iterations. The inverse dimensionless velocity $1/\tilde{v}$ corresponds to the number of steps for a steered schedule over a unit distance. White space indicates that at least one chain never switched mode during observation time. The left-most panel shows the same results as Fig. 8 (with a different color scale).