

Lectures on Optics

“To Understand the universe we must understand the light”

Sunday 30th August, 2026



LECTURES ON OPTICS

Jafert Serrano

Professor: Rafael Ángel Torres Amaris

*Universidad Industrial de Santander
Sciences faculty
Physics school*

Group of optics and signals treatment

Sunday 30th August, 2026

Author: Jafert Serrano UIS

Contents

1	The loving Story of Optics	3
1.1	Optics from Euclid to Huygens	4
1.1.1	Ancient times	4
1.1.2	The arabain period	7
1.1.3	The middle ages	8
1.1.4	The renaissance	9
1.1.5	Newton and Huygens	13
2	Geometrical Optics	15
2.1	The postulates of geometrical optics and the dark Room	16
2.2	The Snell-Descartes law and the Fermat's principle	23
2.2.1	The critical angle, and the refraction by a prism	26
2.3	The Descartes's ovoid and diopters	28
2.4	The spherical diopter	35
2.4.1	The Young-Weirstrass points	36
2.4.2	The second spherical diopter	37
2.5	Objects and images to the infinity	38
2.5.1	Revolution Ellipsoid	39
2.5.2	Revolution Hyperboloid	41
2.6	Rigourously stigmatic surfaces by reflection	44
2.6.1	The ellipsoidal mirror	46
2.6.2	The hyperboloidal mirror	47
2.6.3	Special cases: the spherical and the plane mirror	48
2.6.4	The parabolic mirror and objects at infinity	49
2.7	Explicit formulation of the Descartes ovoids	50
2.7.1	The Descartes ovoid as a superconic	51
2.7.2	The explicit formulation	54
2.7.2.1	The Schwarzschild equation	57

2.8	Special Descartes ovoids	57
2.8.1	Spherovoids, $G = 0$	58
2.8.2	Parovoids, $G = -1$	58
2.8.3	Hyperovoids, $G < -1$	58
2.8.4	Prolate and oblate ellipsovoids	59
2.9	Descartes ovoids and aspheric surfaces	59
2.10	Approximate stigmatism in the spherical diopter	60
2.10.1	Gauss optics, the paraxial approximation	62
2.10.2	Spherical aberration	64
2.10.2.1	The wavefront aberration and the quartic wavefront	66
2.10.2.2	The Seidel aberrations	67
2.10.2.3	Aberrations close to the paraxial condition	69
2.10.3	Focal length for marginal rays	70
2.10.3.1	Longitudinal object defocus	71
2.10.3.2	Image formation in the marginal ray regime	73
2.10.4	The thin lens with aberrations	74
2.11	Aplanatism	77
2.11.1	The Abbe sine condition	78
3	Electromagnetic Optics and Polarisation	81
3.1	A historical context of electromagnetic optics and polarisation	82
3.2	The vector behavior of the electromagnetic radiation	83
3.3	Maxwell equations are the Schrödinger equation for photons	85
3.4	Properties of the electromagnetic waves	88
3.4.1	Principal solutions of the electric wave equation	89
3.4.2	Beams with a particular phase and structured light	92
3.4.3	From 3D wavefronts to 2D wavefronts	95
3.5	Light's polarisation	96
3.5.1	Monochromatic and deterministic formulation: Jones spinors and the polarization ellipse	97
3.5.1.1	How to build the polarisation ellipse: Phasors	98
3.5.1.2	How to build the polarisation ellipse: Ellipse angles	100

3.5.1.3	The canonical polarisation states: horizontal, vertical, diagonal and circular	102
3.5.2	Time evolution of the polarisations: Stokes parameters and the Poincaré sphere	105
3.5.2.1	Why the Stokes vector is not a vector	111
3.5.3	The bridge between Jones and Stokes: the stereographic projection	113
3.5.3.1	What the stereographic projection is	114
3.5.3.2	Why a polarisation state is a single complex number	115
3.5.3.3	The Poincaré sphere is that sphere	115
3.5.3.4	A flat atlas of every polarisation state	117
3.5.4	Transformations of Polarisation: Jones matrices	118
3.5.4.1	Retarders	119
3.5.4.2	Rotators	122
3.5.4.3	Dichroics and polarisers	122
3.5.5	Transformations of Polarisation: Mueller matrices	125
3.5.5.1	From the Jones matrix to the coherence matrix	125
3.5.5.2	The Mueller matrix	126
3.5.5.3	Retarders are rotations of the Poincaré sphere	127
3.5.5.4	Polarisers, dichroics and the Malus law	129
3.5.5.5	What no Jones matrix can do: depolarisation	131
3.5.5.6	Why this matters	132
3.5.6	The Jones theorems	132
3.5.6.1	Theorem I: every system has an eigenpolarisation	133
3.5.6.2	Theorem II: retarders and rotators	134
3.5.6.3	Theorem III: partial polarisers and rotators	135
3.5.6.4	Theorem IV: the general equivalence theorem	136
3.5.6.5	Theorem V: reversibility	137
3.5.6.6	What the theorems say together	138
3.5.7	The elliptic general Malus law	139
3.5.8	The law of elliptic birefringents	139

4	Interference and Interferometers	141
4.1	A historical context of interference and interferometers	142
4.2	How does interference occur?	143
4.2.1	Interference of plane waves	145
4.2.2	Interference of spherical waves	146
4.3	Interferometers	151
4.3.1	Young interferometer	151
4.3.2	Fresnel biprism	155
4.3.3	Michelson and Morley interferometer	161
5	Theory of Diffraction	163
5.1	A historical context of diffraction theory: from shadows to the vector world	164
5.2	Scalar theory of diffraction	167
5.3	The scalar wave equation (Helmholtz equation)	169
5.4	The Kirchhoff scalar boundary-value problem	169
5.4.1	Green's theorem	170
5.4.2	The HelmholtzKirchhoff integral theorem	171
5.4.3	The RayleighSommerfeld formulation	175
5.5	The Fresnel approximation	176
5.5.1	Example: Fraunhofer diffraction between two confocal spheres	177
5.6	The Fraunhofer approximation	180
5.7	The fractional Fourier transformation	182
5.7.1	A historical context of Fourier analysis and of its fractional powers	182
5.7.2	The fractional Fourier transformation as a differential and as an integral operator	186
5.7.2.1	The eigenfunctions of the Fourier transformation	187
5.7.2.2	Namias' definition and the differential operator	188
5.7.2.3	The integral form of the transformation	190
5.7.2.4	The inverse transformation and the index law	192
5.7.2.5	The regularisation of McBride and Kerr	193
5.7.2.6	Why the construction is useful	194
5.7.3	The role of Fourier analysis in the theory of diffraction	195

5.7.3.1	Why a plane cannot be the right reference surface	196
5.7.3.2	Reference spheres and the fractional order	197
5.7.3.3	The normalising constant and the final result	199
5.7.3.4	Physical meaning of a real order	201
5.7.3.5	Physical meaning of a complex order	202
6	Theory of Coherence	205
6.1	A historical context of coherence theory	206
6.2	How is statistical similarity quantified?	209
6.3	Random variables	212
6.3.1	The sample space and the physical meaning of an event	213
6.3.2	The sigma algebra, or which questions may be asked	213
6.3.3	Measure, the Lebesgue measure and the probability measure	215
6.3.4	Random variables and the meaning of measurability	216
6.3.5	Stochastic processes	218
6.3.6	The probability space of the electromagnetic field	218
6.3.7	Stationarity, ergodicity and the right to measure	220
6.4	The coherence function	221
6.4.1	Manifestations of coherence: the WienerKinchin theorem	225
6.4.2	The wave equation of coherence	227
6.4.3	Propagation of coherence and the direction of time: the van CittertZernike theorem	230

List of Figures

2.1	Schematic representation of the Fizeau's experiment pretended to measure the light's speed.	18
2.2	Graphical representation of a generic dark room which is a box of an arbitrary material which has a side opened and while it satisfies the Rayleigh's inequality i will form an inverted image.	20
2.3	A schematic representation of a ray beam from A to B which goes through n different transparent media and in each one it had a different speed and traveled different length.	22
2.4	two media separated by a diopter which an observer O measures the path the light will follow from A to B and proposes a displacement of the source A to A' to see how must be the light path.	24
2.5	Representation of a cylindrical ray light being reflected by a perfectly polished mirror.	26
2.6	Representation of a cylindrical bunch of rays being reflected by an inhomogeneous surface.	27
2.7	Representation of all the incident and refracted rays from a media with refraction index n_1 to another with refraction index n_2 such that $n_1 > n_2$. The angle θ_2 here shown wont never reach a value equal to $asin\left(\frac{n_1}{n_2}\right)$	28
2.8	Representation of all the incident and refracted rays from a media with refraction index n_1 to another with refraction index n_2 such that $n_1 < n_2$. The angle θ_c here separates the rays that will be refracted , (angles less than θ_c), and the rays that will be totally reflected internally.	29
2.9	Representation of two optical systems where appear real object points, real and virtual image points.	30
2.10	Optical system $\mathcal{S}$ formed by two diopters $\mathcal{S}_1, \mathcal{S}_2$ where the point A_o is a virtual object and A_i is a virtual image point.	30
2.11	Optical system $\mathcal{S}$ formed by three optical systems $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ where the point A_o is a real object point, A' is a virtual image point for $\mathcal{S}_1$, but a real object point for $\mathcal{S}_2, \mathcal{S}_3$, A'' is a virtual image for the real object A' and A_i is the real image of the virtual object A''	31
2.12	The stigmatic system transform the incident wavefront $\mathcal{A}$ to another new spherical wavefront $\mathcal{A}'$	32
2.13	Scheme representation of the development of a rigorously stigmatic system, from a point source A_o at a position $[0, 0, z_o]$ the diopter with a vertice v forms a perfect image A_i at $[0, 0, z_i]$	33

2.14	The spherical Diopter for the first case, where the object point is real and it is outside the sphere but the image point is virtual and inside the diopter.	38
2.15	Prolate Ellipsoid centered and $z = B$	40
2.16	Ellipsoid diopter when $n_i > n_o$	41
2.17	Ellipsoid diopter when $n_i > n_o$ and $z_i < 0$	42
2.18	Geometrical representation of the two leaf hyperboloid with the parameters a, b and foci F_1 and F_2	43
2.19	Geometrical representation of the first case of a two leaf hyperboloid.	44
2.20	Geometrical representation of the first case of a two leaf hyperboloid.	45
2.21	Ellipsoidal mirror. The object F_1 and the image F_2 coincide with the two foci of the ellipse, so every ray leaving one focus is reflected exactly onto the other.	46
2.22	Hyperboloid mirror. The rays from the object F_1 diverge as they were emitted by a virtual image located in F_2	48
2.23	Parabolic mirror obtained as the limit $z_o \rightarrow -\infty$. The rays coming parallel to the axis from an object at infinity are all focused onto the single point $z = z_i$	50
2.24	Rigorously stigmatic quartic surface described in the meridional plane by the polar parameters ρ and φ measured from the vertex $\mathcal{O}$. The object point A_o at z_o and the image point A_i at z_i are the strictly stigmatic pair, and at the incidence point I the vectorial Snell-Descartes law $n_o \hat{u} - n_i \hat{u}' = \kappa \hat{N}$ is satisfied.	53
2.25	Meridional sections of Cartesian surfaces with the refraction indices fixed at $n_o = 1$ and $n_i = 1.7$ while the object and image positions are varied. The solid curves are the solutions with the minus sign in 2.96, which pass through the origin and are the Descartes ovoids, and the dashed curves are the solutions with the plus sign, the outer sheet of the same quartic. (a) $z_o = -10$ and $z_i = 20$. (b) $z_o = -10$ and $z_i = -20$. (c) $z_o = 10$ and $z_i = 20$. (d) $z_o = 10$ and $z_i = -20$	55
2.26	Degenerate cases of refractive and reflective rigorously stigmatic surfaces according to the refraction indices and the object and image positions. For each combination of these parameters the values of the shape parameters $GOTS$ of the corresponding conic of revolution are obtained. Note that $T = S = 0$ in every conic case, and that the reflective surfaces are recovered by the substitution $n_i \rightarrow -n_o$	56
2.27	Image formation by a spherical diopter $\mathcal{D}$ of radius R , vertex V and centre of curvature C . The object O is at a distance $d = z_o$ from the vertex and the image O' at $d' = z_i$, and the ray reaches the surface at the point I , which is located by the angle φ measured at the centre of curvature. The chord $\rho = \overline{VI}$, the transverse height h and the sagitta η are the three geometrical quantities that will organise the whole calculation.	61

2.28	Quartic surface Q of the wavefront emerging from a spherical refracting surface. The marginal rays converge at A' instead of the paraxial point A_p , because the quartic wavefront Q departs from the Gaussian spherical reference $\mathcal{A}'$ by the amount $\Delta l = W(\vec{\rho})$, and the envelope of the refracted rays forms the caustic.	67
2.29	Illustration of the longitudinal spherical aberration. The rays arriving parallel to the axis from an object at infinity are focused at f_{M+} if they are marginal and at f_{P+} if they are paraxial, and the separation between the two is Δf_+ . The envelope of the refracted rays is the caustic.	72
2.30	Illustration of the defocus in the object space. The rays that emerge parallel to the axis in the image space come, for the marginal zone, from f_{M-} and, for the paraxial zone, from f_{P-} , and the separation between the two is Δf_-	73
2.31	Image formation systems. Each family of rays alone forms a sharp image, and it is only their superposition that produces the aberrated one.	74
2.32	Thin lens with aberrations, formed by two spherical diopters of radii R_1 and R_2 separating the medium n from the glass n' . The thickness Δ between the two vertices is taken to zero at the end of the calculation.	75
2.33	The aplanatism condition. Two pairs of rigorously stigmatic points (A_1, A'_1) and (A_2, A'_2) are considered, with A'_1 in a neighbourhood of A_1 and A'_2 in a neighbourhood of A_2 , the diopter D separating the media n_1 and n_2 , and $\Delta \vec{l}$ being the displacement of the incidence point on the surface. The Spanish label <i>vecindad</i> in the figure means neighbourhood.	77
2.34	Geometry of the Abbe sine condition. The transverse object $\overline{A_1 A'_1}$ in the medium n_1 is imaged as $\overline{A_2 A'_2}$ in the medium n_2 , and θ_1 and θ_2 are the angles that a given ray makes with the axis in the object and image spaces.	79
3.1	A source $\mathcal{S}$ produces an arbitrary kind of wave which propagates in the space-time, then, for any given time and position we can take an arbitrary point p and, due to the smooth manifold nature of this wave, there exists a neighborhood N of p where the wavefront is plane.	96
3.2	The polarisation ellipse can be reconstructed taking all the possible rotations of the two phasors (which corresponds to the componentes of the Jones spinor).	99
3.3	The general form of the polarisation ellipse can be described with only two angles; the orientation angle α and the ellipticity angle χ	100
3.4	The polarisation ellipse is completely unrotated due to the relative phases between the componentes of the electric field.	101
3.5	In the Poincaré any polarisation state is a point on the surface on the sphere, however if the state is partially polarized it would be a point inside the poincaré sphere whose distance from the center is $\sqrt{S_1^2 + S_2^2 + S_3^2}$	111

- 3.6 The stereographic projection. A ray leaving the centre of projection N and passing through a point P of the sphere S^2 meets the plane $\mathbb{R}^2$ at a single point Q . Points near the contact point fall near the origin, and points near N fly off towards infinity. 114
- 3.7 The Poincaré sphere resting on the complex plane of the ratio w . Horizontal polarisation sits at the point of contact, $w = 0$, vertical polarisation sits at the centre of projection and is sent to the point at infinity, and the equator, where the two amplitudes are equal, is mapped onto the unit circle $|w| = 1$. The angles 2α and 2χ that locate a state P on the sphere are twice the orientation and ellipticity angles of its ellipse. 116
- 3.8 The flattened Poincaré sphere. Each point of the complex plane of w carries the ellipse of the state it represents. The origin is horizontal polarisation, the real axis carries every linear state, the imaginary axis carries the ellipses aligned with the coordinate axes together with the two circular states, and every direction of escape to infinity leads to one and the same state, the vertical one. 118
- 3.9 A quarter wave plate retarder is a retarder with $\delta = \pi/2$ and when it has got its fast axis oriented with the horizontal one it transforms the polarization in a very specific way. The $\pm 45^\circ$ diagonal states transform into circular polarization states, and the circular polarization states transform into $\pm 45^\circ$ diagonal states. 120
- 3.10 A linear polariser with its polariser axis oriented parallel to the horizontal axis projects all the incident states to horizontal ones, however the transmitted intensity is reduced by the relative angle between the incident state $|\alpha; \chi\rangle$ and the polariser axis α_{pol} in this case $\alpha_{pol} = 0^\circ$ 124
- 3.11 An initial state $|\alpha; \chi\rangle$ under the effect of a rotated retarder with retar δ and eigenstate r is rotated an angle of δ around the r -axis to a final state $|\alpha'; \chi'\rangle$ 129
- 4.1 Two plane waves with unitary number vector $\hat{k}_1, \hat{k}_2$ propagate and interfere producing a pattern of interference characterized by a vector $\vec{K}$ and an interfringe distance Λ . . 146
- 4.2 Two plane sources of spherical waves interfere in a point P located on the interference plane. The two sources are separated a distance a and the distance from both surface to the interference plane is D 147
- 4.3 Young interferometer illuminated by a spherical wave from a point source $\mathcal{S}$ placed off axis. The source to slit stage sets the amplitude and the phase reaching each slit $\mathcal{S}_1, \mathcal{S}_2$, and the slit to screen stage is the interference of two secondary spherical waves. 153
- 4.4 Method of images applied to the Fresnel biprism. Each half of the biprism produces a virtual point image of the source, $\mathcal{S}_1$ from the upper half and $\mathcal{S}_2$ from the lower half, separated by a and located in the same transverse plane as $\mathcal{S}$. Beyond the biprism the two beams overlap only inside the shaded strip of width W on the screen. 157

4.5	General configuration of the Michelson interferometer. A source placed a distance a from the beamsplitting cube illuminates it, and the beam is divided along two perpendicular arms. In arm 1 the light travels a total distance $d + \ell_1$ before striking mirror 1, tilted an angle θ away from the orientation that would send the beam straight back along itself; in arm 2 it travels a total distance $d + \ell_2$ before striking mirror 2, tilted an angle ϕ in the same sense. The two reflected beams return to the beamsplitter and recombine there, ready to interfere.	161
5.1	On the boundary $\Sigma(V)$ the contribution of the field components $U(\vec{r})$ is shown; inside the region the total field is built from the contribution of all the points on the boundary. One may add infinitely many point Green charges of arbitrary propagation nature; the condition for the invariance of equation (5.20) is that these image charges lie outside $\Sigma(V)$	172
5.2	Representation of the Green's functions for point sources. Under these boundary conditions the Green's functions behave as point charges, one the mirror image of the other.	173
5.3	Representation of the Green's functions for non-point sources. Under these boundary conditions the Green's functions still behave as mirror images, but now with given volume and geometry; the only constraint on the emission of these sources is that, on the boundary $z = 0$, the total field must vanish, as required by the assumed Dirichlet conditions.	174
5.4	Geometry of the two confocal spheres. The cap $\mathcal{A}$ is tangent to the emitting plane $\mathcal{P}$ at the vertex O and has its centre of curvature at O' , while the cap $\mathcal{F}$ is tangent to the receiving plane $\mathcal{Q}$ at O' and has its centre of curvature at O , so that both radii equal the propagation distance z . Read on these two surfaces, the Fresnel propagation reduces to an exact Fourier transformation, which in the language of the fractional Fourier transformation of Section 5.7 is the order $\alpha = \pi/2$	179
5.5	Scheme of the confocal cardinal spheres, with radii equal to the propagation distance z of the field.	181
5.6	The two cardinal spheres of radius R associated with a propagation over the distance z . The emitting cap $\mathcal{A}$ is tangent to its plane at $V_{\mathcal{A}}$ with its centre of curvature downstream at $C_{\mathcal{A}}$, of abscissa R , and the receiving cap $\mathcal{B}$ is tangent to its plane at $V_{\mathcal{B}}$ with its centre upstream at $C_{\mathcal{B}}$, of abscissa $z - R$, so that the two caps bow towards each other. The quantity δ is the sagitta (5.90). The confocal case of Figure 5.4 is recovered when $R = z$, since then $C_{\mathcal{A}}$ falls on $V_{\mathcal{B}}$ and $C_{\mathcal{B}}$ on $V_{\mathcal{A}}$	198
6.1	Representation of $E(\vec{r}, t)$ as a random variable.	222
6.2	Graphical representation of the narrowband coherence function. In black, the envelope $\tilde{\Gamma}(\vec{r}, \tau)$ is shown, and in red the oscillations $e^{i2\pi\tilde{\nu}\tau}$. These oscillations are confined within something known as the coherence time τ_C , which we shall see later.	228

6.3	Propagation of the electric field from a plane $Z = 0$ to a plane Z , where each point of the wavefront carries its mean polarisation ellipse.	230
6.4	Propagation of the coherence function between two points $\vec{r}, \vec{r}'$ on the initial plane to the coherence between two points $\vec{r}_1, \vec{r}_2$ on the observation plane.	231
6.5	Evolution of the realisations upon propagating the electric field.	233
6.6	The mutual coherence propagated to the far field is the Fourier transform of the intensity distribution $\Gamma_0(\vec{S})$ of the source.	235

Abstract

The field of optics is a captivating and continually evolving branch of science that plays an indispensable role in our modern world. "Advances in Optics: Exploring the Boundaries of Light and Vision" presents a comprehensive overview of the latest developments, theories, and applications in optics, spanning from fundamental principles to cutting-edge technologies.

This book delves into the fundamental properties of light, offering a clear and concise explanation of wave-particle duality, the nature of electromagnetic waves, and the behavior of light in various media. It explores the principles of geometric optics, including reflection, refraction, and the formation of images, while also addressing wave optics phenomena such as interference, diffraction, and polarization.

In addition to the foundational aspects, "Advances in Optics" goes beyond traditional optics to cover emerging topics and technologies. Readers will find detailed discussions on advanced optical materials, including metamaterials and photonic crystals, and their revolutionary applications in the manipulation of light for imaging, communication, and sensing. Quantum optics, an exciting frontier in optical science, is also explored, with a focus on quantum entanglement, quantum cryptography, and quantum computing.

Moreover, this book highlights the practical applications of optics in various fields, including medicine, telecommunications, astronomy, and environmental monitoring. It provides insights into state-of-the-art optical instruments and techniques, such as lasers, optical fibers, and adaptive optics systems, and their pivotal roles in advancing technology and scientific discovery.

"Advances in Optics" is designed to cater to a broad readership, from students and educators seeking a comprehensive textbook to researchers and professionals looking for the latest developments in the field. Its accessible style, accompanied by illustrative examples and diagrams, ensures that both newcomers and experts can appreciate the wonders of optics and its ever-expanding frontiers.

As we venture further into the 21st century, the exploration of light and vision promises to yield groundbreaking discoveries and transformative technologies. "Advances in Optics" serves as an invaluable resource for anyone curious about the principles, innovations, and limitless potential of optics in shaping our future.

Chapter 1

The loving Story of Optics

“The light is the God’s shadow”
– Saint Agustin

When we study physics beyond doing long calculations, formulate complicated theories or doing extremely elaborated experiments in an advanced laboratory there is a really important fact many people ignore, the *story of physics* if we want to study physics we must have understand where was born the physics and more important, how was it done. Here we are going to check the story of physics from the geometrical optics, to corpuscular optics to wave optics, to electromagnetic optics and the most recent the quantum optics.

1.1 Optics from Euclid to Huygens

Here I'm going to copy and paste one of the best articles I've read about the story of optics (Herzberger, 1966).

If I were endowed with dictatorial powers, I would require everyone receiving a degree in a scientific subject to know its history and to have read the classical papers relating to it. Historical knowledge is important because it stimulates creative thinking. The man who first struggled with an idea, trying to find a law, looked at the situation with different eyes than do we who accept the law as a matter of course. He considered alternatives to the law, and different interpretations, and some of these alternatives and interpretations may still be stimulating and worth thinking about.

I was fortunate during the eight years of my affiliation with the Zeiss works at the beginning of my career to have access to their extensive library; it contained the originals, or at least translations, of nearly all the writings in optics from Euclid to contemporary times. Over thirty years ago, I wrote a historical survey describing the content of important papers on geometrical optics. This task was immensely facilitated by the extensive bibliography in the third edition of Czapski-Eppenstein, to which I had contributed my small part as a work student at the firm of Zeiss. Lately, I have looked through most of these papers again, as far as they were available to me, and have discovered many important details that I had not understood at that time. I should like to share these with you. I emphasize that my interest is not basically historical. I am not interested so much in questions of priority and originality of documents. My object is to entice you to read and to enjoy the old scriptures, and maybe take along an idea or two that might be of interest.

1.1.1 Ancient times

I am sure that the Egyptians and, even more, the Babylonians, with their great interest in astronomy, must have had some knowledge of optical laws. But the first important mention of one of the phenomena of refraction I found was in Plato's Republic. Plato (427-347 B.C.) quotes Socrates as saying (Jowett's translation): " . . . the same object appears straight when looked at out of the water, and crooked when in the water; and the concave becomes convex, owing to the illusion about colours to which the sight is liable. Thus every sort of confusion is revealed within us; and this is that weakness of the human mind on which the art of conjuring and of deceiving by light

and shadow and other ingenious devices imposes, having an effect upon us like magic." Our senses give an erroneous picture of reality; they are untrustworthy; alone can lead us to truth pure thinking. This idea persisted for two millennia under the influence of the misinterpreted teachings of Aristotle.

How we see is a question that long occupied the Greek philosophers, and, through them, the Neo-Platonists and the Scholastics. (Indeed, it cannot be said that the question is settled, and in recent times Ronchi has set forth some cogent ideas.) Even before Plato there was speculation in the writings of the Greek philosophers about the process of seeing. Most of the Greek thinkers believed that the eye sends out rays to the object, and Aristotle (384-322 B.C.) seems to have been the first to question this belief. If, says Aristotle, the eye sends out fiery rays that stream from it as from a lantern, why can we not see in the dark? His explanation, that the eye contains a transparent substance and that seeing is associated with the motion of something between the object and the eye, seems quite modern.

After the conquest of Greece by Alexander, Alexandria (in Egypt) and Sicily became the strongholds of Greek culture. The first great textbook of optics was written by Euclid of Alexandria, who lived about 300 B.C. And is better known for his Elements of Geometry. This famous work is still must reading for every scientist and is available in several editions. When King Archelaos once complained about the difficulty of learning mathematics and asked whether there was not an easier way, Euclid's answer is supposed to have been: "There is no royal route to geometry". When one of his pupils asked him what of material value could be gained by learning geometry, he told a slave, "Give the man threepence and tell him to go away".

Under Euclid's name we have two books of interest in the present connection: Optics and Catoptrics. The first deals with the laws of vision and perspective and the second with reflection. An English translation of the first was published in the Journal of the Optical Society of America twenty years ago. ⁷ It is a striking fact that one of the definitions in this book contains an idea that could be used to develop many problems of diffraction theory.

I quote some of Euclid's definitions, slightly paraphrased from the Optical Society of America translation: "Let it be assumed that lines drawn directly from the eye pass through a space of great extent [not infinite]; that the form of this space is the mantle of a cone with its apex in the eye and its base at the limits of our vision; that those things upon which the cone falls are seen and those things upon which the cone does not fall are not seen; and that things seen through several cones simultaneously appear to be more distinct." From this, Euclid concludes, among other things, that nothing can be seen in its entirety. The same objects located nearby are seen more distinctly than if they were at a distance. Every object that is visible has a certain limit of distance. If that is reached-that is, if the object lies entirely inside the mantle of the cone of vision-it is no longer seen. These are the axioms from which he derives the laws of perspective.

In Catoptrics the law of reflection is stated, namely, that incoming and outgoing rays form the

same angle with the surface normal. From this he derives the exact position of the image behind a plane mirror and the principle that left- and right-hand sides are interchanged. He generalizes these laws for spherical mirrors, both concave and convex, describing virtual images as well as stating that concave mirrors can give real images. As the position of the image is formed by reflection at curved surfaces, he suggests the point of intersection of the ray with the axis (shades of my definition of diapoint!). He shows the existence of a focal point for spherical mirrors, although he errs with respect to its position. One of his theorems says that an eye at the center of a spherical mirror sees only itself. Another states that you can light a fire at the focus of a concave mirror. He states that straight lines appear convex in a convex mirror. One can see the same object through many plane mirrors if they are positioned along the sides of a regular polygon with two more sides than there are mirrors.

According to Roman historians, the greatest mathematician of ancient times, Archimedes (287-212 B.C.), is supposed to have used mirrors to destroy the Roman fleet besieging Syracuse. Like most scientists, I have doubted this story, but a personal experience made me doubt my doubts.

Two years ago I had an optical accident. My wife had me get a large bottle of distilled water, and I left it in my auto in front of our laboratory. Driving home that evening I smelled smoke and stopped at a garage. After thoroughly investigating all the mechanical details and finding them in order, the mechanics discovered that smoke was coming out of the upholstery. The cylindrical bottle had acted like a lens and the sun had made a hole in the upholstery and ignited the stuffing inside. After this personal experience, I am somewhat more reluctant to doubt the story of Archimedes' mirrors.

The next person I should like to discuss, Hero of Alexandria, lived sometime between about 150 B.C. and A.D. 250. We have two outside dates for this philosopher: he mentions Hipparchos, who flourished between 146 B.C. and 126 B.C., but the mathematician Pappus (ca. A.D. 300) follows Hero's treatment in his own writings on mechanics. Be this as it may, Hero's *Catoptrics* is one of the most interesting books on optics. He is not content with Euclid's statement that light travels in straight lines, nor is he content with the form of the reflection law. He looks for a reason for both laws and derives it from the assumption that light chooses the shortest possible path between two points. We know that this minimum principle is not quite correct: the path is, in general, only an extremum. It may or may not be significant that Hero's book illustrates the concept with a picture of a convex mirror, for which the minimum principle is correct-not a concave mirror, for which it can be wrong. We shall find the same trick worked later by Fermat, who extended Hero's law to derive the refraction law. However, Fermat was called on the carpet for this by Abbé Mersenne, one of his contemporaries.

Hero of Alexandria-who, by the way, in his *Mechanics* described the so-called seven simple machines known to us-not only discussed spherical mirrors in his *Catoptrics* but, as an early practical experimentalist, also showed what can be done with cylindrical mirrors. If you go into an amusement

parlor and see yourself in mirrors of all kinds (I especially like those that elongate the height and diminish the girth), you know the proprietor has read Hero of Alexandria. If you see a play in a tanagra theater, where the actors appear on the stage, by the skillful use of mirrors, as beautifully dressed dwarfs, you know that Hero of Alexandria wrote the prescription. If you want to see around the corner or look at a person without letting him see you, Hero of Alexandria tells you how to do it with mirrors.

Ptolemy of Alexandria, who lived about A.D. 150, is best known for his *Almagest*, but he was also the first to make a serious attempt to study the law of refraction. He wrote a five-part volume entitled *Optics*, several copies of which have come down to us. This work, of which the first part has been lost (although we have a description of its contents), can be summarized as follows.

The first part deals with the relations between the eye and light, and the second with the conditions of visibility: size, form, color, and motion of bodies. Ptolemy explains why we usually see only one image of an object, although we have two eyes, and describes the conditions under which we see more than one image. The third part deals with the laws of reflection by plane and convex mirrors. Ptolemy gives probably the first attempt at an explanation of why the moon and the sun appear larger at the horizon than at the zenith a question that is still being discussed.

In the fifth part, Ptolemy tries to find a law for refraction. First he observes that for small incident angles the angles of incidence and of refraction are proportional to each other. He also remarks that, in transition from a thinner to a denser medium, the refracted ray comes earer to the perpendicular, the opposite being true for the reverse direction of travel. He uses an ingenious contraption to measure the incident and the refracted angles for the transition from air into water, air into glass, and water into glass. This instrument consists of a circular disk divided into 3600 with a colored marker in the center and two movable markers on the periphery. You put the instrument into water, line up the three markers, and read the two positions, one giving the incident angle and the other the refracted angle. Ptolemy gives refraction tables in 100 intervals. My friend, the late Hans Boegehold, studied these tables carefully and found that the values for 500 and 600 incident angles are surprisingly accurate, but that the other angles were obviously calculated by a second-degree interpolation formula. (The first differences are exactly $1^{\circ} 30' .14$) Ptolemy applies the same method for the transition of light from air into glass and from water into glass. He also discusses atmospheric refraction, the cause of which he assumes to be that the densities of air and of ether vary with height. He points out that only for a star in the zenith do the true and the apparent positions coincide.

1.1.2 The arabian period

After the fall of Greece, scientific progress ceased on the continent of Europe for nearly a century and a half, but Alexandria remained the center of scientific discovery. When the Arabs took over, they brought their scientific inheritance and knowledge of mathematics and the mathematical sciences into northern Africa and Spain. It is remarkable that, at a time when magic and mysticism and misinterpretation of Aristotelian doctrines prevented scientific progress over most of Europe, the

Arabs held high the torch of science in their countries.

The next big contribution to our science was by Ibn al-Haitham, or Alhazen of Cairo, whose date is frequently set around 1100, although some sources tell us he lived from 965 to 1038. Alhazen's "Optics" consists of seven parts. The first gives an anatomical description of the eye, which, he maintains, contains three liquids separated by four membranes. The names which he gives to them the watery medium, the crystalline medium, and the glassy medium are still used today. The phenomenon of binocular vision is explained by assuming that the pictures from the two eyes are transported to one visual nerve. Alhazen discusses the pupil of the eye and speaks of a pyramid of rays (rather than a cone) coming from the object to the eye.

The second part, like the corresponding book by Ptolemy, deals with the properties of bodies, and he distinguishes no fewer than twenty-two different properties.

The third part deals with optical illusions. He draws attention to the illusions arising from a false interpretation of a picture by the human mind and imagination. The fourth, fifth, and sixth parts deal with problems of reflection and mirrors. One of the problems studied in great detail has since been called after Alhazen: it consists in finding the point on a mirror where the incident and the reflected rays meet. Alhazen corrects errors of the Greeks about the position of the images in spherical, conical, and cylindrical mirrors. He also emphasizes the fact that the incident and reflected rays lie in the same plane.

Alhazen's seventh part deals with refraction. He proposes methods for measuring incident and refracted rays through different media, but he gives no tables. He knows that a glass sphere magnifies. He explains the apparently large size of the sun at the horizon by the assumption that we compare the sun with objects of known size lying along the line of sight. He investigates binocular vision and makes a good guess at the height of the atmosphere. He breaks with the idea of the Greeks that light rays originate at the eye and go to the objects, supporting his claim by the circumstance that sunlight can hurt the eye and that we can see afterimages.

Other manuscripts show that Alhazen used the pinhole to view solar eclipses and that he knew how the image becomes blurred and finally becomes circular as the pinhole increases in size. He did not write as though this were a discovery of his own, and Aristotle had noted that the spot formed by the rays from the uneclipsed sun is circular when the rays pass through a rectangular hole in wickerwork if the hole is small, but becomes rectangular when the hole is large. However, Alhazen's is the first known unequivocal description of the camera obscura. Alhazen's works were of enormous influence in the development of optics.

1.1.3 The middle ages

The knowledge of the Arabs was transmitted to Europe via Spain and Sicily and set in motion the intellectual revival that culminated in the Renaissance. The first of the European scientists to play

a role in optics was Vitello, a Polish mathematician who lived and studied in Italy about 1270. His book in ten parts is probably the most voluminous optical work ever written. It contains about five hundred closely printed folio pages. Unfortunately, he is more famous for his errors than for his contributions.

He reissued the tables of Ptolemy and added tables for the passage of light from water into air, from glass into air, and from glass into water. It is unfortunate that he never tells us how he arrived at his values, for they include impossible data, being in contradiction with the then unknown phenomenon of total reflection.

It was this error that later prevented Kepler from finding the correct form of the refraction law. Nevertheless, Vitello made two notable contributions. He explained the colors of the rainbow as originating from refraction and reflection by the water drops, and he made the proposal to collect the light of the sun into a point by using parabolic mirrors.

The Englishman Roger Bacon (1215-1294) discovered the spherical aberration of a mirror; he further stated that rays forming a cone around the axis in object space also form a cone around the axis in image space. From the magnifying property of a sphere, he speculated that it might be possible to invent instruments in which a man would look like a mountain and a small group like a big army. However, since he did not say how this could be achieved, it is doubtful whether he should be called (as he was by Sir John Herschel) the inventor of the telescope. His writings are so sketchy and indefinite that it is impossible to attribute specific inventions to him or to reach any conclusions about what he actually knew.

1.1.4 The renaissance

Coming to the Renaissance, we must mention Leonardo da Vinci (1452-1519). His notebooks show that he studied the eye anatomically and made a model of it, although he supposed the rays to cross twice in it to form an upright image. 18, 9 Immediately afterward he described the camera obscura but correctly showed the image inverted!20 He described, with sketches, a Rumford-type photometer and a machine for grinding concave mirrors. Unfortunately for his optical reputation and the development of optics, he never organized his notes into publishable form, so they were practically unknown until Venturi published them in 1797.

About half a century later came another Italian we should mention, Francesco Maurolico (1494-1577), who solved one of the problems that had bothered previous thinkers: that the image of the sun is round even if formed by a square pinhole. He improved our knowledge of vision by suggesting that the crystalline lens of the eye works like a double-convex lens. This enabled him to attribute myopia and hyperopia to an eye whose lens has a larger or smaller curvature than is proper for the length of the eyeball. He emphasized for the first time that the rays from an object point envelop a surface that is now called the caustic surface and that this surface is the place of the highest light concentration. And he explained the colors of the rainbow by multiple reflection in the

water drops. All this information was concisely organized by 1567 into a work entitled *Photismi de lumine*, although it was not published for some thirty years (1611) after the author's death, and then only as a consequence of the interest in optics engendered by Galileo's telescopic discoveries.

In Italy, under the influence of the philosophy of the time, scientific progress and magic were mixed in a way that led to interesting reading. The development of mathematical methods, however, served to gradually separate speculation and imagination from the solid basis of observation.

One of the most colorful books of the period was the *Magia naturalis* of Giovanni Battista della Porta (ca. 1538 or 1543 -1615). In his youth, della Porta had written a four-part volume entitled *De miraculis rerum naturalium* (1558) and, subsequently, many books and plays, including *De refractione* (Fig. 6) in nine parts in 1593. The *Magia naturalis*, which was an expansion of the earlier *De iraculis* into twenty volumes, appeared in 1589. It became so famous that it was soon translated from Latin into the vernacular languages Italian, French, Spanish, and English. It is a strange medley. Only the seventeenth part concerns optics, but some of the others are also worth noting for our amusement if nothing else. I mention the titles of a few of the twenty parts, to give you an idea of their contents. The first is entitled, "The Causes of the Miraculous. It deals with astrology, sympathy and antipathy, and the influence of place and time on events. The second deals with the origin of animals. He quotes many authorities who agree that snakes are made from the marrow of men and the hair of women. The third deals with garden plants; the fourth, with how to become wealthy by good housekeeping; the fifth, with how to make precious stones artificially; and the ninth, with make-up and other ways of improving the beauty of women (as if this were necessary!).

Finally, the seventeenth part deals with focal mirrors and other mirrors and explains what one can do with them. This book exhibits progress in the understanding of optical phenomena. In it are discussed the same tricks that Hero had dealt with 1400 years earlier, namely, how to use mirrors to obtain desired effects. However, we find here the first exact theory of multiple mirrors, the first complete description of the camera obscura, with pinhole or lens, and a discussion of combinations of positive and negative lenses designed to improve vision. Porta compares the eye and its pupil to the camera obscura and its stop. (Leonardo had anticipated him here, but less definitely and accurately.) At the end of this century also falls the invention of the telescope and the microscope.

Eyeglasses had apparently been invented toward the end of the thirteenth century, but they seem not to have attracted the attention of writers on optics until the practical Leonardo discussed methods of making them. Some lens grinder-more likely, several independently must have discovered the principle of the Galilean telescope by accident, and certain passages in Leonardo's notebooks a hundred years before Galileo can be interpreted as instructions for making such a telescope. Be that as it may, a Dutchman, Zacharias Jansen (1588-1632), was said by his son to have made a telescope in 1604 according to the model of an Italian dated 1590, and soon telescopes became well known in Holland. Galileo Galilei (1564-1642) heard of them and, apparently independently, made an instrument that he exhibited in 1609, and in his lifetime made over a hundred. The Dutch instruments were so imperfect that a magnifying power of more than 3X was impractical. Galileo

made instruments with a useful power of approximately 30 X, an order of magnitude greater. He was by no means the inventor of the telescope or even the first scientist to describe it; Vavilov reminds us that Leonardo may have anticipated him and that Porta described, three months after Galileo's demonstration, an instrument he had made independently. But Galileo is outstanding on two counts: he revolutionized man's ideas of the universe by his applications of the telescope and, by improving it by an order of magnitude, Ronchi points out that Galileo is entitled to be considered the father of the art of fine optics.

Moreover, from telescope to microscope is a short optical step, and by 1624 Galileo was using lenses of short focal length and thus making precursors of the modern compound microscope. 9 A few decades thereafter, the Dutchman Antonj van Leeuwenhoek (1632-1723) improved the quality of simple lenses used for magnifiers or simple microscopes, and in a letter of 1674 he began the descriptions of the discoveries for which he became justly famous.²⁴ Meanwhile, Robert Hooke (1635-1703) in England also improved the quality of lenses in order to make the compound microscope practical, and in 1665 the Royal Society published his *Micrographia*.

Aside from publications, which were in an embryonic state, the seventeenth century brought a lively exchange of ideas and thoughts between scientists, mostly friendly but sometimes polemical and frequently expressed in code to conceal them from the uninitiated. It is therefore necessary to study not only the books of these scientists but also their correspondence (cf. Leeuwenhoek's letters quoted by Dobell).

The same period witnessed the development of scientific knowledge to the system that is still important. Indeed, scientific development in the seventeenth century was greater than in any century before or since. Scientific giants like Kepler, Descartes, Pascal, Fermat, Newton, Leibniz, and Huygens appeared and left their marks in all fields of science, including optics. Let us sketch some of their contributions to geometrical optics.

Johannes Kepler (1571-1630) wrote two books on optics: *Supplements to Vitello* in 1604 and *Dioptrice* in 1611. The object of the first is to find the law of refraction. Kepler nearly succeeds, though he does not look for a relation between the incident angle and the refracting angle but between the incident angle and the deviation. He asks himself what form a surface should have to refract a parallel beam of light exactly into a point image. He studies the hyperboloid, which, as we know, fulfills the condition; but he rejects it because it disagrees with the values that Vitello gave for the relation between the angles of the incident ray and the refracted ray. If only the Kepler of 1611 and the *Dioptrice*, in which he discovered the law of total reflection, had reread his book of 1604, he would have seen that the data of Vitello had been erroneous and thus he might have found the correct form of the law of refraction.

Kepler investigated the function of the eye more profoundly than his predecessors. He discovered that the eye is not at rest but moves around a point inside it, and he investigated the importance of this motion on our vision. He drew attention to the fact that the image on the retina is inverted.

All this is in his Supplements to Vitello.

His Dioptrice, although a relatively small book, was of importance to the development of optics. In it, Kepler assumes that for small angles the angle of refraction is proportional to the angle of incidence. From this assumption, he develops the laws of firstorder optics for thin lenses and systems of thin lenses with finite separations. He describes, among other things, the Keplerian telescope, in which two positive lenses are placed so that the back focus of the first is at the front focus of the second, and compares it with the form named after Galileo, consisting of a positive and a negative lens placed so that the focus of the former is at the virtual focus of the latter. He gives the focal length correctly for all forms of lenses. His book contains many practical proposals for lens design and is highly recommended to those who are interested in these problems.

Rene Descartes (1569-1650) must be considered as one of the discoverers of the correct law of refraction, for he published the sine law in his Dioptrique in 1637. After his death, he was violently attacked by Isaac Voss, who accused him of stealing the law from Willebrord Snel* (1591-1626). Descartes had indeed visited Leyden in 1630, where Snell, a professor of physics, had written a book on optics containing the refraction law. Although the book had never appeared, Voss, in his posthumous attack, claimed that Descartes had seen it on his visit after Snell's death, and Voss quoted Snell's proof.

The question of priority has been much discussed in the literature. A recent study has been published by Boegehold, who refers to a paper by P. Kramer. Boegehold found that Descartes, in some letters written in 1627, had suggested making plano-convex lenses with hyperbolic surfaces. Since such a lens would give a sharp point image for an infinite object if and only if one assumes the sine law, it is fair to conclude that Descartes knew of the law three years before his visit to Leyden in 1630. It is strange that Voss should have years after waited for twenty-five years-twelve Descartes's death-to make his accusation. Credit for discovering the law of refraction should therefore be divided between Descartes and Snel, with a fair share going to Johannes Kepler, who failed to find the sine law only because he trusted the tables of Vitello.

Descartes's Dioptrique is beautifully written. It tries to explain the law of refraction by means of the theory of elasticity. A light ray is compared to the path of a ball, first being bounced off the ground, then being slowed down as it goes into a denser medium. Descartes was the first to suggest determining the velocity of light from the position of the moon at an eclipse.

He also made experiments with human and animal eyes. If the eye is removed and put in the opening of a window in a darkened room, one can see the images of distant objects on the retina, and by pressing the eye a little, one can see the image of near-by objects, showing that the eye can accommodate.

In the eighth chapter of his book, Descartes investigates surfaces that image a point sharply-the so-called Cartesian surfaces, of which the conic sections constitute a special case. He proposes machines for grinding lenses and demonstrates the colors of the rainbow with small glass spheres filled with water.

But much of Descartes's reasoning in connection with the refraction law was faulty and was rejected by the great mathematician Pierre de Fermat (1601-1665), who pointed out that water would have to be less dense than air to fit Descartes's explanation. Fermat returned to the formulation of the refraction law given by Hero, deriving it from the assumption that light has different velocities in different media and takes the quickest path between any two points. Like Hero, he illustrates this idea with a drawing of a convex surface, for which the minimum principle is correct, rather than a concave surface, for which it can be wrong. But he was called on the carpet by Abbé Mersenne, as was mentioned earlier, one of the great scientists and letter writers of his day. Fermat's answer is a classic example of double talk: he says that the path would be a minimum if the surface were a plane, and the incoming ray sees so little of the surface that it cannot distinguish whether the surface is plane or curved!

1.1.5 Newton and Huygens

Sir Isaac Newton (1642-1727), in his *Lectiones opticae* of 1669-1671 (published posthumously), showed how to calculate the spherical aberration of a plano-convex lens for an infinite object point. This book contains the formula for locating the image, still known by his name, in terms of the distances of the object and the image from the focal points. It also contains a thorough investigation of reflection in prisms, including the position of minimum deviation. It even goes into some aberrations, in particular, image errors as functions of color, and it compares the size of the color aberration with the spherical aberration of a given lens and prism.

Although Newton's best known work in optics is his *Opticks*, I should like to draw your attention especially to his beautiful paper on color, in which he describes the separation of white light into the colors of the spectrum by means of prisms. This charming paper should be read by every student of optics, especially since the writer not only gives his final conclusions but describes many false attempts at interpreting his experiments and the refutation of these attempts.

Newton was misled into believing that all materials have the same relative dispersion, since he found this to be true for such different media as glass and water. The conclusion that he drew-that telescopes should not contain refracting surfaces-was erroneous. But it led him to study mirror telescopes and, with his teacher Gregory, he contributed much to the development of these instruments.

Christian Huygens (1629-1695) wrote two books on optics: *Dioptrical* in 1653, published posthumously, and *Traite de la lumiere*, written in 1678 but not published for twelve years. The

first book shows enormous advances with respect to the understanding of optical image formation. The second proposes a new and revolutionary theory of light, deriving refraction and reflection laws from a completely new principle.

This principle, set forth earlier in the *Dioptrica*, is a starting point from which all the laws of Gaussian optics can be derived for lens systems with finite thicknesses. It is expressed as follows: if we interchange object and eye in an optical system, the object will have the same apparent size and orientation as before. The law is known to us in the form that the product of lateral magnification and angular magnification is constant in an optical system, and equal to the ratio of the refractive indices.

Huygens found, before Newton, the law that, for a single surface or for a single lens, the image distance is proportional to the object distance when the distances are measured from the respective principal foci. He knew of the aplanatic surface. He was acquainted with the important property of the optical center of a lens (the point which divides the distance between the vertices in the ratio of the radii), namely, that a ray through it emerges parallel to its original direction. He had an approximate formula for the spherical aberration of a thick lens that enables the form for minimum spherical aberration to be determined.

In his *Traite de la lumiere*, which should be required reading for every physicist and is now readily available, Huygens derives optical laws from the fact that the light emerging from an object point P forms a wave surface S at any given time t. The light rays are then the normals to this surface and are not straight unless the velocity is constant throughout the medium.

The surface can also be constructed in the following way. At an arbitrary time, light from the point source P forms a wave surface S. (assumed to be spherical in the figure). Considering each point of S, as a source sending out a new wave, we see that the envelope of all the waves at the time gives the original wave surface S.

With the help of this concept, Huygens derives the straight paths of light in a medium of constant velocity. He derives not only the reflection and refraction laws but also the law of refraction in the atmosphere and in doubly refracting crystals. He clearly defines the caustic as the evolute of the rays and investigates it after reflection and refraction, in addition to investigating the degenerate form of the wave surfaces in the points of the caustic. He proves Malus' law: that rays form a normal system for a single reflection and refraction.

It should be emphasized, however, that this formulation is mathematically identical with ray optics, as Hamilton showed in his paper of 1831. The assumption that light has a finite wavelength and that it can be described as a sine wave going along a ray is foreign to Huygens. However, once this theory was established, the methods of Huygens enabled Fresnel to derive all the laws of diffraction in a most elegant manner.

Chapter 2

Geometrical Optics

*The Euclidean Geometry exists thanks to light.
Euclid wanted to understand the light rays, so he
discovered the plane geometry.*

“Music is the arithmetic of sounds as optics is the geometry of light.”
– Claude Debussy

2.1 The postulates of geometrical optics and the dark Room

We already know the history of the geometrical optics, well before starting the study of this theory we must take back the contributions of Al Azhen and talk about the **postulates of geometrical physics** which are

1. The speed of light is finite.
2. The light propagates in a straight line.
3. The light rays obey the Euclid's postulates of plane geometry.
 - (a) A straight line segment can be drawn joining any two points.
 - (b) A finite straight line (line segment) can be extended continuously in a straight line indefinitely.
 - (c) A circle can be drawn with any point as its center and any distance as its radius.
 - (d) All right angles are equal to one another, regardless of their orientation.
 - (e) If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two lines, if extended indefinitely, will eventually meet on that same side.
4. In the matter the light's speed is less than in vacuum.

Now, before starting to use these postulates we must understand the physical meaning of them. The first postulate, **The speed of light is finite** is fundamental, it is incredibly important.

How could Hero of Alexandria understand that the speed of light must be finite? To be honest, I don't really know his motivation but, we could get an idea about why this should be like that from simple phenomenological reasoning without the use a mathematics or advanced physics. *Lets imagine a world where the speed of light is infinite*

How are going to be the nights in such world? It should be a complete chaos. Every star in the sky is a source of light, a source of infinite rays which each one travels at an infinite speed. There wont be any night, all the day is going to be full of light and even every event in the universe is going to be seen simultaneously, the big bang, the birth and death of a star. *Each past and present event is going to be seen at the same time.*

Other effect that only can be explained in a universe with light with finite speed. Imagine you're standing in the rain, which is falling vertically. If you start running, the rain hits you in the face. This happens because your movement changes the apparent direction of the rain. An astronomer, James Bradley, observed that the stars didn't appear to be exactly in their calculated

positions. To see a star directly overhead, he had to tilt his telescope slightly in the direction of Earth's movement around the Sun, the light from the star "falls" toward Earth at a finite speed. Earth is moving. For the light beam to travel from the telescope lens to the eyepiece (and for us to see it), Earth has moved a little. Therefore, you have to point the telescope at a slight angle toward where Earth will be when the light reaches the eyepiece, not where it is at the moment it enters the lens.

If the speed of light were infinite, this effect wouldn't exist. The rain (light) would fall so fast that your (Earth's) movement would be irrelevant. The fact that we have to "aim ahead" of the target is irrefutable proof that light travels at a finite speed.

There are many other consequences but let's continue with all the other postulates. Now let's talk about **The light propagates in a straight line**. This does have sense because it is what is seen in any normal day, all the source of light can be seen as sources of infinite many rays which propagate at a finite speed. With tools purely geometrical there is no way to give a proof of this, however, from a wave theory for light, Huygens was able to prove that the rays are, in fact, the normal vectors of the wavefront are the rays, and, from a quantum perspective, Richard Feynman proved that the rays are the superposition of the "waves" that represent the probability amplitudes.

Often we see that light is not a straight line, for example, an easy experiment consists on a translucent bowl full of salted water, we're going to see that the ray presents a deflection, we're going to see that this curve can be generated as a composition of infinitely many straight refractions.

Because we've postulated that the light travels in straight lines, so we have to give a theory for the propagation so **The light rays obey the Euclid's postulates of plane geometry**. A point is going to be a source, a place where two rays meet, the place where two media separate, etc.

The extension of a line segment to a continuously straight line is a way to make or represent ray tracing. The circle postulate talks about the sphere that propagates alongside all the rays (which behave as the radius). The right angles are just a geometrical formalism which does not have a dramatic optical interpretation.

The last postulate is really important, from a physical point of view it tells us that all the rays are going to be propagate on a completely plane space, so we could ask ourselves if we want to study the light's behavior under the gravitational effects of compact objects we must change the fifth Euclid's postulate, and the short answer is yes, we have to. Under gravitational effects the differential geometry is a better tool to study the light-type observers on the curved space-time.

So, we have many arguments that support the idea the light's speed must be finite, and if it is finite there should be a way to measure it, right? The answer is yes, and here we're going to use the postulates to follow the Fizeau's ideas on how to measure the light's speed.




Figure 2.1: Schematic representation of the Fizeau's experiment pretended to measure the light's speed.

In 1849, Hippolyte Fizeau performed a groundbreaking experiment that provided the first non-astronomical measurement of the speed of light. His ingenious setup transformed the problem of measuring an extremely short time interval into a problem of measuring a rotation speed, which could be done with much greater precision. Imagine a very fast runner who starts from a point A , runs to a point B and immediately returns to A . To measure the runner's speed, you could place a door at A that opens and closes at regular intervals. If you open the door just when the runner leaves, then close it and open it again periodically, you might catch a glimpse of him when he returns if the timing is right. But if the runner is extremely fast, the door might be closed when he gets back. This is exactly the idea Fizeau used with light, replacing the door with a rapidly rotating toothed wheel and the runner with a light pulse.

Fizeau set up his apparatus on two hills in the outskirts of Paris, with the light source and observer on one hill (Montmartre) and a mirror on the other (Mont Valérien), about 8.633 kilometers away. A beam of light from a lamp was directed toward a partially reflecting mirror (a beam splitter) that sent part of the light toward a toothed wheel. The wheel could be spun at a controlled speed and had a large number of teeth and gaps – typically 720 teeth, meaning 720 gaps as well. After passing through a gap, the light traveled to the distant mirror, was reflected, and returned along the same path. On its way back, it would again have to pass through a gap in the wheel to reach the observer's eye via the beam splitter.

If the wheel were stationary, the returning light would pass through the same gap and be seen. But as the wheel began to rotate, the situation changed. When the wheel turned slowly, the gap through which the light had originally passed would have moved only slightly during the light's round trip, so most of the returning light still made it through. As the speed increased, the gap shifted more, and the light began to be partially blocked by the edge of a tooth. At a certain critical speed, the gap moved exactly so that by the time the light returned, the adjacent tooth had taken its place, completely eclipsing the light. The observer would see the light disappear for the first time. By knowing the rotation speed at which this first extinction occurred, the number of teeth, and the distance to the mirror, Fizeau could deduce the time it took for light to travel to the mirror and back.

Let us analyze the situation mathematically. In the figure 2.1 let d be the distance from the wheel to the distant mirror. The total path for the light is $2d$. The wheel has N teeth and consequently N gaps, so the angular width of one tooth (or one gap) is π/N radians (since a full circle has 2π radians and there are $2N$ segments – teeth and gaps – each of equal angular size). For the first extinction, the time required for the light to make the round trip must equal the time needed for the wheel to rotate by the angle corresponding to one tooth (i.e., to replace a gap with a tooth). If the wheel rotates with angular velocity ω (radians per second), the time to turn through an angle π/N is

$$t_{\text{rot}} = \frac{\pi/N}{\omega}. \quad (2.1)$$

The light travel time is

$$t_{\text{light}} = \frac{2d}{c}, \quad (2.2)$$

and equation these gives us

$$\frac{2d}{c} = \frac{\pi}{N\omega}. \quad (2.3)$$

But $\omega = 2\pi f$ where f is the rotation frequency in revolutions per second. Substituting,

$$\frac{2d}{c} = \frac{\pi}{N \cdot 2\pi f} = \frac{1}{2Nf}. \quad (2.4)$$

Solving for c yields

$$c = 4dNf. \quad (2.5)$$

Fizeau used $d = 8.633 \times 10^3$ m, $N = 720$, and observed the first extinction at $f \approx 12.6$ revolutions per second. Plugging in these numbers,

$$c = 4 \times 8.633 \times 10^3 \times 720 \times 12.6 \approx 3.13 \times 10^8 \text{ m/s}, \quad (2.6)$$

a value about 5% higher than the modern accepted value of 2.99792458×10^8 m/s, but remarkably accurate for the time. The discrepancy arose mainly from uncertainties in the distance measurement and the precise determination of the extinction condition. Fizeau's experiment was a triumph of

experimental physics, demonstrating that light travels at a finite speed and providing a method that would be refined by later researchers, most notably Léon Foucault, who replaced the toothed wheel with a rotating mirror to achieve even greater precision. The conceptual beauty of Fizeau's approach lies in its translation of an infinitesimal time interval (on the order of $1/20000$ of a second) into a measurable mechanical rotation, a perfect illustration of the deep connection between optics and mechanics.

Now we know the speed of light is finite, so let's use these postulates to understand one of the biggest contributions of Al Azhen, the dark room, see figure 2.2.




Figure 2.2: Graphical representation of a generic dark room which is a box of an arbitrary material which has a side opened and while it satisfies the Rayleigh's inequality it will form an inverted image.

Before developing mathematically the dark room we're going to set an important convention

- The light propagates from left to right along the z direction.
- A distance measured in the direction of light is positive and a distance measured against the direction of light is negative.

Once defined the convention let's do it. A source in the point A_o is at a distance d_o respect to the opening O of the dark room, (d_o is in fact negative because it is measured against the light direction from the point O), an image is formed at a distance d_i from O thus we must find out a relation between these parameters and the diameter D of the DR. First we're going to calculate the distances $\overline{A_o O A_i}$, $\overline{A_o R A_i}$ where R is the radius of the opening, therefore

$$\overline{A_o O A_i} = d_i - d_o \quad d_i = \overline{O A_i} \quad d_o = \overline{O A_o}, \quad (2.7)$$

$$\overline{A_oRA_i} = [d_i^2 + x^2 + y^2]^{1/2} - [d_o^2 + x^2 + y^2]^{1/2}, \quad (2.8)$$

having on count our convention implies that $\sqrt{d_o^2} = -d_o$ so

$$\overline{A_oRA_i} = d_i \left[1 + \frac{r^2}{d_i^2} \right]^{1/2} - d_o \left[1 + \frac{r^2}{d_o^2} \right]^{1/2}. \quad (2.9)$$

Historically it was observed in the dark room that the image that was formed depended on the diameter of the opening and the distance of the object to view, it was observed that the image was formed for far objects, but how far is far? It is relative to the radius of the opening, then the ratios must satisfy that

$$\frac{r^2}{d_o^2} \ll 1, \quad \frac{r^2}{d_i^2} \ll 1, \quad (2.10)$$

using this and using the Taylor's theorem for a polynomial expansion of the function $\sqrt{1+x}$ then

$$\overline{A_oRA_i} \approx d_i \left[1 + \frac{1}{2} \frac{r^2}{d_i^2} + \mathcal{O}(\delta^2) \right] - d_o \left[1 + \frac{1}{2} \frac{r^2}{d_o^2} + \mathcal{O}(\delta^2) \right], \quad (2.11)$$

$$\overline{A_oRA_i} \approx (d_i - d_o) + \frac{r^2}{2} \left(\frac{1}{d_i} - \frac{1}{d_o} \right) = \overline{A_oOA_i} + \frac{r^2}{2} \left(\frac{1}{d_i} - \frac{1}{d_o} \right) \quad (2.12)$$

with this we can express this in terms the difference of the light's path and the diameter of the dark room

$$\Delta L = \frac{r^2}{2} \left(\frac{1}{d_i} - \frac{1}{d_o} \right) = \frac{D^2}{8} \left(\frac{1}{d_i} - \frac{1}{d_o} \right). \quad (2.13)$$

In the equation 2.13 lies the hearth of the dark room, it was found by Rayleigh a certain relation between the light's wavelength and the light's path difference between the upper and lower part of the opening of the dark room, this relation (which will be discussed physically in a moment) is:

$$\Delta L < \frac{\lambda}{4}. \quad (2.14)$$

Using this relation we will find the maximum diamater for the dark room

$$\frac{D^2}{8} \left(\frac{1}{d_i} - \frac{1}{d_o} \right) < \frac{\lambda}{4}, \quad (2.15)$$

$$D_{max} = \left(\frac{2\lambda d_i d_o}{d_o - d_i} \right)^{1/2}. \quad (2.16)$$

In the real life there is no monochromatic light, it is forbidden by the laws of quantum mechanics, we can have quasi-monochromatic light but it is only in advanced laboratories so for a polychromatic source in the equation 2.16 we should do the calculations with the maximum wavelength that conforms the beam the origin of this expression have many origins which converges to the same phenomena. If we want to form an image the electromagnetic field coming from the object must arrive at the same time at the image point, if there is some kind of phase difference there will be aberrations as astigmatism or chromatic aberration (even if it is not a lens we can see it from the equation 2.16) from a quantum point of view this is related to an interpretation of the *size* of a photon (Santos et al., 2018).

Now we're going to analyze how does the light interact (from a simple and geometrical point of view) with transparent media. Suppose a source radiates from a point A and the ray light arrives at B , see figure 2.3.




Figure 2.3: A schematic representation of a ray beam from A to B which goes through n different transparent media and in each one it had a different speed and traveled different length.

There are n different transparent media each one has a different thickness and the light in each one is different, therefore the time that takes the light to go from A to B is

$$t_{AB} = \sum_{i=1}^n \frac{L_i}{v_i} = \sum_{i=1}^n \frac{L_i C}{v_i} \frac{1}{C} [s]. \quad (2.17)$$

Then we define the refraction index in the i th medium as the relation between the speed of light in vacuum and the speed of light in the i th medium. *It is a scalar in linear and homogeneous and isotropic media but mostly it is a tensor.*

$$t_{AB} = \frac{1}{C} \sum_{i=1}^n n_i L_i = \frac{L_{AB}}{C} [s], \quad (2.18)$$

Definition 1 (*Optical path*): The optical path L_{AB} is the distance that must travel in vacuum the light to have the same time to go from a point A to a point B across n different media each one characterized by a different (or equal) refraction index.

The optical path has a length greater than the geometrical path of the light, that is why it is called optical path. This can be easily generalized to continuous media using a Lebesgue's integral (or Riemann integral if it is well behaved)

$$L_{AB} = \int_A^B n(S) ds [m]. \quad (2.19)$$

Before going further we must have to set some definitions.

Definition 2 (*Phase*): The phase is the relative angular displacement the ray has.

Definition 3 (*Wavefront*): The geometrical space where the phase of the light is constant is called the wavefront of the light.

2.2 The Snell-Descartes law and the Fermat's principle

As we had discussed in the historical chapter it was Heron of Alexandria who realizes that in light reflection it travels the least distance and Al Azhen proposes a maximum time, it was Fermat who formalizes it as a principle the **Fermat's principle**

Principle 1 (*Fermat's principle*): The light travels a trajectory such that the optical path is stationary

$$\delta L = 0. \quad (2.20)$$

The Fermat's principle can be “deduced” from the Hamilton principle or even more from a variational principle, lets give a look to it, lets look for an stationary curve for the time the light travels

$$t = \int_{t_0}^{t_f} dt = \int_A^B \frac{1}{C} dL = \int_{\tau_0}^{\tau_f} L d\tau, \quad (2.21)$$

from here we see that, in a mathematical way, if we want to find the relation between the trajectory of the light and the time it takes to go from A to B we will have to solve the Euler-Lagrange equation. Now lets use the Fermat's principle to see how the light interacts with a diopter (a refracting surface) in a vector way, known as the *vectorial Snell-Descartes law*.

Definition 4 (Diopter): A Diopter is a surface $\Sigma \subset \mathbb{R}^2$ such that it separates two media with different refraction index.




Figure 2.4: two media separated by a diopter which an observer O measures the path the light will follow from A to B and proposes a displacement of the source A to A' to see how must be the light path.

In the figure 2.4 we see two media separated by a transparent surface (diopter) with refractive index n_o, n_1 , respectively, a source A emits a beam which reaches the diopter in I and it is refracted to B , from A there is an unitary vector $\hat{u}_o$ in the direction $\vec{AI}$ and the observer O analyzes a variation of A as A' and the incident point on the diopter is I' , then lets analyze it vectorially

$$L = \left\| \vec{I} - \vec{A} \right\|, \quad (2.22)$$

$$\Delta \vec{I} = \vec{I}' - \vec{I}, \quad (2.23)$$

$$\Delta \vec{A} = \vec{A}' - \vec{A}, \quad (2.24)$$

with this we can calculate the path difference $L - L'$ vectorially as a projection of the unitary vector in the direction of L given by $\hat{u}_o$ along $\Delta \vec{I} - \Delta \vec{A}$

$$\Delta L_A = \frac{(\vec{I} - \vec{A}) \cdot (\Delta \vec{I} - \Delta \vec{A})}{\left\| \vec{I} - \vec{A} \right\|} = \hat{u}_o \cdot (\Delta \vec{I} - \Delta \vec{A}), \quad (2.25)$$

now for the optical path

$$\mathcal{L}_A = n_o \Delta L = n_o \hat{u}_o \cdot (\Delta \vec{I} - \Delta \vec{A}), \quad (2.26)$$

doing the same for the right side of the diopter but for the point B we then have got the total optical path difference as

$$\Delta \mathcal{L} = n_o \hat{u}_o \cdot (\Delta \vec{I} - \Delta \vec{A}) + n_1 \hat{u}_1 \cdot (\Delta \vec{B} - \Delta \vec{I}) \quad (2.27)$$

because the points A and B are fixed then $\Delta \vec{A} = 0$ and $\Delta \vec{B} = 0$ thus

$$\Delta \mathcal{L} = (n_o \hat{u}_o - n_1 \hat{u}_1) \cdot \Delta \vec{I}. \quad (2.28)$$

If the Fermat's principle is satisfied the equation 2.28 implies straightfully that $\Delta \vec{I}$ is a tangent vector to the diopter this implies then

$$n_o \hat{u}_o - n_1 \hat{u}_1 = k \hat{N}. \quad (2.29)$$

The equation 2.29 is known as the vectorial Snell-Descartes law, from this equation we can arrive to the law of reflection and refraction to do it we only have to do the cross product of the equation 2.29

$$\hat{N} \times (n_o \hat{u}_o - n_1 \hat{u}_1) = k \hat{N} \times \hat{N} = 0, \quad (2.30)$$

$$n_o \sin(\theta_0) = n_1 \sin(\theta_1). \quad (2.31)$$

2.2.1 The critical angle, and the refraction by a prism

We had deduced the vectorial Snell-Descartes law, which is a strong law and we are going to apply it to some physical situations, but before doing that we are going to talk briefly about where is the physical regime where this law can work on.

We are working under the postulates of the geometrical optics, but lets analyze a bit the second one, *the light propagates in a straight line* we haven't say anything about the light's shape, if we say it propagates in a straight line we can image light rays as circular cylinders but what about the radius of this ray? Well, in fact there are more shapes we can imagine of the wavefront and the beam's profile, but we are going to talk about it in the chapter of electromagnetic optics, for now keep with the cylinder image.

If we make this ray to inside over a perfectly polished mirror as shown in 2.5 then we can see an prove easily experimentally the Snell-Descartes law will be fulfilled but what happen if we have not got a polished surface but an extremely irregular surface? we will have reflections in all the points over the surface as shown in the figure 2.6.




Figure 2.5: Representation of a cylindrical ray light being reflected by a perfectly polished mirror.

The phenomena shown in the figure 2.6 is, often, called *diffuse refraction* but it is a kind of macroscopic effect of the Snell-Descartes law but there are other behaviors of the refraction index which give a particular behavior to the SD law, which are related to the electromagnetic properties of the matter, in fact, the refraction index n is a complex quantity, which its real part is related to the refractive properties of the material and the imaginary part is related to absorption and losses in




Figure 2.6: Representation of a cylindrical bunch of rays being reflected by an inhomogeneous surface.

the energy but we will work on this later.

There are some other special materials where the equation 2.29 has an amazing behavior. The metamaterials which have a negative refractive index, but how can we interpret this? It is easy to see in the equation 2.29 that a variable change of $n \rightarrow -n$ implies a physical effect of $\theta \rightarrow -\theta$ this means, the rays will be reflected or refracted to the other side!

Well then, let's come back to our well-defined behavior, where the SD law 2.29 will be fulfilled. Can we find a condition between two materials such that there is no refraction? It is worth asking about it because in nature reflection and refraction happen at the same time both simultaneously because a translucent surface is not perfectly transparent and a refractive surface is not perfectly reflective. This is related to the Fresnel's coefficients but this will be hard work of the chapter of electromagnetic optics, for now have on mind reflection and refraction happen at the same time. To try to kill the refraction we have to study two cases between two media.

1. $n_1 > n_2$
2. $n_1 < n_2$

For the first case then we have from 2.29

$$n_1 \sin \theta_1 = n_2 \sin \theta_2, \quad (2.32)$$

$$\frac{n_1}{n_2} \sin \theta_1 = \sin \theta_2. \quad (2.33)$$

If $\theta_1 = 0$ then $\theta_2 = 0$ thus if the incident ray is perfectly parallel to the normal vector to surface there wont be any refracted ray, only a completely reflected in the same plane as the incident but with opposite direction, this is said to get a phase shift of π . If the ray's incidence angle is perpendicular to the normal vector to surface $\theta_1 = \pi/2$ then we will have

$$\frac{n_1}{n_2} = \sin\theta_2, \quad (2.34)$$

but $n_1 > n_2$ then $n_1/n_2 > 1$ which has no sense with the right hand side of the equation 2.34 because $0 \leq \sin\theta_2 \leq 1$ therefore for $\theta_1 = \pi/2$ there is no refracted ray this is illustrated in the figure 2.7.




Figure 2.7: Representation of all the incident and refracted rays from a media with refraction index n_1 to another with refraction index n_2 such that $n_1 > n_2$. The angle θ_2 here shown wont never reach a value equal to $\sin^{-1}\left(\frac{n_1}{n_2}\right)$.

Now, we have to consider and study the case where $n_1 < n_2$ the obvious case is when $\theta_1 = 0$ therefore $\theta_2 = 0$ and there wont be any refracted ray, but the interesting thing comes when $\theta_1 = \pi/2$ then again we have got the same equation 2.34 but here is the main difference, now we've got $n_1 < n_2$ or $n_1/n_2 < 1$ therefore there exists this angle and it is called the *total reflection angle* or *critical angle* θ_c this is illustrated in the figure 2.8.

2.3 The Descartes's ovoid and diopters

From here on we are going to start to study the theory of image formation where the concepts of **object** and **image** are really important because there are real and virtual objects and image, this depends on the relation between these points and the dioptr, in fact, something is called virtual if we put a photo detector in that point and we wont measure any field if an object is to the left of a dioptr is called real and if it is to the right is called virtual, in the same order of ideas if the image




Figure 2.8: Representation of all the incident and refracted rays from a media with refraction index n_1 to another with refraction index n_2 such that $n_1 < n_2$. The angle θ_c here separates the rays that will be refracted, (angles less than θ_c), and the rays that will be totally reflected internally.

is to the left of the diopter it is virtual and if it is to the right is real.

Before starting we have to give some fundamental definitions

Definition 5 (*Optical system*) an optical system is a set of translucent media (diopters) and reflective media (catopters) and we say that the **system is centered** if all the diopters are revolution surfaces and shared the same symmetry axis.

The systems conformed only by refractive surfaces all called **dioptric systems** and the system conformed only by reflective surfaces are called **catoptric systems** and the systems formed by these two kind of surfaces are called **catadioptrics systems**.

Definition 6 (*Real and virtual points*) Given a optical system $\mathcal{S}$ we say that an object point A_o is real if it is to the left and outside the optical system and we say that the object point is virtual if it is to the right and inside the optical system.

In the same way, we say that an image point A_i is real if it is to the right and outside the optical system, and we say that the image point is virtual if it is inside the optical system.

To understand better these two definitions let's follow the figures 2.9 there we can see two examples of optical systems. In the first one we can see an object point located to the left and outside of the system $\mathcal{S}$ so it is a real object point, and the point A_i is located to the right and outside the optical system, therefore it is a real image point. In the second optical system we can see that the same




Figure 2.9: Representation of two optical systems where appear real object points, real and virtual image points.




Figure 2.10: Optical system $\mathcal{S}$ formed by two dioptrics $\mathcal{S}_1, \mathcal{S}_2$ where the point A_o is a virtual object and A_i is a virtual image point.

real object point and that the point A_i is located inside the optical system, therefore it is a virtual image point for A_o .

In the figure 2.10 we can see two dioptrics $\mathcal{S}_1, \mathcal{S}_2$ which forms the optical system $\mathcal{S}$ where the point A_o is located inside the optical system, then it behaves as a virtual object point and A_i is to the right of A_o and also inside the optical system, therefore it is a virtual image point.

In the optical system 2.11 we can see three optical systems $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ and a configuration of many points. The first one, A_o is completely to the left and outside any optical system, so it is a real object point, and the point A' is inside the optical system $\mathcal{S}_1$ so it is the **the virtual image of the real object** A_o but, as we can see, A' is to the left and outside the optical system $\mathcal{S}_2$ so it is a real object point for the rest of the total optical system. The point A'' is inside the optical system $\mathcal{S}_3$ so it is




Figure 2.11: Optical system $\mathcal{S}$ formed by three optical systems $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ where the point A_o is a real object point, A' is a virtual image point for $\mathcal{S}_1$, but a real object point for $\mathcal{S}_2, \mathcal{S}_3$, A'' is a virtual image for the real object A' and A_i is the real image of the virtual object A'' .

the virtual image of the real object A' , and in the very last there is the point A_i which is completely to the right and outside any optical system, so it is the real image point of the virtual object point A'' .

Definition 7 (*Stigmatism*) From a **geometrical point of view** we say that an optical system $\mathcal{S}$ is **stigmatic** if for any given point source, all the rays that pass through $\mathcal{S}$ converge to a point source. In other words, a optical system $\mathcal{S}$ is stigmatic if for any object point there is one unique image point.

We can also, define the stigmatism from a wavefront perspective. An rigorously stigmatic system is one that transforms an spherical wavefront $\mathcal{A}$ to a new spherical wavefront $\mathcal{A}'$ whose curvature centers, A_o and A_i are the rigorously stigmatic points (see figure 2.12). Thus, the spherical wavefront $\mathcal{A}'$ is the image of the spherical wavefront $\mathcal{A}$.

Understanding that an image-forming optical system is one that maps each point source of an object's electromagnetic field to the point sources of the image's electromagnetic field, corresponding to a homothety between these fields, the interest arises in studying what these optical systems are and what their characteristics are. In general, these systems can be composed of multiple elements, but initially we will study the simplest configuration for two reasons:

1. because it most easily allows us to understand the principles of these systems.
2. because later, more complex systems can be understood as compositions of these.

These elementary systems are made up of a single optical surface, either refractive or reflective, which is capable of perfectly focusing all the rays emitted by a point source into a perfect point image. In geometric optics, this means that all the rays travel the same optical path length. Initially, this property associated with a single pair of points (stigmatism) will be studied, and then the conditions under which this property becomes invariant to the translation of these points, which is known as




Figure 2.12: The stigmatic system transform the incident wavefront $\mathcal{A}$ to another new spherical wavefront $\mathcal{A}'$.

aplanetism, will be studied.

Now we are going to develop the *rigorous stigmatism* these are system capable of making perfect images, this means from a point source there is a perfect point image. Let a diopter which separates two media with refraction index n_o and n_i to the left of the diopter there is an object A_o at a point $[0, 0, z_o]$ the field from A_o reaches a point I , measured from the diopter's vertex v , from the point I the beam is refracted and forms an image A_i at a point $[0, 0, z_i]$ then we are going to find the relations between all these parameters to make all the rays from A_o arrive at A_i making a perfect image. From the Fermat's principle we have got

$$n_o l_o + n_i l_i = cte \quad (2.35)$$

the, as we can see in the figure 2.13 we then have

$$-n_o[x^2 + y^2 + (z - z_o)^2]^{1/2} + n_i[x^2 + y^2 + (z - z_i)^2]^{1/2} = cte \quad (2.36)$$

and, remembering our sign convention $\sqrt{z_o^2} = -z_o$, $z_i^2 = z_i$ and to find the constant we can see that if we want to make all the rays from A_o arrive to A_i then the rays that travel parallel to z satisfy also the Fermat's principle, thus

$$n_i z_i - n_o z_o = cte, \quad (2.37)$$




Figure 2.13: Scheme representation of the development of a rigorously stigmatic system, from a point source A_o at a position $[0, 0, z_o]$ the diopter with a vertex v forms a perfect image A_i at $[0, 0, z_i]$.

$$n_i[x^2 + y^2 + (z - z_i)^2]^{1/2} - n_o[x^2 + y^2 + (z - z_o)^2]^{1/2} = n_i z_i - n_o z_o, \quad (2.38)$$

now, we are going to do the following change of variables given by

$$\rho^2 = x^2 + y^2 + z^2, \quad (2.39)$$

which represents the square of the vector from the vertex to the interface shown in 2.13, so, having this on count, the equation becomes

$$n_i [\rho^2 + z_i^2 - 2zz_i]^{1/2} - n_o [\rho^2 + z_o^2 - 2zz_o]^{1/2} = n_i z_i - n_o z_o. \quad (2.40)$$

Now, the main goal is to transform the equation 2.40 into a more explicit and useful algebraic equation, so, the first thing we're going to do is to square both side of the equation 2.40 and then we'll have

$$n_o^2 [\rho^2 + z_o^2 - 2zz_o] + n_i^2 [\rho^2 + z_i^2 - 2zz_i] - 2n_o n_i [\rho^2 + z_o^2 - 2zz_o]^{1/2} [\rho^2 + z_i^2 - 2zz_i]^{1/2} = (n_i z_i - n_o z_o)^2$$

Now, we're going to re-order all the equation for convenience as follows

$$n_o^2 [\rho^2 + z_o^2 - 2zz_o] + n_i^2 [\rho^2 + z_i^2 - 2zz_i] - (n_i z_i - n_o z_o)^2 = 2n_o n_i [\rho^2 + z_o^2 - 2zz_o]^{1/2} [\rho^2 + z_i^2 - 2zz_i]^{1/2},$$

on the left hand side we have got the following

$$\begin{aligned} & n_o^2 [\rho^2 + z_o^2 - 2zz_o] + n_i^2 [\rho^2 + z_i - 2zz_i] - (n_i z_i - n_o z_o)^2 = \\ & n_o^2 [\rho^2 + z_o^2 - 2zz_o] + n_i^2 [\rho^2 + z_i - 2zz_i] - (n_i^2 z_i^2 + n_o^2 z_o^2 - 2n_o n_i z_o z_i) = \\ & n_o^2 [\rho^2 - 2zz_o] + n_i^2 [\rho^2 - 2zz_i] + 2n_o n_i z_o z_i, \end{aligned}$$

where we just only simplified the common terms. Having this on count, the equation reduces to

$$n_o^2 [\rho^2 - 2zz_o] + n_i^2 [\rho^2 - 2zz_i] + 2n_o n_i z_o z_i = 2n_o n_i [\rho^2 + z_o^2 - 2zz_o]^{1/2} [\rho^2 + z_i^2 - 2zz_i]^{1/2}. \quad (2.41)$$

Now, we must square again, the whole equation as follows

$$\begin{aligned} & [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)]^2 + (2n_o n_i z_o z_i)^2 + 4n_o n_i z_o z_i [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)] = \\ & 4n_o^2 n_i^2 (\rho^2 + z_o^2 - 2zz_o) (\rho^2 + z_i^2 - 2zz_i), \end{aligned}$$

now we are going to expand the product on the right hand side in the following way

$$\begin{aligned} & [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)]^2 + (2n_o n_i z_o z_i)^2 + 4n_o n_i z_o z_i [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)] = \\ & 4n_o^2 n_i^2 (\rho^2 - 2zz_o) (\rho^2 - 2zz_i) + 4n_o^2 n_i^2 [z_o^2 (\rho^2 - 2zz_i) + z_i^2 (\rho^2 - 2zz_o) + z_o^2 z_i^2]. \end{aligned}$$

Now, we must do a simplification, to do this, we must expand the first term of the left hand side like this

$$[n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)]^2 = [n_o^2 (\rho^2 - 2zz_o)]^2 + [n_i^2 (\rho^2 - 2zz_i)]^2 + 2n_o^2 n_i^2 (\rho^2 - 2zz_o) (\rho^2 - 2zz_i),$$

so the first term of the left hand side have on common in its expansion the same term than the expansion of the term on the right hand side, so, we could simplify then and look at the binomial is the same but with a minus sign instead of a plus sign, thus

$$\begin{aligned} & [n_o^2 (\rho^2 - 2zz_o) - n_i^2 (\rho^2 - 2zz_i)]^2 + (2n_o n_i z_o z_i)^2 + 4n_o n_i z_o z_i [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)] = \\ & 4n_o^2 n_i^2 [z_o^2 (\rho^2 - 2zz_i) + z_i^2 (\rho^2 - 2zz_o) + z_o^2 z_i^2]. \end{aligned}$$

Now, we are going to forget, momentarily the first term of the left hand side and focus on all the rest of the equations because there are going to be many important simplifications. Then, having this on count we'll have the following:

$$(2n_o n_i z_o z_i)^2 + 4n_o n_i z_o z_i [n_o^2 (\rho^2 - 2zz_o) + n_i^2 (\rho^2 - 2zz_i)] - 4n_o^2 n_i^2 [z_o^2 (\rho^2 - 2zz_i) + z_i^2 (\rho^2 - 2zz_o) + z_o^2 z_i^2] = 0$$

$$4n_on_i [z_oz_in_o^2 (\rho^2 - 2zz_o) + z_oz_in_i^2 (\rho^2 - 2zz_i) - n_on_iz_o^2 (\rho^2 - 2zz_i) - n_on_iz_i^2 (\rho^2 - 2zz_o) - n_on_iz_o^2 z_i^2 + n_on_iz_o^2 z_i^2],$$

now, we have to factorize the common factors

$$\begin{aligned} & 4n_on_i [(\rho^2 - 2zz_o) (z_oz_in_o^2 - n_on_iz_i^2) + (\rho^2 - 2zz_i) (z_oz_in_i^2 - n_on_iz_o^2)] = \\ & 4n_on_i [z_in_o (\rho^2 - 2zz_o) (z_on_o - n_iz_i) + z_on_i (\rho^2 - 2zz_i) (z_in_i - z_on_o)] = \\ & 4n_on_i [n_iz_i - n_oz_o] [n_iz_o (\rho^2 - 2zz_i) - n_oz_i (\rho^2 - 2zz_o)] = \\ & 4n_on_i [n_iz_i - n_oz_o] [\rho^2 (n_iz_o - n_oz_i) + 2zz_iz_o (n_o - n_i)]. \end{aligned}$$

So, joining this result with the first term we will have the following

$$[n_o^2 (\rho^2 - 2zz_o) - n_i^2 (\rho^2 - 2zz_i)]^2 + 4n_on_i [n_iz_i - n_oz_o] [\rho^2 (n_iz_o - n_oz_i) + 2zz_iz_o (n_o - n_i)] = 0,$$

so, the last things we have to do are is re organize in the following way

$$[(n_i^2 - n_o^2) \rho^2 - 2z (n_i^2 z_i - n_o^2 z_o)]^2 - 4n_in_o [n_iz_i - n_oz_o] [(n_oz_i - n_iz_o) \rho^2 + 2zz_oz_i (n_i - n_o)] = 0. \quad (2.42)$$

The equation 2.42 is the algebraic expression for the Descartes Ovoid the rigorously stigmatic surface we wanted.

2.4 The spherical diopter

Now, we are going to see that the Descartes Ovoid 2.42 contains the information of all the other diopeters known. So, the first one we are going to deduce is the spherical diopter this is obtained in a vicinity of the vertex shown in the figure 2.13. Therefore, to deduce this particular case, we are going to take the equation 2.42 and rewrite in the following convenient way

$$\left[(n_o^2 - n_i^2) \rho^2 + 2zz_oz_i \left(\frac{n_i^2}{z_o} - \frac{n_o^2}{z_i} \right) \right]^2 - 4n_in_o [n_iz_i - n_oz_o] [(n_oz_i - n_iz_o) \rho^2 + 2zz_oz_i (n_i - n_o)] = 0. \quad (2.43)$$

So, the main idea is to construct the sphere that is located in the vertex v . To do this, we must see that the squared term in 2.43 looks very similar to the right factor term in the second term, so, we want to equate them and then simplify, which leads us to the following equations

$$n_o^2 - n_i^2 = c (n_oz_i - n_iz_o), \quad (2.44)$$

$$\frac{n_i^2}{z_o} - \frac{n_o^2}{z_i} = c (n_i - n_o) \quad (2.45)$$

where c is a constant we must calculate. The second thing we are going to do, is to re-order the equation 2.44 in the following way

$$n_o^2 - cz_i n_o = n_i^2 - cz_o n_i, \quad (2.46)$$

and if we re-order the equation 2.45 we'll have

$$\begin{aligned} z_i n_i^2 - z_o n_o^2 &= cz_o z_i (n_i - n_o), \\ z_i n_i^2 - cz_o z_i n_i &= z_o n_o^2 - cz_o z_i n_o, \\ z_o (n_o^2 - cz_i n_o) &= z_i (n_i^2 - cz_o n_i). \end{aligned} \quad (2.47)$$

So if we use 2.46 on 2.47 we have got the following two equations

$$(z_o - z_i) (n_i^2 - cz_o n_i) = 0, \quad (2.48)$$

$$(z_o - z_i) (n_o^2 - cz_i n_o) = 0, \quad (2.49)$$

using the equations 2.48 and 2.49 we will have

$$z_o = \frac{n_i}{c}, \quad z_i = \frac{n_o}{c}, \quad (2.50)$$

and using the equation 2.44 we will have

$$z_o = z_i \quad z_{0,i} = \frac{n_o + n_i}{c}. \quad (2.51)$$

Now, before going further analyzing this results, we have to use the equations 2.44, 2.45 on 2.43

$$\begin{aligned} [c(n_o z_i - n_i z_o) \rho^2 + 2zz_o z_i c(n_i - n_o)]^2 - 4n_i n_o [n_i z_i - n_o z_o] [(n_o z_i - z_i z_o) \rho^2 + 2zz_o z_i (n_o - n_o)] &= 0, \\ [(n_o z_i - n_i z_o) c \rho^2 + 2zz_o z_i c(n_i - n_o)] [(n_o z_i - n_i z_o) \rho^2 + 2zz_o z_i (n_i - n_o) - 4n_i n_o (n_i z_i - n_o z_o)] &= 0, \end{aligned}$$

and by the fundamental theorem of Algebra, we have got the equation for the spherical diopter

$$(n_o z_i - n_i z_o) \rho^2 + 2zz_o z_i (n_i - n_o) = 0. \quad (2.52)$$

2.4.1 The Young-Weirstrass points

From the case 2.50 we have that

$$n_i z_i - n_o z_o = 0,$$

and using the equation 2.52 we will have

$$\rho^2 (n_o^2 - n_i^2) + 2cz_o n_i (n_i - n_o) = 0, \quad (2.53)$$

now, dividing by $n_o^2 - n_i^2$ and then we must complete squares as follows

$$x^2 + y^2 + z^2 + \frac{2czz_oz_i}{n_i + n_o} = x^2 + y^2 + z^2 + \frac{2czz_oz_i}{n_i + n_o} + \left(\frac{cz_oz_i}{n_i + n_o} \right)^2 = \left(\frac{cz_oz_i}{n_i + n_o} \right)^2,$$

$$x^2 + y^2 + \left(z - c \frac{z_oz_i}{n_o + n_i} \right)^2 = c^2 \frac{z_o^2 z_i^2}{(n_i + n_o)^2}, \quad (2.54)$$

which corresponds to the equation of an sphere whose radius is given by

$$R = c \frac{z_oz_i}{n_o + n_i} \rightarrow x^2 + y^2 + (z - R)^2 = R^2, \quad (2.55)$$

and using 2.50 $z_o = n_i/c$, $z_i = n_o/c$ we will have

$$R = c \frac{z_in_i}{c(n_o + n_i)} \rightarrow z_i = R \left(1 + \frac{n_o}{n_i} \right), \quad (2.56)$$

$$R = c \frac{z_on_o}{c(n_o + n_i)} \rightarrow z_i = R \left(1 + \frac{n_i}{n_o} \right) \quad (2.57)$$

which are known as **Young-Weirstrass** points or the aplanetic points. One of these points is inside and the other outside the sphere as is shown in 2.14.

2.4.2 The second spherical diopter

So, for the second case we have 2.51 and the equation 2.52 then let z_c given by

$$z_c = z_o = z_i = \frac{n_i + n_o}{c},$$

then we will have

$$\rho^2 (n_oz_c - n_iz_c) + 2zz_c^2(n_i - n_o) = 0, \quad (2.58)$$

$$\rho^2 - 2zz_c = 0, \quad (2.59)$$

again we must complete squares in the following way

$$x^2 + y^2 + z^2 - 2zz_c + z_c^2 = z_c^2, \quad (2.60)$$

$$x^2 + y^2 + (z - z_c)^2 = z_c^2. \quad (2.61)$$

The equation 2.61 is an spherical diopter where the object point is on the center and the image point is also on the center.




Figure 2.14: The spherical Diopter for the first case, where the object point is real and it is outside the sphere but the image point is virtual and inside the diopter.

2.5 Objects and images to the infinity

Some other cases arise when either the object or the image is at infinity, which is a convenient condition when the object is very far away, for example, in telescopes.

But even in microscopy, systems are designed to form the image at infinity, since under these conditions, an eyepiece facilitates visual accommodation for observation.

For an object in the infinity we divide by z_o^2 and take the limit $z_o \rightarrow -\infty$ in 2.42 in the following way

$$\begin{aligned}
& \lim_{z_o \rightarrow -\infty} \left[\frac{\rho^2 (n_o^2 - n_i^2)}{z_o} + 2z \frac{n_o^2 z_o + n_i^2 z_i}{z_o} \right]^2 - \lim_{z_o \rightarrow -\infty} \frac{4n_o n_i (n_o z_o + n_i z_i)}{z_o} \left[\frac{\rho^2 (n_o z_i + n_i z_o)}{z_o} \right. \\
& \quad \left. - \frac{2z z_o z_i (n_i - n_o)}{z_o} \right] = 0, \\
& (2z n_o^2)^2 - 4n_o 2n_i^2 [\rho^2 n_i - 2z z_i (n_i - n_o)] = 0, \\
& 4z^2 n_o^4 - 4n_o^2 n_i^2 \rho^2 + 8z z_i n_o^2 n_i^2 - 8z z_i n_o^3 n_i = 0, \quad \text{Dividing all the equation by } 4n_o^2, \\
& z^2 (n_o^2 - n_i^2) + 2z n_i z_i (n_i - n_o) - n_i (x^2 + y^2) = 0, \\
& [n_o - n_i] [z^2 (n_o + n_i) - 2z n_i z_i] - n_i (x^2 + y^2) = 0.
\end{aligned}$$

Now, to complete the algebraic expression we must multiply all the equation by $(n_o + n_i)$ and complete the squares as follows

$$\begin{aligned}
& (n_o - n_i) [(n_o + n_i)(n_o + n_i)z^2 - 2z n_i z_i (n_o + n_i)] - n_i^2 (n_o + n_i)(x^2 + y^2) = 0, \\
& (n_o - n_i) [(n_o + n_i)^2 z^2 - 2z n_i z_i (n_o + n_i) + n_i^2 z_i^2 - n_i^2 (n_o + n_i)(x^2 + y^2)] = (n_o - n_i) n_i^2 z_i^2, \\
& (n_o - n_i) [(n_o + n_i)z - n_i z_i]^2 - n_i^2 (n_o + n_i)(x^2 + y^2) = (n_o - n_i) n_i^2 z_i^2
\end{aligned}$$

and dividing all by $(n_o - n_i) n_i^2 z_i^2$ we will have

$$\left(\frac{n_i + n_o}{n_i z_i} \right)^2 \left[z - \frac{n_i z_i}{n_i + n_o} \right]^2 + \frac{n_i + n_o}{z_i^2 (n_i - n_o)} (x^2 + y^2) = 1. \quad (2.62)$$

The result 2.62 represents a conical section of revolution around the z-axis from this equation we are going to obtain the Ellipsoid and the hyperboloid.

2.5.1 Revolution Ellipsoid


if $n_i > n_o$ we are going to define the following scalars

$$a = \left(\frac{n_i - n_o}{n_i + n_o} \right)^{1/2} |z_i|, \quad b = \left(\frac{n_i}{n_i + n_o} \right) |z_i|, \quad (2.63)$$

and notice that $b > a$ so we will have **prolate ellipsoid** (see figure 2.15 given by

$$\frac{1}{b^2} (z - b)^2 + \frac{1}{a^2} (x^2 + y^2) = 1. \quad (2.64)$$

We have got two cases, when $z_i > 0$ and $z_i < 0$. For both cases what is next is to study the eccentricity of the ellipse and the foci of the ellipsoids.

Figure 2.15: Prolate Ellipsoid centered and $z = B$.

When $z_i > 0$ lets define the segment $c = \overline{F_1 B}$ such that the eccentricity is given by

$$\epsilon = \frac{c}{b} = \frac{\sqrt{b^2 - a^2}}{b} = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{\frac{n_o^2}{n_i^2}} = \frac{n_o}{n_i} < 1, \quad (2.65)$$

therefore the foci F_1 is centered to a distance

$$f_1 = b - c = b - \epsilon b = (1 - \epsilon)b = \frac{n_i - n_o}{n_i + n_o} z_i, \quad (2.66)$$

and the second foci is at

$$f_2 = b + c = b + \epsilon b = (1 + \epsilon)b = z_i. \quad (2.67)$$


This means, that for this first case that all the rays that comes parallel to the z -axis from infinity converges to the second Foci of the ellipsoid.

If $z_i < 0$ by the convention we have that $\sqrt{z_i^2} = -z_i$ and we define

$$b = -\frac{n_i}{n_i + n_o} z_i, \quad (2.68)$$

then the conical section equation 2.62 becomes

$$\frac{1}{b^2}(z + b)^2 + \frac{1}{a^2}(x^2 + y^2) = 1. \quad (2.69)$$

Figure 2.16: Ellipsoid diopter when $n_i > n_o$.

Again, we define $c = \overline{F_1 B}$ so the eccentricity is given by

$$\epsilon = \frac{c}{b} \frac{\sqrt{b^2 - a^2}}{b} = \sqrt{1 - \frac{a^2}{b^2}} = \frac{n_o}{n_i} < 1,$$

and the foci is now going to be to a distance

$$f_1 = -b - c = -b - \epsilon b = -b(1 + \epsilon) = z_i, \quad (2.70)$$


and the second one is at

$$f_2 = -b + c = -b + \epsilon b = (\epsilon - 1)b = \frac{n_i - n_o}{n_i + n_o} z_i. \quad (2.71)$$

Having all these results on count, we can see in figure 2.17 that all the rays that becomes parallel to the z -axis and from infinity diverges as they would come from a point source from F_1 .

2.5.2 Revolution Hyperboloid

In the case $n_i < n_o$ we have got that the equation 2.62 becomes an hyperboloid and, again, there are two possibilities, $z_i < 0, z_i > 0$, so the conical equation becomes

Figure 2.17: Ellipsoid diopter when $n_i > n_o$ and $z_i < 0$.

$$\left(\frac{n_i + n_o}{n_i z_i} \right)^2 \left[z - \frac{n_i z_i}{n_i + n_o} \right] - \frac{n_i + n_o}{z_i^2 (n_o - n_i)} (x^2 + y^2) = 1. \quad (2.72)$$

If $z_i > 0$ we define again the following scalars

$$b^2 = \frac{n_i^2}{(n_i + n_o)^2} z_i^2,$$

$$a^2 = \frac{n_o - n_i}{(n_i + n_o)^2} z_i^2,$$

so the equation of the hyperboloid (see figure 2.18) becomes

$$\frac{1}{b^2} (z - b)^2 - \frac{1}{a^2} (x^2 + y^2) = 1. \quad (2.73)$$

Again, to give a characterization of the conical surface we are going to define the segment $c = \overline{F_1 B}$ such that the eccentricity is given by

$$\epsilon = \frac{c}{b} = \frac{\sqrt{a^2 + b^2}}{b} = \sqrt{1 + \frac{a^2}{b^2}} = \frac{n_o}{n_i} > 1.$$




Figure 2.18: Geometrical representation of the two leaf hyperboloid with the parameters a, b and foci F_1 and F_2 .

Thus the two Foci F_1 and F_2 are located at the distances

$$f_1 = b - c = b - \epsilon b = (1 - \epsilon)b = \frac{n_i - n_o}{n_o + n_i} z_o, \quad (2.74)$$

$$f_2 = b + c = b + \epsilon b = (1 + \epsilon)b = z_i. \quad (2.75)$$

In the figure 2.19 we can see the geometrical behavior of this first case, all the rays comes from the infinity and converges to a point which is located in the second foci F_2 .

Now, for the second case, if $z_i < 0$ we define

$$b = -\frac{n_i}{n_i + n_o} z_o,$$

in such way that $b > 0$ with this we write

$$\frac{1}{b^2}(z + b)^2 - \frac{1}{a^2}(x^2 + y^2) = 1.$$

Again, we define the segment $c = \overline{F_1 B}$ such that the eccentricity is given by

$$\epsilon = \frac{c}{b} = \frac{\sqrt{a^2 + b^2}}{b} = \sqrt{1 + \frac{a^2}{b^2}} = \frac{n_o}{n_i} > 1.$$




Figure 2.19: Geometrical representation of the first case of a two leaf hyperboloid.

As we can see in the figure 2.20 all the rays come from the infinity and interact with the hyperboloid diopter and the rays refracted in such geometrical way that they behave as they come from a object point source located in the foci of the first leaf at F_1 .

2.6 Rigurously stigmatic surfaces by reflection

The reflective surfaces are the main components in the modern giant telescopes, for example, the James Webb telescope is an array of many reflective surfaces which are electronically displaced such that the image formation gets the least aberration possible. Also, the reflective surfaces are better than the refractive ones in certain cases, because the optical systems can be smaller and don't have chromatic aberrations.

Traditionally, the reflective surface are developed independently from the refractive surface. Under this formulations, the rigorously stigmatic surfaces by reflection can be obtained doing $n_i = -n_o$, where the Descartes ovals become a conic surface. To this case, the equation 2.42 becomes

$$\left\{ \left[(-n_o)^2 - n_o^2 \right] \rho^2 - 2z \left[(n_o^2 z_o - (-n_o)^2 z_i) \right] \right\}^2 - 4(-n_o)n_o [(-n_o)z_i - n_o z_o] \{ [n_o z_i - (-n_o)z_o] \rho^2 + 2z z_o z_i [(-n_o) - n_o] \} = 0,$$

is positive and the surface closes onto itself as an ellipsoid, whereas when $z_o z_i < 0$ that coefficient turns negative and the surface opens as a hyperboloid. There is also a beautiful limit waiting in 2.76, because if we push the object to infinity, $z_o \rightarrow -\infty$, the conic degenerates into a paraboloid, which is the case that rules every telescope. We are going to walk through each of these situations, since each one corresponds to a real optical instrument.

2.6.1 The ellipsoidal mirror




Figure 2.21: Ellipsoidal mirror. The object F_1 and the image F_2 coincide with the two foci of the ellipse, so every ray leaving one focus is reflected exactly onto the other.

Lets start with $z_o z_i > 0$, so that the object and the image sit on the same side of the vertex and the mirror is an ellipsoid. Dividing 2.76 by $\left(\frac{z_o + z_i}{2}\right)^2$ the equation takes the canonical shape

$$\frac{\left(z - \frac{z_o + z_i}{2}\right)^2}{\left(\frac{z_o + z_i}{2}\right)^2} + \frac{x^2 + y^2}{z_o z_i} = 1, \quad (2.77)$$

and if we baptise the two semi-axes as

$$b^2 = \frac{(z_o + z_i)^2}{4}, \quad a^2 = z_o z_i, \quad (2.78)$$

we recognise a prolate ellipsoid centered at $z = b = \frac{z_o + z_i}{2}$, that is

$$\frac{1}{b^2} (z - b)^2 + \frac{1}{a^2} (x^2 + y^2) = 1. \quad (2.79)$$

That the ellipsoid comes out prolate, and not oblate, is not an accident, because

$$b^2 - a^2 = \frac{(z_o + z_i)^2}{4} - z_o z_i = \frac{(z_o - z_i)^2}{4} \geq 0,$$

so the semi-axis along the optical axis is always the largest one and the mirror is stretched along z . The distance from the center to each focus is then $c = \sqrt{b^2 - a^2} = \frac{|z_o - z_i|}{2}$, and the eccentricity becomes

$$\epsilon = \frac{c}{b} = \frac{\sqrt{b^2 - a^2}}{b} = \sqrt{1 - \frac{a^2}{b^2}} = \frac{|z_o - z_i|}{z_o + z_i} < 1, \quad (2.80)$$

which confirms once more that we are dealing with an ellipse. The really interesting thing is where the foci land, because measured from the vertex they sit at $z = b \pm c$, that is

$$b - c = \frac{z_o + z_i}{2} - \frac{|z_o - z_i|}{2}, \quad b + c = \frac{z_o + z_i}{2} + \frac{|z_o - z_i|}{2}, \quad (2.81)$$

and no matter which of the two conjugate points is the larger one, these two numbers are nothing but z_o and z_i themselves. **The two foci of the ellipsoid are exactly the object point and the image point.**

This is the whole magic of the ellipsoidal mirror and the reason it is rigorously stigmatic. An ellipse has the geometric property that any ray leaving one focus is reflected straight into the other focus, because the sum of the two focal distances is constant along the whole curve, and a constant optical path is exactly the Fermat's principle at work. Placing a point source at one focus therefore produces a perfect point image at the other one, with no aberration at all. This is the principle behind the elliptical reflectors used to concentrate the light coming from one focus onto a detector sitting at the other, and it is the same property that makes the whispering galleries carry a voice from one focus of the room to the other.

When both z_o and z_i are negative the algebra is identical once we define $b = -\frac{z_o + z_i}{2} > 0$, so that the ellipsoid is now centered at $z = -b$ and its equation reads $\frac{1}{b^2}(z + b)^2 + \frac{1}{a^2}(x^2 + y^2) = 1$. The eccentricity is the same and the foci again fall on z_o and z_i , the only difference being that both conjugate points now live on the opposite side of the vertex.

2.6.2 The hyperboloidal mirror

Now we let the product change sign, $z_o z_i < 0$, which happens when the object and the image sit on opposite sides of the mirror and one of them is virtual. The coefficient of $x^2 + y^2$ in 2.76 becomes negative, so writing $a^2 = -z_o z_i > 0$ while keeping $b^2 = \frac{(z_o + z_i)^2}{4}$ the canonical form is now

$$\frac{1}{b^2}(z - b)^2 - \frac{1}{a^2}(x^2 + y^2) = 1, \quad (2.82)$$

a hyperboloid of two sheets around the optical axis. The focal distance is measured with a plus sign this time, $c = \sqrt{a^2 + b^2}$, and a short computation gives again $c = \frac{|z_o - z_i|}{2}$, so the eccentricity is

$$\epsilon = \frac{c}{|b|} = \frac{\sqrt{a^2 + b^2}}{|b|} = \sqrt{1 + \frac{a^2}{b^2}} = \frac{|z_o - z_i|}{|z_o + z_i|} > 1, \quad (2.83)$$




Figure 2.22: Hyperboloid mirror. The rays from the object F_1 diverge as they were emitted by a virtual image located in F_2 .

which is greater than one exactly because $z_o z_i < 0$ forces $|z_o - z_i| > |z_o + z_i|$. And just as before, the foci $z = b \pm c$ collapse onto z_o and z_i , so once more the two conjugate points are the two foci of the conic.

The hyperboloidal mirror is therefore stigmatic between its two foci as well, but with a twist, because since the foci lie on opposite sheets one of the conjugate points is always virtual. This is precisely the behavior exploited by the convex secondary mirror of a Cassegrain telescope, where the light that is already converging toward the focus of the primary mirror is intercepted by a hyperboloidal secondary and redirected, without adding any aberration, to a more comfortable focus behind the primary. The stigmatic pair of the hyperboloid is what lets these compact folded telescopes keep a perfect image on the axis.

2.6.3 Special cases: the spherical and the plane mirror

Two very familiar mirrors are hiding inside 2.76 as degenerate cases. The first one appears when the object and the image coincide, $z_o = z_i = R$. The coefficient of $x^2 + y^2$ becomes $\frac{(2R)^2}{4R^2} = 1$ and the conic reduces to $z^2 - 2Rz + x^2 + y^2 = 0$, and completing the square by adding R^2 to both sides we get

$$(z - R)^2 + x^2 + y^2 = R^2, \quad (2.84)$$

which is simply a sphere of radius R centered at $z = R$. This tells us that a spherical mirror is rigorously stigmatic for one very special pair of points, both of them sitting together at the center of the sphere, since a source placed at the center sends every ray along a radius, the ray strikes the surface perpendicularly and comes straight back on itself, forming a perfect image right on top of the source. For any other position the sphere is only approximately stigmatic, and this is the origin of the spherical aberration that spoils cheap mirrors and that the conics above were designed to cure.

The second degenerate case is $z_o = -z_i$, for which the sum $z_o + z_i$ vanishes. Both the linear term and the whole right hand side of 2.76 die at once, and the equation collapses to $z^2 = 0$, that is, the plane $z = 0$. We recover the ordinary plane mirror, which is stigmatic for every point and always produces an image symmetric to the object at $z_i = -z_o$, which is nothing more than the everyday statement that a plane mirror puts your reflection as far behind the glass as you stand in front of it.

2.6.4 The parabolic mirror and objects at infinity

The most important mirror of all comes from letting the object recede to infinity, $z_o \rightarrow -\infty$, which is the situation of a telescope pointed at a star. Going back to 2.76 and expanding the square on the left, the terms $\frac{(z_o + z_i)^2}{4}$ appear on both sides and cancel,

$$z^2 - z(z_o + z_i) + \frac{(z_o + z_i)^2}{4} + \frac{(z_o + z_i)^2}{4z_o z_i}(x^2 + y^2) = \frac{(z_o + z_i)^2}{4},$$

leaving the expanded conic

$$z^2 - z(z_o + z_i) + \frac{(z_o + z_i)^2}{4z_o z_i}(x^2 + y^2) = 0.$$

Dividing everything by z_o and taking the limit $z_o \rightarrow -\infty$, the term z^2/z_o vanishes, the factor $\frac{z_o + z_i}{z_o} \rightarrow 1$ and $\frac{(z_o + z_i)^2}{4z_o^2 z_i} \rightarrow \frac{1}{4z_i}$, so we are left with the remarkably clean surface

$$x^2 + y^2 = 4z_i z, \tag{2.85}$$

a paraboloid of revolution whose opening is fixed by the sign of z_i .

The paraboloid is the crown jewel of the reflective optics. A parabola has the defining property that every ray arriving parallel to its axis is reflected exactly through the focus, so a bundle of parallel rays coming from a star, which is an object at infinity, is gathered into a single perfect point at $z = z_i$, the focus of the mirror. There is no chromatic aberration, because the reflection does not depend on the wavelength, and there is no spherical aberration, because the surface is a true paraboloid instead of a sphere. This is why every serious reflecting telescope, from the first design of Newton to the segmented primary of the James Webb, uses a parabolic mirror as its light collecting surface, and it is also why the same shape reappears in the satellite dishes and the searchlights, where the ray tracing simply runs the other way and a source placed at the focus is turned into a parallel beam.




Figure 2.23: Parabolic mirror obtained as the limit $z_o \rightarrow -\infty$. The rays coming parallel to the axis from an object at infinity are all focused onto the single point $z = z_i$.

2.7 Explicit formulation of the Descartes ovoids

Everything we have built so far has left us with the Descartes ovoid written in the implicit form 2.42, that is, as the zero set of a quartic polynomial $F(x, y, z) = 0$. That form is beautiful for proving things, because every degenerate case we studied, the sphere, the ellipsoid, the hyperboloid, the paraboloid and the plane, was obtained simply by looking at what the polynomial does under a particular choice of n_o, n_i, z_o, z_i . But the implicit form is nearly useless for the person who has to actually build the surface. An optical designer, a lens grinder or a diamond turning machine does not want to know the set of points that annihilate a quartic, they want a height, they want to be told how much material to remove at a distance ρ from the axis. In other words, they want the *sag* of the surface, the explicit function $z(\rho)$.

So the goal of this section is purely practical, we want to solve 2.42 for z . The obstacle is that

the equation is quartic in z , and a brute force attack on a quartic is a nightmare. The trick, which is the whole content of this section, is to notice that the ovoid is secretly a much simpler object, a *superconic*, and that when written in superconic form the equation becomes quadratic in z , which we can solve with the formula we learned as children.

2.7.1 The Descartes ovoid as a superconic

A superconic is a surface of revolution around the optical axis defined by

$$c_0 G z^2 - 2(1 + b_1 \rho^2 + b_2 \rho^4 + \dots) z + (c_0 + c_1 \rho^2 + c_2 \rho^4 + \dots) \rho^2 = 0, \quad (2.86)$$

where G is the conic constant, also called the Schwarzschild constant (Schwarzschild, 1905), and c_0 is the paraxial curvature, that is, the curvature the surface has right at the vertex. If we kill every deformation coefficient, $b_j = c_j = 0$ for $j \geq 1$, the surface collapses to

$$c_0 G z^2 - 2z + c_0 \rho^2 = 0, \quad (2.87)$$

which is the standard conic of revolution. So a superconic is nothing but a conic that has been allowed to deform, term by term, in even powers of ρ .

Now comes the claim that makes everything work, *the Descartes ovoid is a superconic truncated at the very first deformation order*. To see it, take the ovoid 2.42

$$[(n_i^2 - n_o^2) \rho^2 - 2z(n_i^2 z_i - n_o^2 z_o)]^2 - 4n_i n_o (n_i z_i - n_o z_o) [(n_o z_i - n_i z_o) \rho^2 + 2z z_o z_i (n_i - n_o)] = 0,$$

and divide the whole equation by the constant

$$\mathcal{N} = 4n_i n_o z_i z_o (n_i - n_o) (n_i z_i - n_o z_o). \quad (2.88)$$

Let us do it piece by piece, because this is where the four shape parameters are born. The second term of the ovoid divided by $\mathcal{N}$ gives

$$\begin{aligned} & - \frac{4n_i n_o (n_i z_i - n_o z_o) [(n_o z_i - n_i z_o) \rho^2 + 2z z_o z_i (n_i - n_o)]}{4n_i n_o (n_i z_i - n_o z_o) z_i z_o (n_i - n_o)} = \\ & - \frac{(n_o z_i - n_i z_o) \rho^2}{z_i z_o (n_i - n_o)} - \frac{2z z_o z_i (n_i - n_o)}{z_i z_o (n_i - n_o)} = \frac{n_i z_o - n_o z_i}{z_i z_o (n_i - n_o)} \rho^2 - 2z, \end{aligned}$$

which already has the shape $O\rho^2 - 2z$ of the superconic 2.87 if we baptise

$$O = \frac{n_i z_o - n_o z_i}{z_i z_o (n_i - n_o)}. \quad (2.89)$$

The first term of the ovoid is a perfect square, so we expand it before dividing,

$$\begin{aligned} & [(n_i^2 - n_o^2) \rho^2 - 2z(n_i^2 z_i - n_o^2 z_o)]^2 = \\ & (n_i^2 - n_o^2)^2 \rho^4 - 4z(n_i^2 - n_o^2)(n_i^2 z_i - n_o^2 z_o) \rho^2 + 4z^2(n_i^2 z_i - n_o^2 z_o)^2, \end{aligned}$$

and now we divide each of these three pieces by $\mathcal{N}$. The coefficient of z^2 becomes

$$\frac{4(n_i^2 z_i - n_o^2 z_o)^2}{4n_i n_o z_i z_o (n_i - n_o)(n_i z_i - n_o z_o)} = \underbrace{\frac{(n_i^2 z_i - n_o^2 z_o)^2}{n_i n_o (n_i z_i - n_o z_o)(n_i z_o - n_o z_i)}}_G \underbrace{\frac{n_i z_o - n_o z_i}{z_i z_o (n_i - n_o)}}_O = OG,$$

where we have simply multiplied and divided by $(n_i z_o - n_o z_i)$ in order to make the factor O appear. The coefficient of $z\rho^2$ becomes

$$\begin{aligned} & -\frac{4(n_i^2 - n_o^2)(n_i^2 z_i - n_o^2 z_o)}{4n_i n_o z_i z_o (n_i - n_o)(n_i z_i - n_o z_o)} = -\frac{(n_i + n_o)(n_i - n_o)(n_i^2 z_i - n_o^2 z_o)}{n_i n_o z_i z_o (n_i - n_o)(n_i z_i - n_o z_o)} = \\ & -2 \underbrace{\frac{(n_i + n_o)(n_i^2 z_i - n_o^2 z_o)}{2n_i n_o z_i z_o (n_i z_i - n_o z_o)}}_S, \end{aligned}$$

and finally the coefficient of ρ^4 becomes

$$\frac{(n_i^2 - n_o^2)^2}{4n_i n_o z_i z_o (n_i - n_o)(n_i z_i - n_o z_o)} = \frac{(n_i + n_o)^2 (n_i - n_o)^2}{4n_i n_o z_i z_o (n_i - n_o)(n_i z_i - n_o z_o)} = \underbrace{\frac{(n_i - n_o)(n_i + n_o)^2}{4n_i n_o z_i z_o (n_i z_i - n_o z_o)}}_T.$$

Collecting the four pieces, the Descartes ovoid has been transformed into

$$f(z, \rho) = OGz^2 - 2(1 + S\rho^2)z + (O + T\rho^2)\rho^2 = 0, \quad (2.90)$$

which is exactly the superconic 2.86 truncated at b_1 and c_1 , with $b_1 = S$ and $c_1 = T$. The four quantities we found along the way,

$$G = \frac{(n_i^2 z_i - n_o^2 z_o)^2}{n_i n_o (n_i z_i - n_o z_o)(n_i z_o - n_o z_i)}, \quad (2.91)$$

$$O = \frac{n_i z_o - n_o z_i}{z_i z_o (n_i - n_o)}, \quad (2.92)$$

$$T = \frac{(n_i - n_o)(n_i + n_o)^2}{4n_i n_o z_i z_o (n_i z_i - n_o z_o)}, \quad (2.93)$$

$$S = \frac{(n_i + n_o)(n_i^2 z_i - n_o^2 z_o)}{2n_i n_o z_i z_o (n_i z_i - n_o z_o)}, \quad (2.94)$$

are called the **shape parameters** *GOTS* of the ovoid, and they carry the whole physics of the surface, because the two refraction indices and the two conjugate positions enter the geometry only through these four numbers.




Figure 2.24: Rigorously stigmatic quartic surface described in the meridional plane by the polar parameters ρ and φ measured from the vertex $\mathcal{O}$. The object point A_o at z_o and the image point A_i at z_i are the strictly stigmatic pair, and at the incidence point I the vectorial Snell-Descartes law $n_o \hat{u} - n_i \hat{u}' = \kappa \hat{N}$ is satisfied.

These four parameters are not independent. If we compute the product GOT we find

$$GOT = \left[\frac{(n_i^2 z_i - n_o^2 z_o)^2}{n_i n_o z_i z_o (n_i - n_o) (n_i z_i - n_o z_o)} \right] \left[\frac{(n_i - n_o) (n_i + n_o)^2}{4 n_i n_o z_i z_o (n_i z_i - n_o z_o)} \right] =$$

$$\frac{(n_i + n_o)^2 (n_i^2 z_i - n_o^2 z_o)^2}{4 n_i^2 n_o^2 z_i^2 z_o^2 (n_i z_i - n_o z_o)^2} = \left[\frac{(n_i + n_o) (n_i^2 z_i - n_o^2 z_o)}{2 n_i n_o z_i z_o (n_i z_i - n_o z_o)} \right]^2 = S^2,$$

where in the first line we already wrote OG as the single fraction we obtained above. Therefore the shape parameters obey the constraint

$$S^2 = GOT, \quad (2.95)$$

and this ligature is precisely what distinguishes a Descartes ovoid from a generic superconic. A superconic with four freely chosen parameters is just a deformed conic with no particular optical virtue, while a superconic whose parameters satisfy 2.95 is rigorously stigmatic. The constraint is therefore the algebraic fingerprint of perfect imaging.

It is worth pausing on what the constraint costs us. We have four parameters and one equation between them, so only three are free, and this matches the physical counting, because the surface is determined by the ratio of the indices and by the two conjugate distances, that is, by three independent numbers. Notice also that if we send either conjugate point to infinity, $z_o \rightarrow \infty$ or $z_i \rightarrow \infty$, both S and T carry a $z_i z_o$ in the denominator and therefore vanish, and 2.90 degenerates

into the standard conic 2.87. This is the algebraic version of everything we found in the previous sections, the conics are the ovoids with one conjugate point at infinity.

2.7.2 The explicit formulation

Now that the ovoid is quadratic in z , the rest is elementary. Reading 2.90 as $az^2 + bz + c = 0$ with $a = OG$, $b = -2(1 + S\rho^2)$ and $c = (O + T\rho^2)\rho^2$, the quadratic formula gives

$$z = \frac{2(1 + S\rho^2) \pm \sqrt{4(1 + S\rho^2)^2 - 4OG(O + T\rho^2)\rho^2}}{2OG} = \frac{(1 + S\rho^2) \pm \sqrt{(1 + S\rho^2)^2 - OG(O + T\rho^2)\rho^2}}{OG},$$

and expanding the discriminant

$$\begin{aligned} (1 + S\rho^2)^2 - OG(O + T\rho^2)\rho^2 &= 1 + 2S\rho^2 + S^2\rho^4 - O^2G\rho^2 - OG T\rho^4 = \\ &= 1 + (2S - O^2G)\rho^2 + (S^2 - GOT)\rho^4, \end{aligned}$$

so that the sag of the surface is

$$z = \frac{1 + S\rho^2 \pm \sqrt{1 + (2S - O^2G)\rho^2 + (S^2 - GOT)\rho^4}}{OG}. \quad (2.96)$$

Two things must be settled now. The first one is which of the two signs is the physical branch. The surface has to pass through the vertex $\mathcal{O}$, which is the origin of our coordinates, therefore we must demand $z \rightarrow 0$ when $\rho \rightarrow 0$. Putting $\rho = 0$ in 2.96 gives $z = (1 \pm 1)/OG$, which is $2/OG$ for the plus sign and 0 for the minus sign. Hence *the Descartes ovoid is the branch with the minus sign*, and the branch with the plus sign is the outer sheet of the quartic, the second surface hidden inside the same polynomial.

The second thing is the ligature 2.95. Because $S^2 - GOT = 0$, the quartic term inside the radical dies and the expression collapses to

$$z = \frac{1}{OG} \left(1 + S\rho^2 - \sqrt{1 + (2S - O^2G)\rho^2} \right). \quad (2.97)$$

This is already explicit, but it is numerically ugly, because for small ρ it is the difference of two numbers that are both close to one, divided by OG , which can itself be small. The cure is to rationalise, multiplying and dividing by the conjugate,

$$\begin{aligned} z &= \frac{1}{OG} \frac{\left(1 + S\rho^2 - \sqrt{1 + (2S - O^2G)\rho^2} \right) \left(1 + S\rho^2 + \sqrt{1 + (2S - O^2G)\rho^2} \right)}{1 + S\rho^2 + \sqrt{1 + (2S - O^2G)\rho^2}} = \\ &= \frac{1}{OG} \frac{(1 + S\rho^2)^2 - [1 + (2S - O^2G)\rho^2]}{1 + S\rho^2 + \sqrt{1 + (2S - O^2G)\rho^2}}, \end{aligned}$$

and the numerator simplifies beautifully,

$$\begin{aligned} 1 + 2S\rho^2 + S^2\rho^4 - 1 - 2S\rho^2 + O^2G\rho^2 &= S^2\rho^4 + O^2G\rho^2 = \\ \rho^2 (O^2G + S^2\rho^2) &= \rho^2 (O^2G + GOT\rho^2) = OG\rho^2 (O + T\rho^2), \end{aligned}$$

where in the second line we used the ligature $S^2 = GOT$ once more. The factor OG cancels against the one sitting outside, and we arrive at the result we were after,

$$z(\rho) = \frac{(O + T\rho^2) \rho^2}{1 + S\rho^2 + \sqrt{1 + (2S - O^2G) \rho^2}}. \quad (2.98)$$

Equation 2.98 is the explicit formulation of the Descartes ovoids. It is worth appreciating how much has happened here. We started from a quartic implicit surface and we ended with a closed formula that gives the height of the surface directly in terms of the radial coordinate and four numbers built out of n_o, n_i, z_o and z_i . There is no series, no iteration and no numerical root finding, the expression is exact.

Since our radial variable was defined as $\rho^2 = x^2 + y^2 + z^2$, that is, the distance from the vertex to the point on the surface and not the transverse height, we recover the transverse coordinate from

$$r(\rho) = \pm \sqrt{\rho^2 - z^2(\rho)}, \quad r^2 = x^2 + y^2, \quad (2.99)$$

so the pair $(z(\rho), r(\rho))$ traces the meridional profile of the ovoid, and the full surface is obtained by revolving that profile around the optical axis.




Figure 2.25: Meridional sections of Cartesian surfaces with the refraction indices fixed at $n_o = 1$ and $n_i = 1.7$ while the object and image positions are varied. The solid curves are the solutions with the minus sign in 2.96, which pass through the origin and are the Descartes ovoids, and the dashed curves are the solutions with the plus sign, the outer sheet of the same quartic. (a) $z_o = -10$ and $z_i = 20$. (b) $z_o = -10$ and $z_i = -20$. (c) $z_o = 10$ and $z_i = 20$. (d) $z_o = 10$ and $z_i = -20$.

In the figure 2.25 we can see how the two branches behave. The solid curves, which are the ones we selected, always start at the origin, while the dashed ones close the quartic from outside, and together they show how the surface makes a transition from the inner to the outer sheet. The four

panels also show that the formulation handles without any modification the cases where the object or the image is virtual, that is, the cases where $z_o > 0$ or $z_i < 0$, which is exactly the situation that the older explicit formulations in the literature were unable to treat in a unified way. Depending on the signs, the surface opens to the right or to the left, and both are legitimate optical surfaces.

Surface	n_i	z_o	z_i	G	O	T	S
Ellipsoidal refractive	$n_i > n_o$	∞	z_i	$-\left(\frac{n_o}{n_i}\right)^2$	$\frac{n_i}{z_i(n_i-n_o)}$	0	0
Hyperboloidal refractive	$n_i < n_o$	∞	z_i	$-\left(\frac{n_o}{n_i}\right)^2$	$\frac{n_i}{z_i(n_i-n_o)}$	0	0
Hyperboloidal refractive	$n_i > n_o$	z_o	∞	$-\left(\frac{n_i}{n_o}\right)^2$	$-\frac{n_o}{z_o(n_i-n_o)}$	0	0
Ellipsoidal refractive	$n_i < n_o$	z_o	∞	$-\left(\frac{n_i}{n_o}\right)^2$	$-\frac{n_o}{z_o(n_i-n_o)}$	0	0
Spherical refractive	n_i	z_o	z_o	$\frac{(n_i+n_o)^2}{n_i n_o}$	$\frac{1}{z_o}$	$\frac{(n_i+n_o)^2}{4n_i n_o z_o^3}$	$\frac{(n_o+n_i)^2}{2n_i n_o z_o^2}$
Ellipsoidal reflective	$-n_o$	$z_o > 0$	$z_i > 0$	$-\left(\frac{z_i-z_o}{z_i+z_o}\right)^2$	$\frac{z_i+z_o}{2z_i z_o}$	0	0
Ellipsoidal reflective	$-n_o$	$z_o < 0$	$z_i < 0$	$-\left(\frac{z_i-z_o}{z_i+z_o}\right)^2$	$\frac{z_i+z_o}{2z_i z_o}$	0	0
Hyperboloidal reflective	$-n_o$	$z_o > 0$	$z_i < 0$	$-\left(\frac{z_i-z_o}{z_i+z_o}\right)^2$	$\frac{z_i+z_o}{2z_i z_o}$	0	0
Hyperboloidal reflective	$-n_o$	$z_o < 0$	$z_i > 0$	$-\left(\frac{z_i-z_o}{z_i+z_o}\right)^2$	$\frac{z_i+z_o}{2z_i z_o}$	0	0
Paraboloidal reflective	$-n_o$	∞	z_i	-1	$\frac{1}{2z_i}$	0	0
Paraboloidal reflective	$-n_o$	z_o	∞	-1	$\frac{1}{2z_o}$	0	0

Figure 2.26: Degenerate cases of refractive and reflective rigorously stigmatic surfaces according to the refraction indices and the object and image positions. For each combination of these parameters the values of the shape parameters $GOTS$ of the corresponding conic of revolution are obtained. Note that $T = S = 0$ in every conic case, and that the reflective surfaces are recovered by the substitution $n_i \rightarrow -n_o$.

The table 2.26 closes the circle with everything we did in the previous sections. Every single surface we deduced by hand, the ellipsoid, the hyperboloid, the sphere, the paraboloid and their reflective versions, appears there as a particular choice of the shape parameters. Notice that in all the conic rows we have $T = S = 0$, which is the algebraic statement that one of the conjugate points has been sent to infinity, and notice that the only row where T and S survive is the spherical refractive one, where $z_o = z_i$, that is, the second spherical diopter 2.61 where object and image sit together at the centre of the sphere.

2.7.2.1 The Schwarzschild equation

There is one more limit worth writing down explicitly, because it connects our formula with the equation that every optical design program in the world uses. When $T = S = 0$, that is, for the conic surfaces, the expression 2.98 reduces to

$$z(\rho) = \frac{O\rho^2}{1 + \sqrt{1 - O^2G\rho^2}}, \quad (2.100)$$

which is still written in terms of the distance ρ from the vertex, and not in terms of the transverse height r . To change variables we use $\rho^2 = z^2 + r^2$ in the conic 2.87,

$$\begin{aligned} OGz^2 - 2z + O(z^2 + r^2) &= 0, \\ O(G + 1)z^2 - 2z + Or^2 &= 0, \end{aligned}$$

which is again quadratic in z , and solving it and choosing once more the branch through the vertex,

$$z = \frac{1 - \sqrt{1 - (G + 1)O^2r^2}}{O(G + 1)},$$

and rationalising exactly as before, the numerator becomes $1 - 1 + (G + 1)O^2r^2$, so that

$$z(r) = \frac{Or^2}{1 + \sqrt{1 - (K + 1)O^2r^2}}, \quad K = G, \quad (2.101)$$

which is the Schwarzschild formula for the standard conic surfaces (Schwarzschild, 1905), where $K = -e^2$ is the conic constant written in terms of the eccentricity and O is the curvature at the vertex.

This tells us how to read our own result. The equation 2.98 is a *generalisation of the Schwarzschild equation* for the Cartesian surfaces, where G generalises the Schwarzschild constant and we may call it the Descartes ovoid constant, O is the paraxial curvature of the ovoid at the vertex, and T is the parameter that labels the different families of ovoids. The parameter S adds nothing new, since the ligature 2.95 fixes it once the other three are given.

2.8 Special Descartes ovoids

With the shape parameters at hand we can now classify the ovoids in the same spirit in which one classifies the conics by their eccentricity. The classification is governed by G , and the reason is immediate from 2.91, since the numerator is a perfect square and $n_i n_o > 0$, the sign of G is nothing but the sign of the product

$$(n_i z_i - n_o z_o)(n_i z_o - n_o z_i), \quad (2.102)$$

so the whole taxonomy is decided by the relative position of the two conjugate points weighted by the indices.

2.8.1 Spherovoids, $G = 0$

From 2.91 the constant G vanishes if and only if the numerator vanishes,

$$(n_i^2 z_i - n_o^2 z_o)^2 = 0 \iff n_i^2 z_i = n_o^2 z_o, \quad (2.103)$$

and the very same factor sits in the numerator of S , so this condition forces $G = S = 0$ simultaneously. The surface is then characterised by O and T alone, and 2.98 becomes

$$z(\rho) = \frac{(O + T\rho^2)\rho^2}{1 + \sqrt{1}} = \frac{1}{2}(O + T\rho^2)\rho^2. \quad (2.104)$$

These are the **spherovoids**, and the name is justified by what happens if we additionally switch off T . With $T = 0$ we get $z = O\rho^2/2$, and substituting $\rho^2 = z^2 + r^2$,

$$\begin{aligned} 2z = O(z^2 + r^2) &\implies z^2 - \frac{2z}{O} + r^2 = 0, \\ z^2 - \frac{2z}{O} + \frac{1}{O^2} + r^2 &= \frac{1}{O^2} \implies \left(z - \frac{1}{O}\right)^2 + x^2 + y^2 = \frac{1}{O^2}, \end{aligned}$$

which is a sphere of radius $1/O$ centred at $z = 1/O$, in perfect agreement with the interpretation of O as the curvature at the vertex. So the sphere is the spherovoid with no deformation, and T is precisely the parameter that deforms the sphere into the rest of the family.

2.8.2 Parovoids, $G = -1$

Setting $G = -1$ in 2.91 gives the condition

$$(n_i^2 z_i - n_o^2 z_o)^2 = -n_i n_o (n_i z_i - n_o z_o) (n_i z_o - n_o z_i), \quad (2.105)$$

and since $G = -1$ makes $-O^2 G = +O^2$, the sag reads

$$z(\rho) = \frac{(O + T\rho^2)\rho^2}{1 + S\rho^2 + \sqrt{1 + (2S + O^2)\rho^2}}. \quad (2.106)$$

These are the **parovoids**, the ovoidal relatives of the paraboloid, which is recovered when $T = S = 0$ and $K = G = -1$ in 2.101, since then the radical becomes $\sqrt{1}$ and $z = Or^2/2$, the parabola of curvature O .

2.8.3 Hyperovoids, $G < -1$

When

$$(n_i^2 z_i - n_o^2 z_o)^2 < -n_i n_o (n_i z_i - n_o z_o) (n_i z_o - n_o z_i), \quad (2.107)$$

the constant satisfies $G < -1$ and the sag keeps the general form

$$z(\rho) = \frac{(O + T\rho^2)\rho^2}{1 + S\rho^2 + \sqrt{1 + (2S - O^2G)\rho^2}}, \quad (2.108)$$

with $-O^2G > O^2$. These are the **hyperovoids**, the deformed versions of the two sheet hyperboloid we obtained in the section of objects at infinity, and they are the surfaces one uses when the conjugate pair demands an eccentricity larger than one.

2.8.4 Prolate and oblate ellipsovoids

Finally, if the product 2.102 is positive,

$$(n_i z_i - n_o z_o)(n_i z_o - n_o z_i) > 0, \quad (2.109)$$

then $G > 0$ and the surface is a **prolate ellipsovoid**, stretched along the optical axis, while in the intermediate window

$$0 > G > -1, \quad (2.110)$$

the surface is an **oblate ellipsovoid**, flattened along the axis. In both cases the sag is given by 2.108, only the sign and the magnitude of G change.

It is illuminating to line up the whole family with the conic constant of the standard surfaces, because the parallel is exact. A conic with $K > 0$ is an oblate ellipsoid, $K = 0$ is a sphere, $-1 < K < 0$ is a prolate ellipsoid, $K = -1$ is a paraboloid and $K < -1$ is a hyperboloid. The ovoids repeat the same ladder with G in the role of K , with the difference that the ovoid carries in addition the deformation T , which is what allows it to be rigorously stigmatic for a *pair of finite points* instead of only for a point and infinity. That extra parameter is exactly the price of bringing the second conjugate point from infinity to a finite distance.

2.9 Descartes ovoids and aspheric surfaces

We have now two different ways of writing surfaces that are better than the sphere, and it is important to understand how they relate to each other, because they answer two different questions.

The rigorously stigmatic surfaces we have built are perfect, but their perfection is extremely local, it holds for one pair of points and for nothing else, with the single exception of the sphere at its aplanatic points, which we will meet again in a moment. If the object is moved away from that privileged position, the image immediately stops being a point, because each ray leaving the object is refracted towards a slightly different place. In practice the position of the object is arbitrary, so a theory that only knows how to be perfect at one point is not enough, and the study of aberrations becomes mandatory.

The standard engineering answer is to give up rigorous stigmatism and instead deform a conic until the residual distortion is small over a whole range of positions. This is done by adding higher order terms in r to the Schwarzschild equation 2.101, called deformation terms,

$$z(r) = \frac{c_0' r^2}{1 + \sqrt{1 - (1 + K') c_0'^2 r^2}} + \sum_{j=2}^{\infty} A_{2j} r^{2j}, \quad (2.111)$$

where the numbers A_{2j} are the **asphericity coefficients** and $K' = -e^2$ is the Schwarzschild constant of the base conic. The conic surfaces are then the degenerate case of the aspheric surfaces in which every A_{2j} vanishes.

Here lies the essential difference. The aspheric surfaces 2.111 are *open* surfaces, they are a conic plus a polynomial correction that grows without bound, whereas the Descartes ovoids are *closed* quartic surfaces. They are therefore not equivalent, and an aspheric surface with $A_{2j} \neq 0$ is in general not rigorously stigmatic for any pair of points at all. What it is good at is something else, it is good at flattening the residual wavefront error over an extended field, which is why aspheres are used to correct aberrated wavefronts and to design isoplanatic systems. The Descartes ovoids, on the contrary, are particular cases of the superconics, and they buy exactness at one pair of points at the cost of losing it everywhere else.

Both languages are needed. The ovoid tells us what perfect imaging looks like and gives us the reference against which everything else is measured, and the asphere tells us how to distribute the imperfection when perfection is impossible. The rest of this chapter is devoted to the second question, and we shall attack it in the place where it matters most, the spherical diopter, which is by far the most used surface in optics for the simple historical reason that it is the easiest one to grind by hand.

2.10 Approximate stigmatism in the spherical diopter

Rigorous stigmatism and the aplanatism conditions give us the principles that govern distortion free image formation, but their usefulness is limited, because they can only be applied when the object sits at one very specific position with respect to the diopter. In practice the position of the object is arbitrary, and what we want is to describe what an instrument does under those conditions, where distortions appear that are, in principle, the consequence of the stigmatism being only approximate.

Among all the diopters the spherical one is the one with the largest number of applications, partly for historical reasons, since it is the surface that was easiest to obtain by hand or by artisanal methods, and that is precisely what pushed the study of its distortions.

Let us take a pair of conjugate points O and O' such that the optical path $L(O, O')$ does not depend on the point I where the ray crosses the spherical diopter, see figure 2.27, where we write $d = z_o$ and $d' = z_i$ for the object and image distances measured from the vertex. Putting the vertex at the origin, the centre of curvature sits at $z = R$ and the incidence point is

$$I = (R \sin \varphi, 0, R(1 - \cos \varphi)), \quad (2.112)$$




Figure 2.27: Image formation by a spherical diopter $\mathcal{D}$ of radius R , vertex V and centre of curvature C . The object O is at a distance $d = z_o$ from the vertex and the image O' at $d' = z_i$, and the ray reaches the surface at the point I , which is located by the angle φ measured at the centre of curvature. The chord $\rho = \overline{VI}$, the transverse height h and the sagitta η are the three geometrical quantities that will organise the whole calculation.

so that the two segments are

$$l_o^2 = [R(1 - \cos \varphi) - z_o]^2 + R^2 \sin^2 \varphi = R^2 (1 - 2 \cos \varphi + \cos^2 \varphi + \sin^2 \varphi) - 2Rz_o(1 - \cos \varphi) + z_o^2,$$

where the two terms in blue add up to one, and therefore

$$l_o = \left[R^2 + (z_o - R)^2 + 2R(z_o - R) \cos \varphi \right]^{1/2}, \quad (2.113)$$

$$l_i = \left[R^2 + (z_i - R)^2 + 2R(z_i - R) \cos \varphi \right]^{1/2}, \quad (2.114)$$

the second one being obtained in exactly the same way on the image side. Both expressions can be written more compactly as $l_o^2 = z_o^2 + 2R(R - z_o)(1 - \cos \varphi)$ and $l_i^2 = z_i^2 + 2R(R - z_i)(1 - \cos \varphi)$,

and in this form it is evident that the whole dependence on the incidence point enters only through $1 - \cos \varphi$.

The optical path is $L(OO') = n_o l_o + n_i l_i$, and if we demand the Fermat's principle for the ray that passes through I , that is, if we demand that small variations of φ produce no first order change in L , we must impose $dL/d\varphi = 0$. Differentiating 2.113 and 2.114,

$$2l_o \frac{dl_o}{d\varphi} = -2R(z_o - R) \sin \varphi \implies \frac{dl_o}{d\varphi} = \frac{R(R - z_o)}{l_o} \sin \varphi,$$

and identically for l_i , so the stationarity condition reads

$$n_o \frac{R(z_o - R)}{l_o} \sin \varphi + n_i \frac{R(z_i - R)}{l_i} \sin \varphi = 0, \quad (2.115)$$

where we have changed the global sign, which is harmless since the right hand side is zero. Dividing by $R \sin \varphi$, which is legitimate for any ray that is not the axial one, and separating the terms,

$$\begin{aligned} \frac{n_o(z_o - R)}{l_o} + \frac{n_i(z_i - R)}{l_i} &= 0, \\ \frac{n_o z_o}{l_o} - \frac{n_o R}{l_o} + \frac{n_i z_i}{l_i} - \frac{n_i R}{l_i} &= 0, \end{aligned}$$

so that we finally obtain the exact conjugation relation of the spherical diopter,

$$\frac{n_i}{l_i} + \frac{n_o}{l_o} = \frac{1}{R} \left[n_i \frac{z_i}{l_i} + n_o \frac{z_o}{l_o} \right]. \quad (2.116)$$

It is very important to be clear about what 2.116 does and does not say. We have not demanded $L(O, O') = cte$ for every ray, which is what rigorous stigmatism would require, we have only demanded that the Fermat's principle hold for the ray that goes through the particular point I , that is, that for a given angle φ small variations $\Delta\varphi$ produce negligible variations of $L(O, O')$. Therefore O and O' are not necessarily the Young-Weierstrass points. From here on we shall no longer speak of rigorous stigmatism but of approximate stigmatism, aplanatism will be replaced by the weaker notion of isoplanatism, and in general no perfect image is formed at all.

2.10.1 Gauss optics, the paraxial approximation

An enormous simplification is obtained if we restrict the rays that are allowed through the system so that $\sin \varphi \approx \varphi$ and $\cos \varphi \approx 1$. This is the paraxial or Gaussian approximation, and geometrically it means that we keep only the rays that hug the optical axis, so that the diopter is replaced by its tangent plane as far as the distances are concerned. In this regime $l_o \approx -z_o$, the minus sign coming from our convention since $z_o < 0$ while l_o is a positive length, and $l_i \approx z_i$. Substituting into 2.116,

$$\frac{n_i}{z_i} + \frac{n_o}{-z_o} = \frac{1}{R} \left[n_i \frac{z_i}{z_i} + n_o \frac{z_o}{-z_o} \right] = \frac{n_i - n_o}{R},$$

that is,

$$\frac{n_i}{z_i} - \frac{n_o}{z_o} = \frac{n_i - n_o}{R}, \quad (2.117)$$

which is the conjugation formula of the spherical diopter in the paraxial approximation, and we say that O and O' are conjugate points, O' being the paraxial image of O . The right hand side depends only on the surface and the media, never on the position of the object, so it deserves a name of its own, the **vergence**

$$V = \frac{n_i - n_o}{R}. \quad (2.118)$$

The focal distances are defined by sending either the object or the image to infinity. Taking $z_i \rightarrow \infty$ in 2.117 the left hand side keeps only the object term and we obtain the object focus, and taking $z_o \rightarrow \infty$ we obtain the image focus,

$$f_o = \frac{n_o R}{n_o - n_i}, \quad (2.119)$$

$$f_i = \frac{n_i R}{n_i - n_o}. \quad (2.120)$$

Notice that f_o and f_i are not equal unless $n_o = n_i$, and that their ratio is $f_i/f_o = -n_i/n_o$. This asymmetry, which does not exist for a mirror, is the geometrical signature of the fact that the light travels at different speeds on the two sides of the surface.

Under these conditions the optical path is approximately independent of the point I , and this is why we speak of approximate stigmatism. The image is formed with distortions which on one hand are minimal, and on the other hand are invariant under a translation in a transverse plane, and this last property deserves its own name.

Definition 8 (*Isoplanatism*) *An approximately stigmatic system whose distortions remain invariant under a transverse translation of the object is called **isoplanatic**.*

The physical content of isoplanatism is worth spelling out, because it is the bridge between geometrical optics and the Fourier optics we will need much later. If the defects of the instrument are the same at every point of the field, then the instrument acts on the object as a linear and space invariant filter, and the image is the convolution of the object with a single impulse response. If the defects change from point to point the system stops being a convolution and the whole machinery of transfer functions collapses. So isoplanatism is exactly the condition under which an optical instrument can be treated as a linear filter.

By restricting the rays that are allowed through the system, then, we have managed to impose conditions under which the spherical diopter is approximately stigmatic even away from its Young-Weierstrass points. The same approximations can be taken for the hyperboloidal and ellipsoidal diopters, and for the reflective surfaces, with no change in the reasoning.

2.10.2 Spherical aberration

Paraxial optics is a beautiful lie. It is exact only in the limit of a vanishingly thin pencil of rays around the axis, and any real instrument has a finite aperture. So let us now allow the rays that depart moderately from the paraxial condition to also pass through the system, which means we must increase the precision of the approximations and keep the first correction to the paraxial result.

The first step is purely geometrical. From the figure 2.27 we read $\cos \varphi = (R - \eta)/R = 1 - \eta/R$, where η is the sagitta, the axial distance from the vertex to the foot of the incidence point. Since $\rho^2 = \eta^2 + h^2$ and $R^2 = (R - \eta)^2 + h^2$, subtracting the second from the first,

$$\begin{aligned}\rho^2 - R^2 &= \eta^2 - (R - \eta)^2 = \eta^2 - R^2 + 2R\eta - \eta^2 = 2R\eta - R^2, \\ \rho^2 = 2R\eta &\implies \frac{\eta}{R} = \frac{\rho^2}{2R^2},\end{aligned}$$

and therefore

$$\frac{\rho^2}{2R^2} = 1 - \cos \varphi. \quad (2.121)$$

This little relation is the key of the entire section, because it converts the angular variable φ , which is how the geometry is naturally written, into the aperture variable ρ , which is how the aberrations are naturally measured. Substituting 2.121 into the compact form of 2.113,

$$\begin{aligned}l_o^2 &= z_o^2 + 2R^2(1 - \cos \varphi) - 2Rz_o(1 - \cos \varphi) = z_o^2 + 2R^2 \frac{\rho^2}{2R^2} - 2Rz_o \frac{\rho^2}{2R^2}, \\ l_o^2 &= z_o^2 + \rho^2 - \frac{z_o \rho^2}{R} = z_o^2 \left[1 + \frac{\rho^2}{z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right],\end{aligned}$$

so that, remembering $\sqrt{z_o^2} = -z_o$,

$$l_o = -z_o \left[1 + \frac{\rho^2}{z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right]^{1/2}, \quad (2.122)$$

$$l_i = z_i \left[1 + \frac{\rho^2}{z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right]^{1/2}. \quad (2.123)$$

These two expressions are still exact, no approximation has been made yet, we have merely traded φ for ρ . Substituting them into the exact conjugation 2.116 and using $n_o/l_o = -(n_o/z_o)[\dots]^{-1/2}$,

$$\begin{aligned}&\frac{n_i}{z_i} \left[1 + \frac{\rho^2}{z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right]^{-1/2} - \frac{n_o}{z_o} \left[1 + \frac{\rho^2}{z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right]^{-1/2} = \\ &\frac{1}{R} \left\{ n_i \left[1 + \frac{\rho^2}{z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right]^{-1/2} - n_o \left[1 + \frac{\rho^2}{z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right]^{-1/2} \right\}.\end{aligned} \quad (2.124)$$

Now, and only now, we approximate. For $\rho \ll |z_o|$, $\rho \ll z_i$ and $\rho \ll |R|$ the quantity inside the brackets is close to one, and the binomial expansion $(1+x)^{-1/2} \approx 1 - x/2$ gives

$$\frac{1}{l_o} = -\frac{1}{z_o} \left[1 + \frac{\rho^2}{z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right]^{-1/2} \approx -\frac{1}{z_o} \left[1 - \frac{\rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right], \quad (2.125)$$

$$\frac{1}{l_i} = \frac{1}{z_i} \left[1 + \frac{\rho^2}{z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right]^{-1/2} \approx \frac{1}{z_i} \left[1 - \frac{\rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right], \quad (2.126)$$

where the improvement over the Gauss optics is transparent, in the paraxial case we simply put $1/l_o = -1/z_o$ and here we keep the first correction in ρ^2 . Substituting into 2.124,

$$\begin{aligned} & \frac{n_i}{z_i} \left[1 - \frac{\rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right] - \frac{n_o}{z_o} \left[1 - \frac{\rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right] = \\ & \frac{1}{R} \left\{ n_i \left[1 - \frac{\rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \right] - n_o \left[1 - \frac{\rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \right] \right\}, \end{aligned} \quad (2.127)$$

and expanding both sides and moving everything that does not contain ρ^2 to the left,

$$\begin{aligned} \frac{n_i}{z_i} - \frac{n_o}{z_o} &= \frac{n_i - n_o}{R} - \frac{1}{R} \frac{n_i \rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) + \frac{1}{R} \frac{n_o \rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \\ &+ \frac{n_i}{z_i} \frac{\rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) - \frac{n_o}{z_o} \frac{\rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right). \end{aligned} \quad (2.128)$$

The four terms in ρ^2 pair up two by two, the first with the third and the second with the fourth, and each pair produces the same binomial squared,

$$\begin{aligned} \frac{n_i \rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right) \left[\frac{1}{z_i} - \frac{1}{R} \right] &= \frac{n_i \rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right)^2, \\ -\frac{n_o \rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right) \left[\frac{1}{z_o} - \frac{1}{R} \right] &= -\frac{n_o \rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right)^2, \end{aligned}$$

so that the whole calculation collapses into

$$\frac{n_i}{z_i} - \frac{n_o}{z_o} = \frac{n_i - n_o}{R} + \rho^2 \left[\frac{n_i}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 - \frac{n_o}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \right]. \quad (2.129)$$

Equation 2.129 is the conjugation formula of the spherical diopter corrected to first order in the aperture. It is the paraxial formula 2.117 plus one extra term proportional to ρ^2 , and that extra term is the **spherical aberration**. Its meaning is direct, since the right hand side now depends on ρ , the position z_i where the ray crosses the axis depends on how far from the vertex the ray struck the surface. Rays through different zones of the aperture cross the axis at different points, and there is no single image point at all.

The same relation can be rearranged by putting each medium on its own side,

$$n_i \left[\frac{1}{z_i} - \frac{1}{R} - \frac{\rho^2}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 \right] = n_o \left[\frac{1}{z_o} - \frac{1}{R} - \frac{\rho^2}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \right], \quad (2.130)$$

which says that a certain quantity has the same value on both sides of the diopter, and this is what we call a metaxial invariant in the presence of aberrations.

Definition 9 (*Marginal invariant*) For a spherical refracting surface we define

$$\mathcal{M} = n \left[\frac{1}{z} - \frac{1}{R} - \frac{\rho^2}{2z} \left(\frac{1}{z} - \frac{1}{R} \right)^2 \right], \quad (2.131)$$

which takes the primary aberrations into account, and which for $\rho \ll |z|$ and $\rho \ll |R|$ reduces to the Abbe invariant of the paraxial optics

$$\mathcal{A} = n \left(\frac{1}{z} - \frac{1}{R} \right). \quad (2.132)$$

2.10.2.1 The wavefront aberration and the quartic wavefront

So far we have described the aberration as a defect in the ray positions. There is a second and much more powerful description, in terms of the shape of the wavefront that leaves the surface, and it is the one that will connect this chapter with the physical optics of the following ones. Following Hopkins (Hopkins, 1950), we define the **wave front aberration** as the residual phase

$$W(\vec{\rho}) = \frac{1}{2} \left[\frac{n_i}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 - \frac{n_o}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \right] \rho^4 = C_{40} \rho^4, \quad (2.133)$$

where C_{40} is the quartic aberration coefficient for a spherical wavefront passing through a spherical refracting surface. The bracket is exactly one half of the aberration bracket of 2.129, which is the usual statement that the transverse ray aberration is the gradient of the wavefront aberration.

The geometry behind 2.133 is shown in the figure 2.28, and it is worth reading carefully. A *spherical refracting surface converts a spherical wavefront into a quartic wavefront*, and that is the whole origin of the geometric aberrations. The marginal rays converge at a point A' instead of the point A_p where the paraxial rays arrive, because the quartic surface Q that emerges from the diopter moves away from the Gaussian spherical surface of reference A' by the amount $W(\vec{\rho})$. The rays, being normal to the wavefront, no longer meet at a single point but are tangent to an envelope, the **caustic**, which is the bright cusped curve one sees at the bottom of a cup of coffee.

This gives us a very clean way of restating everything we did in the first half of the chapter. When a point source is placed at a strictly stigmatic point of a refracting quartic surface, that is, of a Descartes ovoid, the emerging wavefront is a perfect spherical wave converging to the conjugate




Figure 2.28: Quartic surface Q of the wavefront emerging from a spherical refracting surface. The marginal rays converge at A' instead of the paraxial point A_p , because the quartic wavefront Q departs from the Gaussian spherical reference A' by the amount $\Delta l = W(\vec{\rho})$, and the envelope of the refracted rays forms the caustic.

point. Those points are in general unique for a given quartic surface, and if the object is removed from them the emerging wavefront ceases to be spherical and becomes a quartic phase wavefront. The sphere is a particular case of the Descartes ovoids where the strictly stigmatic pair are the Young-Weierstrass points, and therefore any object placed anywhere else gives rise to a wavefront with quartic geometry and, by the above, to the spherical geometric aberrations.

2.10.2.2 The Seidel aberrations

Up to now the object has been on the axis. If we let it move off axis the aberration function acquires a much richer structure, and this is where the classical five aberrations appear. Let $\vec{A}$ be the vector from the vertex V to the reference point V_0 on the axis, let $\vec{r}$ be the position on the surface measured from V_0 , and let $\vec{\rho} = \vec{r} + \vec{A}$ as before. Calling $a = |\vec{A}|$ and θ the angle between $\vec{A}$ and $\vec{r}$, the cosine

rule gives $\rho^2 = a^2 + r^2 + 2ar \cos \theta$, and the aberration function measured relative to the axial point is

$$\begin{aligned} W &= -C_{40} (\rho^4 - a^4) \\ &= -C_{40} (r^4 + 4ar^3 \cos \theta + 4a^2r^2 \cos^2 \theta + 2a^2r^2 + 4a^3r \cos \theta), \end{aligned} \quad (2.134)$$

where the a^4 generated by squaring the cosine rule has cancelled against the a^4 we subtracted. Since the triangles V_0VC and $A'B'C$ are similar, a is proportional to the transverse image size $\sigma' = \overline{A'B'}$, that is $a = g\sigma'$ with $g = R/(z_i - R)$, and substituting we may write

$$W(r, \theta, \sigma') = {}_0C_{40}r^4 + {}_1C_{31}\sigma'r^3 \cos \theta + {}_2C_{22}\sigma'^2r^2 \cos^2 \theta + {}_2C_{20}\sigma'^2r^2 + {}_3C_{11}\sigma'^3r \cos \theta, \quad (2.135)$$

where the coefficient ${}_aC_{bc}$ carries the powers of σ'^a , r^b and $\cos^c \theta$ respectively. These five terms are the monochromatic or **Seidel aberrations** (Welford, 1986), and each one deserves a word on what it actually looks like and where an instrument builder runs into it,

1. **Spherical aberration**, r^4 . This is the only term with no power of σ' at all, so it survives even for a point on the axis, exactly the effect we derived earlier in this section, rays through different zones of the aperture come to a focus at different points along the axis. It blurs a point into a small disc surrounded by a soft halo, and it is why fast lenses, telescopes and camera objectives with a large aperture need aspheric surfaces or multi-element corrections to stay sharp.
2. **Coma**, $\sigma'r^3 \cos \theta$. One power of the field and three of the aperture means it grows quickly as we move off axis, and it is the aberration that turns a point image into the comet-shaped smear that gives it its name, because rays passing through different annular zones of the aperture land at different transverse positions and with different magnifications. It is the dominant defect of fast, wide field instruments such as Schmidt cameras, and it is precisely what the Abbe sine condition of section 2.11.1 is designed to eliminate.
3. **Astigmatism**, $\sigma'^2r^2 \cos^2 \theta$. Here an off-axis point does not focus to a point at all, but to two separate, mutually perpendicular line images at two different distances along the axis, the tangential and the sagittal foci, because a tilted pencil of rays sees an effectively different curvature of the surface in the two perpendicular planes. It is the same astigmatism that a badly shaped cornea produces in the human eye.
4. **Petzval field curvature**, σ'^2r^2 . This term carries no dependence on θ , so it does not blur the image at all, it simply relocates where the sharpest focus falls. The consequence is that even a system free of every other aberration forms its image on a curved surface, the Petzval surface, rather than on a plane, so a flat detector such as film or a sensor can only be in focus over a small region of the field while the edges turn soft. This is why photographic objectives use dedicated field-flattening elements.
5. **Distortion**, $\sigma'^3r \cos \theta$. Being linear in r , this term displaces the image point transversally without spreading it out at all, the image stays perfectly sharp but the magnification changes

with field position. This is the origin of the familiar barrel and pincushion warping of straight lines seen near the edges of wide angle photographs, an effect with no blur, only geometric displacement.

The structure of 2.135 is very instructive. The first term does not contain σ' at all, which is why spherical aberration survives even for an object on the axis, and it is the only one of the five that does. The last term is linear in r , which means that it displaces the image without blurring it, and this is why distortion changes the shape of a picture but keeps it sharp. The three terms in between mix aperture and field, and they are the ones that make an instrument that is perfect at the centre of the field become useless at its edge.

2.10.2.3 Aberrations close to the paraxial condition

In practice it is convenient to have the aberration term of 2.129 written in terms of only one of the two conjugate distances, and this will be essential when we compose two surfaces into a lens. Since our solution is close to the paraxial condition, we may use the paraxial formula 2.117 inside the aberration term, which is already a small correction, so that the error we commit is of higher order.

The key identity is the following. From 2.117 we have $1/z_i = (1/n_i)[n_o/z_o + (n_i - n_o)/R]$, and therefore

$$\frac{1}{z_i} - \frac{1}{R} = \frac{1}{n_i} \left[\frac{n_o}{z_o} + \frac{n_i - n_o}{R} \right] - \frac{1}{R} = \frac{1}{n_i} \left[\frac{n_o}{z_o} + \frac{n_i - n_o}{R} - \frac{n_i}{R} \right] = \frac{n_o}{n_i} \left(\frac{1}{z_o} - \frac{1}{R} \right),$$

a remarkably clean relation which says that the Abbe invariants on the two sides are proportional. Writing $u = 1/z_o - 1/R$ for brevity, the aberration bracket becomes

$$\frac{n_i}{2z_i} \left(\frac{n_o}{n_i} \right)^2 u^2 - \frac{n_o}{2z_o} u^2 = \frac{u^2}{2} \left[\frac{n_o^2}{n_i z_i} - \frac{n_o}{z_o} \right] = \frac{u^2}{2} \frac{n_o^2}{n_i^2} \left[\frac{n_o}{z_o} + \frac{n_i - n_o}{R} \right] - \frac{u^2}{2} \frac{n_o}{z_o},$$

where in the last step we used the paraxial formula once more to eliminate $1/z_i$. Factoring n_o/n_i^2 ,

$$\begin{aligned} \frac{u^2}{2} \frac{n_o}{n_i^2} \left[\frac{n_o^2}{z_o} + \frac{n_o(n_i - n_o)}{R} - \frac{n_i^2}{z_o} \right] &= \frac{u^2}{2} \frac{n_o}{n_i^2} \left[\frac{n_o^2 - n_i^2}{z_o} + \frac{n_o(n_i - n_o)}{R} \right] = \\ &= \frac{u^2}{2} \frac{n_o}{n_i^2} (n_i - n_o) \left[\frac{n_o}{R} - \frac{n_i + n_o}{z_o} \right], \end{aligned}$$

and rewriting the prefactor as $n_o/n_i^2 = (n_o/n_i)^2/n_o$ we obtain the identity we wanted,

$$\frac{n_i}{2z_i} \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 - \frac{n_o}{2z_o} \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \approx \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R} - \frac{n_i + n_o}{z_o} \right), \quad (2.136)$$

which is the Airy formula for the aberration at refraction by a single surface. Therefore the conjugation formula written in terms of the object distance alone is

$$\frac{n_i}{z_i} - \frac{n_o}{z_o} = \frac{n_i - n_o}{R} + \rho^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R} - \frac{n_i + n_o}{z_o} \right) = V + \Delta V_+, \quad (2.137)$$

and repeating the whole argument with the roles of the two media exchanged, that is, using $1/z_o - 1/R = (n_i/n_o)(1/z_i - 1/R)$, the same expression written in terms of the image distance alone is

$$\frac{n_i}{z_i} - \frac{n_o}{z_o} = \frac{n_i - n_o}{R} - \rho^2 \frac{1}{2} \left(\frac{n_i}{n_o} \right)^2 \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 \left(\frac{n_o - n_i}{n_i} \right) \left(\frac{n_i}{R} - \frac{n_i + n_o}{z_i} \right) = V - \Delta V_-, \quad (2.138)$$

where V is the vergence 2.118 and $\Delta V_{\pm}$ is the change in the vergence produced by the marginal rays.

It is very convenient to absorb these corrections into an effective radius. Defining

$$\frac{1}{R_+} = \frac{\rho^2}{2n_o} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_o} - \frac{1}{R} \right)^2 \left(\frac{n_o}{R} - \frac{n_i + n_o}{z_o} \right), \quad (2.139)$$

$$\frac{1}{R_-} = \frac{\rho^2}{2n_i} \left(\frac{n_i}{n_o} \right)^2 \left(\frac{1}{z_i} - \frac{1}{R} \right)^2 \left(\frac{n_i}{R} - \frac{n_i + n_o}{z_i} \right), \quad (2.140)$$

both 2.137 and 2.138 collapse into the single compact statement

$$\frac{n_i}{z_i} - \frac{n_o}{z_o} = (n_i - n_o) \left(\frac{1}{R} + \frac{1}{R_{\pm}} \right), \quad (2.141)$$

and we define the **marginal rays vergence** as $V_{\pm} = (n_i - n_o)(1/R + 1/R_{\pm})$.

This is the cleanest way of understanding what the marginal rays do. *To include the marginal rays in the study of the spherical refracting surface, all we have to do is to add to the inverse of the radius the term $1/R_{\pm}$, according to whether we are working in object or in image coordinates.* The surface behaves, for those rays, as if it had a different radius of curvature, and since $R_{\pm}$ depends on ρ^2 , each zone of the aperture sees a different sphere. That is spherical aberration in one sentence.

2.10.3 Focal length for marginal rays

Since the aberration term depends on the distance ρ at which the rays cross the surface, different rays are deviated differently and consequently the system does not have a single focal length, but a value that depends on ρ for each ray. Let us compute it.

We take 2.137 in the form $1/z_i = n_o/(n_i z_o) + (n_i - n_o)/(n_i R) + \rho^2 \Delta V_+/n_i$ and send the object to infinity, $z_o \rightarrow \infty$, which is the situation of a telescope pointed at a star. In that limit the first term vanishes, $1/z_o \rightarrow 0$ inside the squared bracket so that it becomes $1/R^2$, and the last parenthesis becomes n_o/R , hence

$$\frac{1}{f_{M+}} = \frac{n_i - n_o}{n_i R} + \frac{\rho^2}{n_i} \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \frac{1}{R^2} \left(\frac{n_i - n_o}{n_o} \right) \frac{n_o}{R} = \frac{n_i - n_o}{n_i R} + \frac{\rho^2 n_o^2 (n_i - n_o)}{2n_i^3 R^3},$$

and factoring the paraxial piece,

$$\frac{1}{f_{M+}} = \frac{n_i - n_o}{n_i R} \left[1 + \frac{\rho^2}{2R^2} \left(\frac{n_o}{n_i} \right)^2 \right], \quad (2.142)$$

or, using 2.121 to go back to the angular variable,

$$\frac{1}{f_+} = \frac{n_i - n_o}{n_i R} \left[1 + \left(\frac{n_o}{n_i} \right)^2 - \left(\frac{n_o}{n_i} \right)^2 \cos \varphi \right]. \quad (2.143)$$

We can now read off the two limits. The focus depends on φ , so we write $f_+(\varphi)$, and the paraxial focus is recovered as $f_{P_+} = \lim_{\varphi \rightarrow 0} f_+(\varphi)$,

$$f_{P_+} = \frac{n_i R}{n_i - n_o}, \quad (2.144)$$

in agreement with 2.120, as it must be. For the marginal rays we adopt the reasonable working condition $\varphi \leq \pi/3$, which corresponds to $\rho \leq |R|$, and taking $f_{M_+} = \lim_{\varphi \rightarrow \pi/3} f_+(\varphi)$, where $\cos \varphi = 1/2$ and therefore $1 - \cos \varphi = 1/2$,

$$\frac{1}{f_{M_+}} = \frac{1}{f_{P_+}} \left[1 + \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \right] = \frac{1}{f_{P_+}} \frac{2n_i^2 + n_o^2}{2n_i^2},$$

that is,

$$f_{M_+} = \frac{2n_i^2}{2n_i^2 + n_o^2} f_{P_+}. \quad (2.145)$$

Since the fraction is smaller than one, the marginal focus always falls *short* of the paraxial focus, which is the well known fact that the outer zones of a positive spherical surface focus too soon. The gap between the two is called the **longitudinal spherical aberration**, $\Delta f_+ = f_{P_+} - f_{M_+}$, and it is given by

$$\Delta f_+ = f_{P_+} \left[1 - \frac{2n_i^2}{2n_i^2 + n_o^2} \right] = \frac{n_o^2}{2n_i^2 + n_o^2} f_{P_+}, \quad (2.146)$$

which is an explicit closed expression for the longitudinal spherical aberration of a spherical refracting surface, see figure 2.29.

It is worth noticing how much physics is packed into 2.146. The aberration is proportional to the paraxial focal length, so a fast surface with a short focal length is not automatically better, and it depends only on the ratio of the two indices, so the only way to reduce it while keeping the surface spherical is to increase n_i with respect to n_o . For a glass of $n_i = 1.5$ in air the ratio $n_o^2/(2n_i^2 + n_o^2)$ is about 0.18, which means that the marginal focus sits almost a fifth of the focal length in front of the paraxial one, a completely unacceptable amount for any serious instrument. This single number explains why the whole history of optical design is the history of fighting spherical aberration.

2.10.3.1 Longitudinal object defocus

The same calculation can be run on the object side. Taking 2.138 and sending the image to infinity, $z_i \rightarrow \infty$, the squared bracket becomes $1/R^2$ and the last parenthesis becomes n_i/R , so that

$$\frac{1}{f_-} = \frac{n_o - n_i}{n_o R} \left[1 + \frac{\rho^2}{2R^2} \left(\frac{n_i}{n_o} \right)^2 \right], \quad (2.147)$$




Figure 2.29: Illustration of the longitudinal spherical aberration. The rays arriving parallel to the axis from an object at infinity are focused at f_{M+} if they are marginal and at f_{P+} if they are paraxial, and the separation between the two is Δf_{+} . The envelope of the refracted rays is the caustic.

or in terms of the angle,

$$\frac{1}{f_{-}} = \frac{n_o - n_i}{n_o R} \left[1 + \left(\frac{n_i}{n_o} \right)^2 - \left(\frac{n_i}{n_o} \right)^2 \cos \varphi \right]. \quad (2.148)$$

Exactly as before, the paraxial object focus is $f_{P-} = \lim_{\varphi \rightarrow 0} f_{-}(\varphi)$ and the marginal one is $f_{M-} = \lim_{\varphi \rightarrow \pi/3} f_{-}(\varphi)$,

$$f_{P-} = \frac{n_o R}{n_o - n_i}, \quad (2.149)$$

$$f_{M-} = \frac{2n_o^2}{2n_o^2 + n_i^2} f_{P-}, \quad (2.150)$$

and we define the **longitudinal object defocus** by $\Delta f_{-} = f_{M-} - f_{P-}$, so that

$$\Delta f_{-} = -\frac{n_i^2}{2n_o^2 + n_i^2} f_{P-}, \quad (2.151)$$

see figure 2.30.

The interest of this second definition is justified when we consider compound optical systems, as in the case of a lens, because there the object space of one subsystem is the image space of the previous one, so that speaking of the aberrations in the object takes on a very concrete meaning.





Figure 2.30: Illustration of the defocus in the object space. The rays that emerge parallel to the axis in the image space come, for the marginal zone, from f_{M-} and, for the paraxial zone, from f_{P-} , and the separation between the two is Δf_{-} .

2.10.3.2 Image formation in the marginal ray regime


Consequently the marginal and the paraxial rays have different focal lengths, so their images are formed at different distances and with different magnifications, and the superposition of the two is precisely what deteriorates the final composite image.

This suggests a way out which is as simple as it is elegant, shown in the figure 2.31. The optical system could form an image of the same quality with the marginal rays as with the paraxial ones if the two sets of rays were *separated* instead of superposed, which is done by putting a pupil that selects only one annular zone of the aperture. For this we take $\Delta\varphi$ as small as necessary so that the Fermat's principle 2.116 is fulfilled separately in each pupil, around some angle φ , and in this way each zone has a well defined focal length, both in the object and in the image space.

The physical lesson is that spherical aberration is not a defect of any individual ray, every ray obeys the Snell-Descartes law perfectly, it is a defect of the *superposition* of rays coming from zones with different focal lengths. Break the superposition and the blur disappears, at the price of throwing away most of the light. This is exactly the trade off that a photographer makes when stopping down a lens, and it is also the reason why the resolution of an instrument cannot be improved indefinitely by closing the aperture, because at some point the diffraction we shall study in a later chapter takes over.



(a) With paraxial rays, selected by a central pupil, the object is imaged through the foci f_{P-} and f_{P+} .



(b) With marginal rays, selected by an annular pupil, the same object is imaged through f_{M-} and f_{M+} .

Figure 2.31: Image formation systems. Each family of rays alone forms a sharp image, and it is only their superposition that produces the aberrated one.

2.10.4 The thin lens with aberrations

A lens is two spherical diopters back to back, so everything we have done can be composed. Let the first surface have radius R_1 and form, from the object at z_o in the medium n_o , an intermediate image at d_1 inside the glass of index n_i , and let the second surface of radius R_2 take that intermediate image, now sitting at d_2 , and form the final image at z_i back in the medium n_o . Using 2.137 for the first surface and 2.138 for the second,

$$\frac{n_i}{d_1} - \frac{n_o}{z_o} = \frac{n_i - n_o}{R_1} + \rho_1^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_o} - \frac{1}{R_1} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R_1} - \frac{n_i + n_o}{z_o} \right), \quad (2.152)$$

$$\frac{n_o}{z_i} - \frac{n_i}{d_2} = \frac{n_o - n_i}{R_2} - \rho_2^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_i} - \frac{1}{R_2} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R_2} - \frac{n_i + n_o}{z_i} \right). \quad (2.153)$$




Figure 2.32: Thin lens with aberrations, formed by two spherical diopters of radii R_1 and R_2 separating the medium n from the glass n' . The thickness Δ between the two vertices is taken to zero at the end of the calculation.

Calling $\Delta = d_1 - d_2$ the thickness of the lens, see figure 2.32, and adding 2.152 and 2.153, the intermediate distances combine as

$$n_i \left(\frac{1}{d_1} - \frac{1}{d_2} \right) = n_i \frac{d_2 - d_1}{d_1 d_2} = -\frac{n_i \Delta}{d_2 (\Delta + d_2)},$$

so that

$$\begin{aligned} \frac{n_o}{z_i} - \frac{n_o}{z_o} &= (n_i - n_o) \left(\frac{1}{R_1} - \frac{1}{R_2} \right) + \frac{n_i \Delta}{d_2 (\Delta + d_2)} \\ &\quad - \rho_2^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_i} - \frac{1}{R_2} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R_2} - \frac{n_i + n_o}{z_i} \right) \\ &\quad + \rho_1^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{z_o} - \frac{1}{R_1} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left(\frac{n_o}{R_1} - \frac{n_i + n_o}{z_o} \right). \end{aligned} \quad (2.154)$$

For a thin lens we take the limit $\Delta \rightarrow 0$, which kills the second term and makes $\rho_1 = \rho_2 = \rho$, and we define the paraxial focal distance by

$$\frac{1}{f_p} = \frac{n_i - n_o}{n_o} \left(\frac{1}{R_1} - \frac{1}{R_2} \right), \quad (2.155)$$

which is the lens maker's formula. With this,

$$\begin{aligned} \frac{n_o}{z_i} - \frac{n_o}{z_o} = \frac{n_o}{f_p} + \rho^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) & \left[\left(\frac{1}{z_o} - \frac{1}{R_1} \right)^2 \left(\frac{n_o}{R_1} - \frac{n_i + n_o}{z_o} \right) \right. \\ & \left. - \left(\frac{1}{z_i} - \frac{1}{R_2} \right)^2 \left(\frac{n_o}{R_2} - \frac{n_i + n_o}{z_i} \right) \right]. \end{aligned} \quad (2.156)$$

For moderate diameters we may replace, on the right hand side only, the paraxial relation $1/z_i = 1/z_o - 1/f_p$, and then send the object to infinity, $z_o \rightarrow \infty$. In that limit $1/z_o \rightarrow 0$, so the first bracket becomes $(1/R_1)^2 (n_o/R_1) = n_o/R_1^3$ and the second becomes $(1/f_p + 1/R_2)^2 (n_o/R_2 + (n_i + n_o)/f_p)$, therefore the focal distance for marginal rays in a thin lens is

$$\frac{1}{f'_m} = \frac{1}{f_p} + \rho^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left[\frac{1}{R_1^3} - \left(\frac{1}{f_p} + \frac{1}{R_2} \right)^2 \left(\frac{1}{R_2} + \frac{n_i + n_o}{n_o} \frac{1}{f_p} \right) \right]. \quad (2.157)$$

Defining the aberration of the thin lens as the whole ρ^2 term,

$$\frac{1}{f_a} = \rho^2 \frac{1}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{n_i - n_o}{n_o} \right) \left[\frac{1}{R_1^3} - \left(\frac{1}{f_p} + \frac{1}{R_2} \right)^2 \left(\frac{1}{R_2} + \frac{n_i + n_o}{n_o} \frac{1}{f_p} \right) \right] = \frac{n_i - n_o}{n_o R_{1+}} - \frac{n_i - n_o}{n_o R_{2-}}, \quad (2.158)$$

where the two effective radii are

$$\frac{1}{R_{1+}} = \frac{\rho^2}{2} \left(\frac{n_o}{n_i} \right)^2 \frac{1}{R_1^3}, \quad (2.159)$$

$$\frac{1}{R_{2-}} = \frac{\rho^2}{2} \left(\frac{n_o}{n_i} \right)^2 \left(\frac{1}{f_p} + \frac{1}{R_2} \right)^2 \left(\frac{1}{R_2} + \frac{n_i + n_o}{n_o} \frac{1}{f_p} \right), \quad (2.160)$$

the whole result takes the memorable form

$$\frac{1}{f'_m} = \frac{1}{f_p} + \frac{1}{f_a}. \quad (2.161)$$

The aberrated lens behaves, for the marginal rays, as a thin lens of focal length f_p in parallel with a second, purely aberrational lens of focal length f_a . And here appears something the single surface could not offer, the bracket in 2.158 contains R_1 and R_2 separately, not only through the combination $1/R_1 - 1/R_2$ that fixes f_p . Two lenses with the same focal length but different *bending*, that is, with the same difference of curvatures but different individual curvatures, therefore have different spherical aberration. This is the origin of the classical technique of lens bending, and it is why a simple magnifier is always ground as a meniscus with its convex face towards the object rather than as a symmetric biconvex lens (Kingslake and Johnson, 2010).

2.11 Aplanatism

Everything up to this point has been about a single pair of points on the axis. Rigorous stigmatism gives us a perfect image of one point, and approximate stigmatism gives us a slightly imperfect image of one point. But no one photographs a point. What we actually want to image is an *extended* object, and so the question that closes this chapter is the following, under what conditions does the good behaviour of a stigmatic pair survive when we move the object off the axis?

Consider a centred optical system $\mathcal{S}$ and an object AB normal to the axis, with image $A'B'$. The system $\mathcal{S}$ is said to be **aplanatic** if for every point M of AB a ray emerging from M and crossing $\mathcal{S}$ passes through M' , the image of M , so that M and M' are a stigmatic pair. The word comes from the Greek *stigma*, point, and the requirement is that the whole segment, not just its centre, be imaged point by point.




Figure 2.33: The aplanatism condition. Two pairs of rigorously stigmatic points (A_1, A'_1) and (A_2, A'_2) are considered, with A'_1 in a neighbourhood of A_1 and A'_2 in a neighbourhood of A_2 , the diopter D separating the media n_1 and n_2 , and $\Delta \vec{I}$ being the displacement of the incidence point on the surface. The Spanish label *vecindad* in the figure means neighbourhood.

Let us suppose two pairs of points (A_1, A'_1) and (A_2, A'_2) that are rigorously stigmatic, see figure 2.33, in such a way that $L(A_1, A_2) = cte$ and $L(A'_1, A'_2) = cte$, where A'_1 is close to A_1 and A'_2 is close to A_2 . In the language of topology, A'_1 lies in a neighbourhood of A_1 and A'_2 lies in a neighbourhood of A_2 .

Using exactly the same variational argument that gave us the vectorial Snell-Descartes law 2.29,

we write the difference of the two optical paths as

$$\Delta L(A_1, A_2) = L(A'_1, A'_2) - L(A_1, A_2) = (n_1 \vec{u}_1 - n_2 \vec{u}_2) \cdot d\vec{I} + n_2 \vec{u}_2 \cdot \Delta \vec{A}_2 - n_1 \vec{u}_1 \cdot \Delta \vec{A}_1 = cte, \quad (2.162)$$

where $\Delta \vec{A}_1 = \vec{A}'_1 - \vec{A}_1$ and $\Delta \vec{A}_2 = \vec{A}'_2 - \vec{A}_2$. The first term is the one that gave us the law of refraction, and it vanishes identically,

$$(n_1 \vec{u}_1 - n_2 \vec{u}_2) \cdot d\vec{I} = 0,$$

because by the Snell-Descartes law 2.29 the vector $n_1 \vec{u}_1 - n_2 \vec{u}_2$ is parallel to the normal $\hat{N}$, while $d\vec{I}$ is tangent to the diopter. We are therefore left with the **aplanatism condition**

$$\Delta L(A_1, A_2) = n_2 \vec{u}_2 \cdot \Delta \vec{A}_2 - n_1 \vec{u}_1 \cdot \Delta \vec{A}_1 = cte. \quad (2.163)$$

This deserves a comment before we specialise it. Equation 2.163 is a statement about *derivatives* of the optical path with respect to the position of the object, not about the optical path itself. Rigorous stigmatism says that L does not change when we move the incidence point I over the surface. Aplanatism says something much stronger, that the first variation of L with respect to a displacement of the object is also independent of I . The first condition makes one point sharp, the second makes the whole neighbourhood of that point sharp.

2.11.1 The Abbe sine condition

Let us now specialise 2.163 to the case that matters for an object normal to the axis, that is, the case where the displacement is *transverse*. Then $\Delta \vec{A}_1$ and $\Delta \vec{A}_2$ are perpendicular to the axis, and the scalar products pick up the sines of the angles that the rays make with the axis,

$$n_2 \overline{A_2 A'_2} \sin \theta_2 - n_1 \overline{A_1 A'_1} \sin \theta_1 = cte. \quad (2.164)$$

The constant is fixed by the cheapest possible ray, the one that travels along the axis. For that ray $\theta_1 = \theta_2 = 0$, both sines vanish, and therefore the constant is zero. Since the constant must be the same for every ray, we obtain

$$\frac{\sin \theta_1}{\sin \theta_2} = \frac{n_2 \overline{A_2 A'_2}}{n_1 \overline{A_1 A'_1}}, \quad (2.165)$$

or, writing the **transverse magnification** as

$$g_t = \frac{\overline{A_2 A'_2}}{\overline{A_1 A'_1}}, \quad (2.166)$$

the condition takes the form in which Abbe stated it in 1873 (Abbe, 1873),

$$n_1 \overline{A_1 A'_1} \sin \theta_1 = n_2 \overline{A_2 A'_2} \sin \theta_2 \quad \Longleftrightarrow \quad g_t = \frac{n_1 \sin \theta_1}{n_2 \sin \theta_2}. \quad (2.167)$$




Figure 2.34: Geometry of the Abbe sine condition. The transverse object $\overline{A_1A_1'}$ in the medium n_1 is imaged as $\overline{A_2A_2'}$ in the medium n_2 , and θ_1 and θ_2 are the angles that a given ray makes with the axis in the object and image spaces.

Definition 10 (*Aplanatism*) *A rigorously stigmatic system whose stigmatism is maintained for extended objects, with a magnification that is invariant under a transverse translation, is called **aplanatic**.*

The Abbe sine condition is one of those results whose statement is far shorter than its consequences, so it is worth unpacking what it actually demands. The quantity g_t on the left of 2.167 is a single number, fixed once the conjugate planes are chosen. The quantity on the right involves θ_1 and θ_2 , which change from ray to ray as we move over the aperture. So the condition says that *the ratio $\sin \theta_1 / \sin \theta_2$ must be the same for every ray in the pencil*, from the one that grazes the axis to the one that grazes the rim. If it is not, then rays coming from different zones of the aperture image the off axis point at different transverse positions, and the point spreads into the comet shaped blur that gives coma its name. The sine condition is therefore, in practice, the condition for the absence of coma.

It is very illuminating to compare it with the paraxial result. In the paraxial regime $\sin \theta \approx \theta$ and 2.167 degenerates into $n_1 y_1 \theta_1 = n_2 y_2 \theta_2$, which is the Lagrange-Helmholtz invariant of Gauss optics. So the sine condition is not a new law, it is the exact version of a relation we already knew in its linearised form, and the whole difficulty of optical design lies in the gap between θ and $\sin \theta$.

There is one more consequence which is worth stating because it closes a circle we opened many pages ago. We proved, when we studied the spherical diopter, that a sphere is rigorously stigmatic for the pair of Young-Weierstrass points

$$z_o = R \left(1 + \frac{n_i}{n_o} \right), \quad z_i = R \left(1 + \frac{n_o}{n_i} \right),$$

and one can check that at those two points the sphere also satisfies the sine condition exactly, which is precisely why they are called the *aplanatic points* of the sphere. This is the only situation in which a spherical surface is simultaneously free of spherical aberration and of coma, and it is not a curiosity of the textbooks, it is the working principle of every high numerical aperture microscope objective. The front element of an oil immersion objective is a hemisphere used exactly at its aplanatic points, and the immersion oil is there to make n_o match the glass so that the geometry is preserved. In this sense the whole of modern microscopy rests on a pair of points that Young and Weierstrass found in a sphere.

Finally, the sine condition also explains why resolution and aperture are the same problem. If we write $\text{NA} = n \sin \theta$ for the numerical aperture, then 2.167 says that the product of the object size by the object side numerical aperture is invariant, $y_1 \text{NA}_1 = y_2 \text{NA}_2$. To resolve smaller details we must shrink y_1 , and the invariant then forces NA_1 to grow. Abbe understood this in the same years, and it led him to the diffraction limit that will occupy us in the chapter on diffraction. Geometrical optics, pushed to its own boundary, ends up pointing at the wave theory that must replace it.

Chapter 3

Electromagnetic Optics and Polarisation

*Light is the slender thread that ties electricity
and magnetism into a single weave.*

“We can scarcely avoid the inference that light consists in the transverse
undulations of the same medium which is the cause of electric and
magnetic phenomena.”

– James Clerk Maxwell

3.1 A historical context of electromagnetic optics and polarisation

The idea that light is an electromagnetic phenomenon - and that its transverse vibration carries a hidden orientation we call polarisation - is one of the great unifications of nineteenth-century physics. Before light could be understood as a travelling disturbance of electric and magnetic fields, it had to shed two older skins: Newton's corpuscles and the purely mechanical aether of the early wave theorists. The story of polarisation, in particular, is the story of how a seemingly minor curiosity - light reflected from a window behaving differently from ordinary light - forced physicists to accept that the optical vibration is transverse, and ultimately vectorial.

The first quantitative encounter with polarisation came in 1808, when Étienne-Louis Malus, looking through a calcite crystal at sunlight reflected from the windows of the Luxembourg Palace in Paris, noticed that the intensity of the transmitted beam changed as he rotated the crystal. From these observations he extracted the law that bears his name, $I = I_0 \cos^2 \theta$, relating the transmitted intensity to the angle between the analyser and the plane of polarisation Malus (1809). Malus had discovered that reflection could endow light with the same directional property that double refraction revealed, without invoking any crystal at all. A few years later, in 1815, David Brewster established the law that the polarisation upon reflection is complete at a particular angle, the Brewster angle, for which the reflected and refracted rays are perpendicular Brewster (1815).

These facts were deeply puzzling within the longitudinal wave picture inherited from sound. The resolution came from Augustin-Jean Fresnel and François Arago, who between 1816 and 1819 showed experimentally that two beams polarised at right angles do not interfere. Fresnel drew the inevitable, audacious conclusion: the optical vibration must be *transverse* to the direction of propagation. From this single hypothesis he derived the laws governing the amplitudes of reflected and refracted light - the Fresnel equations - which remain in use today. Transversality made polarisation a natural attribute of light: a wave oscillating in a plane perpendicular to its motion can do so along infinitely many directions, and the choice of that direction *is* its state of polarisation.

A decisive step toward a quantitative description of partially polarised light was taken by George Gabriel Stokes in 1852. Recognising that realistic beams are never perfectly polarised, Stokes introduced four measurable quantities - today called the Stokes parameters - that completely characterise the state of polarisation, including the degree of polarisation, of any beam, coherent or not Stokes (1852). Decades later, Henri Poincaré gave these parameters a beautiful geometric home: every state of polarisation corresponds to a point on the surface of a sphere, the Poincaré sphere, on which the poles represent circular polarisation and the equator linear polarisation. This geometric picture is the one that reappears, in the scalar reduction of diffraction theory, when one decomposes the field onto two opposite states of the sphere.

The grand synthesis arrived with James Clerk Maxwell. In his 1865 memoir *A Dynamical Theory of the Electromagnetic Field* Maxwell (1865), Maxwell showed that the equations governing electricity and magnetism admit wave solutions propagating at a speed numerically equal to the measured speed of light, and concluded that "light is an electromagnetic disturbance." Polarisation was thereby identified with the direction of the electric field vector $\vec{E}$, and the whole edifice of physical optics became a chapter of electromagnetism. The experimental confirmation came in 1887, when Heinrich

Hertz generated and detected electromagnetic waves in the laboratory, verifying that they reflect, refract and polarise exactly as light does Hertz (1893).

In the twentieth century the description of polarisation was recast in the language of linear algebra. R. Clark Jones, in a celebrated series of papers beginning in 1941, introduced the calculus of two-component complex vectors and 2×2 matrices that now bears his name, ideally suited to fully polarised, coherent light Jones (1941). For partially polarised or depolarising systems, the complementary 4×4 Mueller calculus acting on the Stokes vector provides the appropriate tool. The modern treatment of all these matters - transversality, the Fresnel equations, the Stokes-Poincaré formalism and the Jones and Mueller calculi - is presented with classic completeness in Born and Wolf's *Principles of Optics* Born and Wolf (1999), the reference against which later treatments are still measured.

From Malus's rotating crystal to the polarisation-encoded qubits of quantum information, the orientation of the optical field has proven to be far more than an ornament: it is a genuine degree of freedom of light, vectorial in nature, and the bridge between the scalar optics of rays and the full electromagnetic description developed in the chapters that follow.

3.2 The vector behavior of the electromagnetic radiation

Until this part of the lectures, we've studied the light as a ray, however, a better model for the light becomes with the 4 Maxwell's equations, which are:

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0}, \quad \text{Gauss Law for the electric field} \quad (3.1)$$

$$\nabla \cdot \vec{B} = 0, \quad \text{Gauss Law for the magnetic field} \quad (3.2)$$

$$\nabla \times \vec{E} = -\frac{\partial}{\partial t} \vec{B}, \quad \text{Faraday-Lenz law} \quad (3.3)$$

$$\nabla \times \vec{B} = \mu_0 \epsilon_0 \frac{\partial}{\partial t} \vec{E} + \mu_0 \vec{J}, \quad \text{Ampère-Maxwell Law..} \quad (3.4)$$

The equations from 3.1 to 3.4 have got such elegant form thanks to the vector notation given by the physicist Oliver Heaviside and thanks to some of the most useful theorems of the mathematical analysis, the divergence (Gauss) and curl (Stokes) theorems, which allow a smooth transition from the integral form of the Maxwell equations to the differential ones.

The Gauss Law for the electric field 3.1 is really easy to read when we know the geometric interpretation of the divergence operator $\nabla \cdot$ which quantifies how much the electric field $\vec{E}$ goes into (also gets out) from a point, but the right hand side of the equation 3.1 tells us that the presence of any *charge density* ρ produces sources and sinks of electric field (positive charge densities produce sources and negative charge densities produce sinks of electric field).

The Gauss Law for the magnetic field 3.2 tells us that **always the sources of magnetic fields are compensated by sinks of magnetic fields**, also it is interpreted as **there are no magnetic fields monopoles**.

The equation 3.3 is known as the Faraday-Lenz law, the Faraday law tells us that every circulation of electric field produces a variation of magnetic field and every time variation of magnetic field produces circulations of electric field, however there is something important, because if we take the equation 3.3 as $\nabla \times \vec{E} = \partial_t \vec{B}$ without the minus sign there won't be energy conservation, so the minus sign is introduced by Lenz and this equation can describe the behavior of eddy currents.

The fourth equation 3.4 describes the circulations of magnetic fields, due to the Gauss Law 3.2 there won't be any polar magnetic field, it is only possible to have axial-like magnetic fields so, this circulations can only be created by two ways, from the presence of density currents $\vec{J}$ or by time variation of the electric fields, Maxwell added this term such that the continuity equation

$$\frac{\partial}{\partial t} \rho + \nabla \cdot \vec{J} = 0, \quad (3.5)$$

is satisfied.

Now, because we're interested in the study of the electromagnetic radiation *light* we must look for a wave equation, to do this, we're going to use the following mathematical identity:

$$\nabla \times (\nabla \times \vec{B}) = \nabla (\nabla \cdot \vec{B}) - \nabla^2 \vec{B}. \quad (3.6)$$

From the Faraday-Lenz law 3.3 we've got applying the curl to both sides

$$\nabla (\nabla \times \vec{E}) = -\nabla \times \left(\frac{\partial}{\partial t} \vec{B} \right),$$

now, before we're going further we're going to suppose that the electromagnetic fields are smooth vector functions, or at least of class $\mathcal{C}^2 := \{f : D^2 f \in \mathcal{C}\}$ (this means all the second derivatives are continuous), therefore because these are continuous we can use the identity 3.6 and interchange the curl with the time derivative on $\vec{B}$, thus

$$\nabla (\nabla \cdot \vec{E}) - \nabla^2 \vec{E} = -\frac{\partial}{\partial t} (\nabla \times \vec{B}),$$

to solve the left hand side we are going to use the Gauss law of the electric field 3.1 and for the right hand side we're going to use the Ampère-Maxwell law 3.4 such that we've got

$$\begin{aligned} \nabla \left(\frac{\rho}{\epsilon_0} \right) - \nabla^2 \vec{E} &= -\frac{\partial}{\partial t} \left(\mu_0 \epsilon_0 \frac{\partial}{\partial t} \vec{E} + \mu_0 \vec{J} \right), \\ \nabla^2 \vec{E} - \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{E} &= \frac{1}{\epsilon_0} \nabla \rho + \mu_0 \frac{\partial}{\partial t} \vec{J}, \end{aligned} \quad (3.7)$$

where $C^2 = \frac{1}{\epsilon_0 \mu_0}$.

The 3.7 is the formal wave equation for the electric field. In the left hand side we find the traditional well known wave equation but on the right hand side we can see some physical information about how can those waves be generated. We can see that any gradient of electric charge densities produces electric waves and that time variations of the current density source them too, with a plus sign, because the derivation used the minus sign of the Lenz law twice: once when $\partial_t \vec{B}$ was replaced by $-\nabla \times \vec{E}$ in going from 3.4 to the line above, and a second time when the whole equation was rearranged to bring $\nabla^2 \vec{E} - \frac{1}{C^2} \partial_t^2 \vec{E}$ to the left, and the two minus signs cancel each other. Now, we've shown that there are electric waves so we're going to proof that there are also magnetic waves.

From the Ampère-Maxwell law 3.4 we've got applying the curl on both sides

$$\nabla (\nabla \times \vec{B}) = \nabla \times \left(\mu_0 \epsilon_0 \frac{\partial}{\partial t} \vec{E} + \mu_0 \vec{J} \right),$$

now, under our assumptions and also $\rho, \vec{J}$ are at least $\mathcal{C}^2$, then we can apply the identity 3.6 and interchange the time derivative with the curl on $\vec{E}$, thus

$$\nabla (\nabla \cdot \vec{B}) - \nabla^2 \vec{B} = \mu_0 \epsilon_0 \frac{\partial}{\partial t} (\nabla \times \vec{E}) + \mu_0 \nabla \times \vec{J}.$$

On the left side we can use the Gauss law for the magnetic field 3.2, and on the right hand side we can use the Faraday-Lenz law thus

$$\begin{aligned} -\nabla^2 \vec{B} &= \mu_0 \epsilon_0 \frac{\partial}{\partial t} \left(-\frac{\partial}{\partial t} \vec{B} \right) + \mu_0 \nabla \times \vec{J}, \\ \nabla^2 \vec{B} - \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{B} &= -\mu_0 \nabla \times \vec{J}. \end{aligned} \quad (3.8)$$

The equation 3.8 is the wave equation for the magnetic field, on the left hand side we can recognize the traditional wave terms and on the right hand side we can see the sources of magnetic waves, because there are no magnetic monopoles there is not term for gradients of such charges, only the circulations of density currents produces this magnetic waves.

Before going any further we must stop and think about something important, if the light is both electric and magnetic radiation, why the equations 3.7 and 3.8 are only coupled by the density current, or even more, in the vacuum, both equations are independent, so if we think on photons we can question ourselves, ***Where are the photons? Are the photons on the electric or magnetic field*** We are so used to learn they're on both, but how?

3.3 Maxwell equations are the Schrödinger equation for photons

The sentence *photons are in the electromagnetic radiation* is easy to believe but hard to proof from a classical point of view. We could be tempted by the devil and add both equations 3.7 and 3.8 and get something like

$$\nabla^2 (\vec{E} + \vec{B}) - \frac{1}{C^2} \frac{\partial^2}{\partial t^2} (\vec{E} + \vec{B}) = \frac{1}{\epsilon_0} \nabla \rho + \mu_0 \left(\frac{\partial}{\partial t} - \nabla \times \right) \vec{J}, \quad (3.9)$$

which is correct, completely correct **in a world where does not exist physics and analysis**.

From an analysis point of view, the electric field $\vec{E}$ and the magnetic field $\vec{B}$ have got completely different natures, the electric field is a polar vector function, and the magnetic field is an axial vector function, so both fields live on different vector spaces and the addition $\vec{E} + \vec{B}$ does not have any sense.

To solve this problem, there is an interesting vector field called the *Riemann-Silberstein field* which is a correct mathematical (and physical as we we'll see) superposition of the electric and magnetic fields given by

$$\vec{F} = \vec{E} + iC\vec{B}. \quad (3.10)$$

That imaginary unit i may be seen innocent, however adding this complex topology to $\vec{E}, \vec{B}$ solves the superposition problem. It is called for Riemann because he was the first one who studied (alongside Gauss) the electromagnetism using complex numbers, and for Silberstein because he used the vector field 3.10 to rewrite the Einstein's field equations in a quaternions formulation. This may be seen as an innocent superposition but we're going to see that this will allow us, from a classical point of view, to arrive to the correct *photon wave function* and showing that the Maxwell equations acting on 3.10 (which behaves as the photon field) are also quantum equations.

Lets rewrite the Maxwell equations 3.1-3.4 in terms of the photon field $\vec{F}$. First, just multiply the Gauss law for the magnetic field 3.2 by iC and add to the electric Gauss law 3.1 and we'll have

$$\nabla \cdot (\vec{E} + iC\vec{B}) = \frac{\rho}{\epsilon_0}. \quad (3.11)$$

And, also multiply the Ampère-Maxwell law 3.4 by iC and add it to the Faraday-Lenz law 3.3 we will have

$$\nabla \times (\vec{E} + iC\vec{B}) = \frac{i}{C} \frac{\partial}{\partial t} (\vec{E} + iC\vec{B}) + iC\mu_0 \vec{J}. \quad (3.12)$$

Joining both equations and using the definitions of $\vec{F}$ we'll have that the four Maxwell's equations becomes two

$$\nabla \cdot \vec{F} = \frac{\rho}{\epsilon_0}, \quad \text{Gauss law for the photon field} \quad (3.13)$$

$$\nabla \times \vec{F} = \frac{i}{C} \frac{\partial}{\partial t} \vec{F} + iC\mu_0 \vec{J}, \quad \text{Faraday-Lenz-Ampère-Maxwell laws for the photon field.} \quad (3.14)$$

The physical interpretation of the equations 3.13 and 3.14 becomes clear. The Gauss law 3.13 tells us that the presence of charge densities produces sources and sinks of photon fields (electromagnetic fields), and the second equation 3.14 says that time variations of the photon field produces circulations of photon fields and also, the mere presence of density currents produces also circulations of this photon fields. Now it is becoming clearer the relation between where are the photons and how can it be seen in the electromagnetic field at classical levels.

Now, we're going to proof that the set of equations 3.13 and 3.14 contains two wave equation. One traditional wave equation for $\vec{F}$ and an Schrödinger wave equation for $\vec{F}$, showing this beauty

behavior both classical and quantum of the Maxwells equation. Lets apply the curl of the second law 3.14 to both sides as follows

$$\nabla \times (\nabla \times \vec{F}) = \frac{i}{C} \nabla \times \left(\frac{\partial}{\partial t} \vec{F} \right) + iC\mu_0 \nabla \times \vec{J},$$

applying the identity 3.6 on the left han side and permutting th time derivative and the curl in the right hand side as

$$\nabla (\nabla \cdot \vec{F}) - \nabla^2 \vec{F} = \frac{i}{C} \frac{\partial}{\partial t} (\nabla \times \vec{F}) + iC\mu_0 \nabla \times \vec{J},$$

using the Gauss law 3.13 and the FLAM law 3.14 we'll have

$$\frac{1}{\epsilon_0} \nabla \rho - \nabla^2 \vec{F} = \frac{i}{C} \frac{\partial}{\partial t} \left(\frac{i}{C} \frac{\partial}{\partial t} \vec{F} + iC\mu_0 \vec{J} \right) + iC\mu_0 \nabla \times \vec{J},$$

carrying out the time derivative on the right hand side, and using $\frac{i}{C} \cdot \frac{i}{C} = -\frac{1}{C^2}$ and $\frac{i}{C} \cdot iC\mu_0 = -\mu_0$, we've got

$$\frac{1}{\epsilon_0} \nabla \rho - \nabla^2 \vec{F} = -\frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{F} - \mu_0 \frac{\partial}{\partial t} \vec{J} + iC\mu_0 \nabla \times \vec{J},$$

and moving $\nabla^2 \vec{F}$ and $\frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{F}$ to the same side, that is multiplying the whole equation by -1 and swapping it around, we arrive at the *classical wave equation for the photon field*

$$\nabla^2 \vec{F} - \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{F} = \frac{1}{\epsilon_0} \nabla \rho + \mu_0 \left(\frac{\partial}{\partial t} - iC \nabla \times \right) \vec{J}. \quad (3.15)$$

As a check, notice that $\nabla^2 - \frac{1}{C^2} \frac{\partial^2}{\partial t^2}$ is a linear operator and $\vec{F} = \vec{E} + iC\vec{B}$ is a linear combination, so 3.15 must equal 3.7 plus iC times 3.8, and indeed $\left[\frac{1}{\epsilon_0} \nabla \rho + \mu_0 \frac{\partial}{\partial t} \vec{J} \right] + iC \left[-\mu_0 \nabla \times \vec{J} \right] = \frac{1}{\epsilon_0} \nabla \rho + \mu_0 \left(\frac{\partial}{\partial t} - iC \nabla \times \right) \vec{J}$, exactly the right hand side above.

We can see that looks very similar to the sinister equation 3.9 but that was a really forced one insted of our beauty 3.15 which was calculated from the Maxwell's equations, and we see the origin of this electromagnetic or photon fields, these waves are originated as from charge density gradients, and time variations and circulations of the density current, now the behavior of the photons, from a classical perspective can be seen in the presence of charges and currents.

Now, to find out the quantum wave equation from the classical equation 3.14 we are going to use, the following algebraic identity:

$$\vec{A} \times \vec{B} = -i (\vec{A} \cdot \vec{S}) \vec{B}, \quad \vec{S} = S_x \hat{i} + S_y \hat{j} + S_z \hat{k} \quad (3.16)$$

$$S_x = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{bmatrix}, \quad S_y = \begin{bmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{bmatrix}, \quad S_z = \begin{bmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The matrices S_x, S_y, S_z are called the photon spin 1 matrices, and $\vec{A}, \vec{B}$ are arbitrary vectors. Using the identity 3.16 on 3.14 we'll have

$$\begin{aligned}\nabla \times \vec{F} &= -i \left(\nabla \cdot \vec{S} \right) \vec{F}, \quad \text{using the product rule} \\ \nabla \times \vec{F} &= -i \left(\vec{S} \cdot \nabla \right) \vec{F}, \\ \frac{i}{C} \frac{\partial}{\partial t} \vec{F} &= -i \left(\vec{S} \cdot \nabla \right) \vec{F} - iC\mu_0 \vec{J},\end{aligned}$$

multiplying both sides by $C\hbar$ we'll have

$$i\hbar \frac{\partial}{\partial t} \vec{F} = C \left(\vec{S} \cdot \frac{\hbar}{i} \nabla \right) \vec{F} - i\hbar C^2 \mu_0 \vec{J}, \quad (3.17)$$

which is the Schrödinger-like equation for the photon field $\vec{F}$ with the perturbation of the density currents $\vec{J}$ if we're on the vacuum, we'll have the well known as the **Birula's photon wave function** Białynicki-Birula (1994)

$$i\hbar \frac{\partial}{\partial t} \vec{F} = C \left(\vec{S} \cdot \frac{\hbar}{i} \nabla \right) \vec{F}. \quad (3.18)$$

Therefore the Maxwell's equation by themselves are classical and quantum, contains the wave functions that describes the photon behavior and, as we will see further in this lectures, we can carry on this result to general relativity using a tensor that I named **Riemann-Silberstein tensor** which even condensates the four equations to just one.

3.4 Properties of the electromagnetic waves

From now on we're going to focus on the electric wave equation 3.7 and not the complete electromagnetic wave equation 3.15 for three reasons:

1. For now, we're going to focus on the radiation propagation on the vacuum, so the electric field intensity $\|\vec{E}\|$ is a lot bigger than the magnetic field intensity $\|\vec{B}\|$ with a relation approximately like $\|\vec{E}\| \approx C\|\vec{B}\|$.
2. The interaction between light and matter is really hard to study when we consider the electric polarization and even more when we consider all the magnetic properties and effects. This topics are going to be study in the lectures on quantum optics.
3. This is still a lecture on fundamentals, so we're going to start from the basic notions and getting richer models while we advance.

3.4.1 Principal solutions of the electric wave equation

Due to the last reasons, we're going to focus on the study of the electric wave equation

$$\nabla^2 \vec{E}(\vec{r}, t) = \frac{1}{C^2} \frac{\partial^2 \vec{E}(\vec{r}, t)}{\partial t^2}. \quad (3.19)$$

We're going to solve this equation in a physical way (latter on this lectures we're going to solve it in a more sophisticated way). We know from a physical point of view that $\vec{E}$ should be a propagation function with a constant velocity $\vec{v}$ (because there is no matter so the momentum is conserved), therefore we could suppose the electric wave have got the following expression

$$\vec{E}(\vec{r}, t) = \vec{u}(\vec{k} \cdot \vec{r} \pm \omega t, t), \quad (3.20)$$

where we have got that $\omega^2/k^2 = C^2$. The two most popular solutions in the basics literature are the plane wave equation

$$\vec{E}(\vec{r}, t) = E(\vec{r}, t) e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi)} \hat{e}(\vec{r}, t), \quad (3.21)$$

where $E(\vec{r}, t)$ is the amplitude, $\vec{k}$ is the waver number, a kind of spatial frequency, ω is the angular frequency, ϕ is the proper fase, an $\hat{e}$ is the unit vector that describes the vector behavior of the electric wave, also we will identify $\hat{e}$ as the *polarisation*.

The other one, very popular is the spherical wave

$$\vec{E}(\vec{r}, t) = \frac{E(t)}{||\vec{r}||} e^{i(kr - \omega t + \phi)} \hat{e}. \quad (3.22)$$

Both expressions are usually written down as if they had fallen from the sky, but each of them is a genuine solution of the wave equation 3.19 and can be obtained directly from it. Let us do it carefully, first for the plane wave and then for the spherical one.

In the vacuum the wave equation 3.19 acts on every Cartesian component of $\vec{E}$ in exactly the same way and without mixing them, so for the moment we can forget the vector character, write $\vec{E}(\vec{r}, t) = \psi(\vec{r}, t) \hat{e}$ with $\hat{e}$ a fixed unit vector, and solve the scalar equation

$$\nabla^2 \psi = \frac{1}{C^2} \frac{\partial^2 \psi}{\partial t^2}. \quad (3.23)$$

A natural way to attack it is to separate the space and time dependence, proposing $\psi(\vec{r}, t) = R(\vec{r})T(t)$. Substituting into 3.23 and dividing by RT we get

$$\frac{\nabla^2 R}{R} = \frac{1}{C^2} \frac{\ddot{T}}{T}.$$

The left hand side depends only on the position and the right hand side only on the time, so the only way for them to be equal for every $\vec{r}$ and every t is that both are the same constant. We call that

constant $-k^2$, where the minus sign is chosen so that the time part stays bounded and oscillatory, as a wave must be. The time equation then becomes

$$\ddot{T} + C^2 k^2 T = 0,$$

and defining $\omega = Ck$, which is precisely the relation $\omega^2/k^2 = C^2$ we anticipated, its solution is $T(t) = e^{\mp i\omega t}$. The spatial part becomes the Helmholtz equation

$$\nabla^2 R + k^2 R = 0. \quad (3.24)$$

Now we ask for the simplest possible geometry of the wavefronts. If we want the surfaces of constant phase to be planes, the field must depend on the position only through a single linear combination of the coordinates, that is, through $\vec{k} \cdot \vec{r}$ for some fixed vector $\vec{k}$. This suggests

$$R(\vec{r}) = e^{i\vec{k} \cdot \vec{r}},$$

and since $\nabla^2 e^{i\vec{k} \cdot \vec{r}} = -\left\|\vec{k}\right\|^2 e^{i\vec{k} \cdot \vec{r}}$, the Helmholtz equation 3.24 is satisfied as long as $\left\|\vec{k}\right\| = k$. The vector $\vec{k}$ that was born as a mere separation constant turns out to point along the direction of propagation and to have a length fixed by the frequency. Putting the space and time parts back together, restoring the amplitude E_0 and a constant phase ϕ , and recovering the polarisation vector, we arrive at

$$\vec{E}(\vec{r}, t) = E_0 e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi)} \hat{e},$$

which is exactly the plane wave 3.21. If instead of the constant E_0 and the fixed $\hat{e}$ we allow a slowly varying amplitude $E(\vec{r}, t)$ and polarisation $\hat{e}(\vec{r}, t)$, we recover the modulated waves that we will use later, whose wavefronts are still approximately planes over regions small compared with the modulation.

There is still one condition we have not used, and it is the one that makes the wave truly electromagnetic. In the vacuum the Gauss law 3.1 reads $\nabla \cdot \vec{E} = 0$, and evaluating it on our solution

$$\nabla \cdot (E_0 e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi)} \hat{e}) = i E_0 (\vec{k} \cdot \hat{e}) e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi)} = 0,$$

which can only hold if $\vec{k} \cdot \hat{e} = 0$. The polarisation is forced to be perpendicular to the direction of propagation, and we recover, now as a theorem instead of a hypothesis, the transversality that Fresnel had to postulate a century and a half earlier.

The plane wave describes a field whose wavefronts are infinite parallel planes, an idealisation that is excellent far from the source but says nothing about how the radiation spreads out from a small emitter. For that situation the natural symmetry is not translational but spherical, so we now look for a field that depends on the position only through the distance $r = \left\|\vec{r}\right\|$ to the source. Working again with the scalar amplitude $\psi(r, t)$, we need the Laplacian of a function that depends on r alone, which in spherical coordinates is

$$\nabla^2 \psi = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \psi}{\partial r} \right).$$

This expression hides a very useful identity. If we expand it and compare it with the second derivative of the product $r\psi$, we find

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \psi}{\partial r} \right) = \frac{1}{r} \frac{\partial^2}{\partial r^2} (r\psi), \quad (3.25)$$

which turns the awkward radial Laplacian into an ordinary second derivative acting on $r\psi$. Using 3.25, the wave equation 3.23 for a spherically symmetric field becomes

$$\frac{1}{r} \frac{\partial^2}{\partial r^2} (r\psi) = \frac{1}{C^2} \frac{\partial^2 \psi}{\partial t^2}.$$

Multiplying both sides by r and noticing that r does not depend on the time, so it can go inside the time derivative on the right, we get

$$\frac{\partial^2}{\partial r^2} (r\psi) = \frac{1}{C^2} \frac{\partial^2}{\partial t^2} (r\psi).$$

If we now baptise the combination $u(r, t) = r\psi(r, t)$, this is nothing but the one dimensional wave equation

$$\frac{\partial^2 u}{\partial r^2} = \frac{1}{C^2} \frac{\partial^2 u}{\partial t^2}, \quad (3.26)$$

whose general solution, by d'Alembert, is the sum of a disturbance travelling outwards and one travelling inwards,

$$u(r, t) = f(r - Ct) + g(r + Ct).$$

The term $g(r + Ct)$ represents a wave collapsing towards the source, so for radiation escaping from an emitter we keep only the outgoing part $u(r, t) = f(r - Ct)$. Undoing the change $u = r\psi$ we finally obtain

$$\psi(r, t) = \frac{f(r - Ct)}{r},$$

and the factor $1/r$, which we had to guess before, now appears on its own as a direct consequence of the geometry. Choosing a monochromatic profile $f(r - Ct) = E_0 e^{ik(r-Ct)} = E_0 e^{i(kr-\omega t)}$, again with $\omega = Ck$, and restoring the phase and the polarisation we reach the spherical wave 3.22,

$$\vec{E}(\vec{r}, t) = \frac{E_0}{r} e^{i(kr-\omega t+\phi)} \hat{e}.$$

It is worth pausing on the $1/r$. The amplitude of the field falls off as the inverse of the distance, so its square, which measures the intensity, falls off as $1/r^2$. Since the area of a spherical wavefront grows exactly as r^2 , the total power crossing every sphere centred on the source is the same, and energy is conserved as the wave expands. This is why the field goes as $1/r$ and not as $1/r^2$: the inverse square law belongs to the intensity, not to the field itself.

The plane wave and the spherical wave look like very different objects, and yet they are intimately related, because far enough from the source the spherical wave is indistinguishable from a plane wave. To see it, place ourselves at a large distance r_0 from the emitter along a fixed direction $\hat{n} = \vec{r}_0/r_0$,

and look at the field only inside a small region around that point, writing $\vec{r} = \vec{r}_0 + \vec{\delta}$ with $\|\vec{\delta}\| \ll r_0$. The distance to the source is then

$$r = \|\vec{r}_0 + \vec{\delta}\| = r_0 \sqrt{1 + \frac{2\hat{n} \cdot \vec{\delta}}{r_0} + \frac{\|\vec{\delta}\|^2}{r_0^2}} \approx r_0 + \hat{n} \cdot \vec{\delta},$$

where we kept only the first order in $\vec{\delta}/r_0$. The phase of the spherical wave becomes, up to the constant kr_0 ,

$$kr \approx kr_0 + k\hat{n} \cdot \vec{\delta} = kr_0 + \vec{k} \cdot \vec{\delta}, \quad \vec{k} = k\hat{n},$$

so it depends on the position through the single linear combination $\vec{k} \cdot \vec{\delta}$, which is exactly the signature of a plane wave travelling along $\hat{n}$. The amplitude, on the other hand, is $E_0/r \approx E_0/r_0$, which over the small region hardly changes and can be treated as the constant E_0/r_0 . Putting both together, in the neighbourhood of the point the field reads

$$\vec{E}(\vec{r}, t) \approx \frac{E_0}{r_0} e^{i(kr_0 + \vec{k} \cdot \vec{\delta} - \omega t + \phi)} \hat{e},$$

which is precisely a plane wave 3.21 of amplitude E_0/r_0 and wavevector $\vec{k} = k\hat{n}$. In physical terms, a small patch of a sphere of enormous radius is almost flat, and the observer far from the source sees the curved wavefronts as if they were planes. This is the regime in which the plane wave, despite carrying infinite energy across an infinite front, becomes a legitimate and extremely convenient approximation, and it is the reason why so much of optics can be done with plane waves even though every real source radiates spherically.

3.4.2 Beams with a particular phase and structured light

The plane wave and the spherical wave are the two solutions that every textbook writes down first, and for a good reason, since they are the simplest geometries the wave equation admits. It would be a serious mistake, however, to believe that they are the only solutions, or even the most important ones. Both are idealisations that no laboratory ever produces exactly, because the plane wave has an infinite transverse extent and carries an infinite power, and the pure spherical wave demands a perfectly point like source radiating in every direction at once. The light we actually generate and use, above all the light of a laser, comes in the form of *beams*, that is, fields well confined around a propagation axis that carry a finite power. What distinguishes one beam from another, and what grants each one its remarkable properties, is not so much its amplitude as the way its *phase* is organised across the wavefront. This subsection is a short tour through the most relevant of these structured beams.

Before writing any of them it helps to have a common equation from which they all descend. Almost every beam of interest travels mostly along one axis, which we take to be z , with only a gentle spreading in the transverse plane, and this is what we call the *paraxial regime*. We therefore look for solutions of the Helmholtz equation 3.24 of the form

$$\vec{E}(\vec{r}, t) = u(x, y, z) e^{i(kz - \omega t)} \hat{e},$$

where the exponential carries the fast oscillation along the axis and u is a slowly varying envelope that sculpts the beam. Substituting this into the Helmholtz equation and cancelling the common factor we obtain

$$\nabla_{\perp}^2 u + \frac{\partial^2 u}{\partial z^2} + 2ik \frac{\partial u}{\partial z} = 0, \quad \nabla_{\perp}^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2},$$

and because the envelope varies slowly over a wavelength we may neglect $\partial_z^2 u$ against $2k \partial_z u$, arriving at the *paraxial wave equation*

$$i \frac{\partial u}{\partial z} = -\frac{1}{2k} \nabla_{\perp}^2 u. \quad (3.27)$$

It is impossible not to notice that 3.27 is formally identical to the Schrödinger equation of a free particle in two dimensions, with the propagation coordinate z playing the role of the time and $1/k$ playing the role of $\hbar/m$. This is not a coincidence but another face of the deep link between Maxwell and Schrödinger we met earlier in this chapter, and it tells us that the whole catalogue of structured beams is, in a precise mathematical sense, the catalogue of the wave packets of a free particle.

The simplest and most important solution of 3.27 is the one whose transverse profile is a Gaussian. Writing $\rho^2 = x^2 + y^2$ for the transverse radius, its electric field is

$$\vec{E} = E_0 \frac{w_0}{w(z)} \exp\left(-\frac{\rho^2}{w(z)^2}\right) \exp\left(i \left[kz - \omega t + \frac{k\rho^2}{2R(z)} - \zeta(z) \right]\right) \hat{e}, \quad (3.28)$$

where the three functions that govern the beam are the width, the radius of curvature and the Gouy phase,

$$w(z) = w_0 \sqrt{1 + \left(\frac{z}{z_R}\right)^2}, \quad R(z) = z \left[1 + \left(\frac{z_R}{z}\right)^2 \right], \quad \zeta(z) = \arctan \frac{z}{z_R}, \quad (3.29)$$

and $z_R = \frac{kw_0^2}{2} = \frac{\pi w_0^2}{\lambda}$ is the *Rayleigh range*, the distance over which the beam stays approximately collimated, while w_0 is the *waist*, the minimal radius the beam reaches at $z = 0$. Reading the phase of 3.28 term by term makes the physics appear. The piece kz is the ordinary plane wave advance along the axis. The quadratic piece $k\rho^2/2R(z)$ is exactly the phase that a spherical wave of radius $R(z)$ carries near the axis, so it tells us that the wavefronts are neither planes nor full spheres but gently curved caps, flat at the waist where $R \rightarrow \infty$, most strongly curved around the Rayleigh range, and flattening again far away where $R(z) \approx z$ and the beam finally looks spherical. The last piece, the *Gouy phase* $\zeta(z)$, is an extra amount of phase, close to π in total, that the beam accumulates as it crosses the focus, an anomaly with no counterpart in the plane wave and measurable by interferometry. The amplitude factor $w_0/w(z)$ in front of the Gaussian keeps the power constant while the beam widens, since as $w(z)$ grows the peak drops so that the integrated intensity is conserved. The beam spreads with a divergence half angle $\theta = \lambda/\pi w_0$, and this reciprocal relation between waist and divergence is nothing but the diffraction limit and, once more, an uncertainty relation between the transverse position and the transverse momentum of the field. The Gaussian beam is the fundamental transverse mode of a laser resonator, the famous TEM₀₀, and it is the most confined beam that diffraction permits, which is why it is the default description of any well behaved laser and the starting point for the design of almost every optical system that works with coherent light.

The Gaussian is only the lowest member of a whole family. Since 3.27 is linear, it admits complete sets of higher order modes that share the same Gaussian envelope but decorate it with nodal structure. In Cartesian coordinates these are the *Hermite-Gauss* modes, indexed by two integers and built from Hermite polynomials, which display a rectangular grid of bright lobes. In cylindrical coordinates the natural family is that of the *Laguerre-Gauss* modes, indexed by a radial number p and an azimuthal number ℓ , whose field is

$$\begin{aligned} \vec{E}_{p\ell} = E_0 \frac{w_0}{w(z)} \left(\frac{\rho\sqrt{2}}{w(z)} \right)^{|\ell|} L_p^{|\ell|} \left(\frac{2\rho^2}{w(z)^2} \right) \exp \left(-\frac{\rho^2}{w(z)^2} \right) e^{i\ell\varphi} \\ \times \exp \left(i \left[kz - \omega t + \frac{k\rho^2}{2R(z)} - (2p + |\ell| + 1) \zeta(z) \right] \right) \hat{e}, \end{aligned} \quad (3.30)$$

where $L_p^{|\ell|}$ are the associated Laguerre polynomials and φ is the azimuthal angle. Two features of 3.30 deserve attention. The Gouy phase now grows in proportion to the mode order $2p + |\ell| + 1$, so different modes advance their phase at different rates, a fact that governs how a superposition of them rotates and reshapes as it propagates. Far more striking is the azimuthal factor $e^{i\ell\varphi}$, which winds the phase through ℓ complete turns around the axis and forces the field to vanish exactly on the axis, giving these beams their characteristic doughnut profile with a dark core. The wavefront is then no longer a stack of caps but a set of intertwined helices, an *optical vortex* of topological charge ℓ . This helical phase is not a mere decoration, because a field carrying $e^{i\ell\varphi}$ transports orbital angular momentum, and each of its photons carries exactly $\ell\hbar$ of it, a quantity entirely separate from the spin angular momentum $\pm\hbar$ associated with circular polarisation. For this reason the Laguerre-Gauss beams have become one of the central tools of modern optics. They are used in the *optical tweezers* to trap microscopic particles and set them spinning, in super resolution microscopy where the dark core of a doughnut beam switches off the fluorescence everywhere except at the very centre, and in classical and quantum communications alike, where the unbounded integer ℓ offers a new and in principle limitless alphabet that lets a single photon encode much more than a single bit.

All the beams written so far spread as they travel, because they are paraxial and diffraction is unavoidable for any field of finite width. It is natural to ask whether the exact Helmholtz equation admits a beam that does not spread at all, and it does. If we demand a field whose transverse intensity profile is literally independent of z , separation of variables in cylindrical coordinates leads to the *Bessel beam*

$$\vec{E} = E_0 J_\ell(k_\perp \rho) e^{i\ell\varphi} e^{i(k_z z - \omega t)} \hat{e}, \quad k_\perp^2 + k_z^2 = k^2, \quad (3.31)$$

where J_ℓ is the Bessel function of order ℓ . Its transverse pattern, a bright central spot when $\ell = 0$ or a vortex ring when $\ell \neq 0$, surrounded by concentric rings, is exactly the same at every plane z , so the beam is said to be *non-diffracting*. The picture behind 3.31 is illuminating, since it is the coherent superposition of plane waves whose wavevectors all lie on a cone of fixed half angle $\arctan(k_\perp/k_z)$, and the interference of that cone regenerates the same profile plane after plane. This conical construction also explains the beam's most surprising property, its *self-healing*, because if an obstacle blocks the central spot the outer rays that build the cone continue past it and reconstruct the pattern a short distance downstream. The ideal Bessel beam pays for these gifts with an infinite

power, since its rings extend forever and each one carries a comparable share of energy, so in practice it is produced with an axicon or a ring aperture and remains non-diffracting only over a finite range. Its long, thin and self-repairing focus makes it invaluable for light sheet microscopy, for precise optical alignment and for material processing whenever a large depth of focus is required.

There is a last beam whose behaviour borders on the paradoxical. Looking again at the paraxial equation 3.27, now in a single transverse dimension, one finds a solution written in terms of the Airy function,

$$u(x, z = 0) = \text{Ai}\left(\frac{x}{x_0}\right) e^{ax/x_0}, \quad a > 0, \quad (3.32)$$

where x_0 sets the transverse scale and the exponential aperture e^{ax/x_0} is introduced to make the total energy finite. As it propagates, the *Airy beam* shows two properties that seem to contradict everything the plane wave taught us. It is non-diffracting, keeping its intensity profile over long distances, and it is *self-accelerating*, since its main lobe does not travel in a straight line but follows a parabolic trajectory $x \propto z^2$, so the beam bends sideways even though no force and no index gradient act on it. Nothing is being violated here, because the centre of mass of the whole field does travel straight, and it is only the intensity pattern that curves, in the same way that the crest of a water wave may move differently from the water itself. Like the Bessel beam it is also self-healing, and these curved, resilient trajectories have been used to route light around obstacles, to guide electric discharges along curved plasma channels, to sweep microscopic particles along a chosen path and, once again, in advanced microscopy.

Seen together, this family teaches a lesson that the plane and the spherical waves, taken alone, keep hidden, namely that what organises a beam and endows it with a useful behaviour is the shape of its phase across the wavefront. A phase linear in the transverse coordinate, $e^{i\vec{k}_\perp \cdot \vec{\rho}}$, simply tilts the beam. A phase quadratic in ρ , $e^{ik\rho^2/2R}$, focuses or defocuses it and is exactly what a thin lens imprints. A phase linear in the azimuth, $e^{i\ell\varphi}$, gives it a vortex and orbital angular momentum. A conical phase produces the non-diffracting Bessel beam, and a cubic phase produces the self bending Airy beam. Modern devices such as the spatial light modulators and the metasurfaces let us paint almost any of these phase profiles onto a wavefront at will, and this ability to engineer the phase is the whole discipline known today as *structured light*. The plane wave and the spherical wave remain useful as building blocks and as limiting cases, but from here on the reader should keep in mind that they are only two convenient points in a vastly richer landscape of solutions.

3.4.3 From 3D wavefronts to 2D wavefronts

As we've seen the electromagnetic radiation has got many presentations, however, there is a very strong formulation where the light is studied in $2D$, in fact, the following chapter treats the electromagnetic radiation as a $2D$ field.

From the mathematical point of view there is a whole theory about arbitrary surfaces or complex geometries, here, we're going to use the properties of *smooth manifolds*¹ This is thanks to the nature

¹A smooth manifold if a set which can be completely covered by subsets and there are smooth functions from the manifold to $\mathbb{R}^n$.




Figure 3.1: A source $\mathcal{S}$ produces an arbitrary kind of wave which propagates in the space-time, then, for any given time and position we can take an arbitrary point p and, due to the smooth manifold nature of this wave, there exists a neighborhood N of p where the wavefront is plane.

of the physical solution of the wave equations, so, given any arbitrary wavefront as shown 3.1 we can choose any point a set and observer on a neighborhood N , and because the wavefront behaves as an smooth manifold then this neighborhood is $\mathbb{R}^2$.

What we are doing is thinkg on this surface like the earth, because we're son tiny we see the earth as plane but it is not. The same phenomena is here, we step on and observer across the point p on the wavefront and for us there is a neighborhood N where it behaves exactly as $\mathbb{R}^2$ ad that is going to allow us to start the study of the polarisation.

3.5 Light's polarisation

In the electromagnetic theory the word *polarisation* is used as a tendency of the electric dipoles to orientate in a given microscopic volume. However, this word also is used to describe the **electric field orientation** in its propagation. Those two definitions then seem completely indepent, however, as we are going to show in the microscopic theory of radiation these two are completely related.

The polarisation of light is one of the most usefull properties of light, there are many applications on real life which uses this property, for example, the cancer cells stole the glucose from the healthy ones, so there is a higher sugar concentration in cancer cells than the healthy ones, so if we irradiate

a sample with linearly polarized light and study the transmitted radiation, we're going to see that the light which went across the cancer cells changed their polarization, rotating a given angle (which depends on the concentration of glucose) so we can detect the presence of cancer cells using the polarisation.

3.5.1 Monochromatic and deterministic formulation: Jones spinors and the polarization ellipse

As we've said, we're going to focus on the electric field behavior in a $2D$ formulation, so the electric field is going to be a vector field $\vec{E} \in \mathcal{C}^2(\mathbb{C}^2)$ which means it is a *complex vector function which their second derivatives are continuous*. Its general form is given by

$$\vec{E}(\vec{r}, t; \omega) = \begin{bmatrix} \mathcal{E}_1 e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi_1)} \\ \mathcal{E}_2 e^{i(\vec{k} \cdot \vec{r} - \omega t + \phi_2)} \end{bmatrix}, \quad (3.33)$$

where we're considering only one frequency ω so this describes the electric field of a monochromatic field and as we can see, this vector field is a plane wave.

We can factor the propagation phase $\exp[i(\vec{k} \cdot \vec{r} - \omega t)]$ in both components and we will have the following vector field

$$\vec{e} = \begin{bmatrix} \mathcal{E}_1 e^{i\phi_1} \\ \mathcal{E}_2 e^{i\phi_2} \end{bmatrix}. \quad (3.34)$$

The vector field 3.34 is known as the *Jones vector* but, in fact, their true nature is vectorial, the field 3.34 is a **spinor**² as we're going to see this spinor behavior in this chapter.

The Jones spinor 3.34 is a vector that describes how the electric field oscillates, in general, under our assumptions³ is an ellipse, well known as the polarization ellipse. To get the ellipse from 3.34 we're going to define the following electric field components, given by

$$\frac{E_1}{\mathcal{E}_1} e^{-i\phi_2} = \frac{1}{2} e^{i\Delta\phi}, \quad (3.35)$$

$$\frac{E_2}{\mathcal{E}_2} e^{-i\phi_1} = \frac{1}{2} e^{-i\Delta\phi}, \quad (3.36)$$

$$\Delta\phi = \phi_1 - \phi_2.$$

Now, we're going to calculate the magnitude of the difference between 3.35 and 3.36, and because these are complex numbers, it is given by

$$\left(\frac{E_1}{\mathcal{E}_1} e^{-i\phi_2} - \frac{E_2}{\mathcal{E}_2} e^{-i\phi_1} \right) \left(\frac{E_1}{\mathcal{E}_1} e^{-i\phi_2} - \frac{E_2}{\mathcal{E}_2} e^{-i\phi_1} \right)^* = \frac{1}{4} \left(e^{i\Delta\phi} - e^{-i\Delta\phi} \right) \left(e^{i\Delta\phi} - e^{-i\Delta\phi} \right)^*,$$

²We can see a spinor as a $1/2$ tensor. From a geometric point of view, we say that a vector $\vec{v}$ is, in fact, a spinor if it needs two complete rotations of 360° to arrive at its original position

³deterministic, $2D$, monochromatic, space-time independent, propagation on vacuum

where $*$ represents the complex conjugate. In the right hand side we can see that the difference between those two exponentials are the sine of the phase difference Δ thus

$$\frac{1}{4} \left(e^{i\Delta\phi} - e^{-i\Delta\phi} \right) \left(e^{i\Delta\phi} - e^{-i\Delta\phi} \right)^* = \frac{1}{4} (2i \sin \Delta\phi) (-2i \sin \Delta\phi) = \sin^2 \Delta\phi. \quad (3.37)$$

In the left hand side we've got the following

$$\left(\frac{E_1}{\mathcal{E}_1} e^{-i\phi_2} - \frac{E_2}{\mathcal{E}_2} e^{-i\phi_1} \right) \left(\frac{E_1}{\mathcal{E}_1} e^{-i\phi_2} - \frac{E_2}{\mathcal{E}_2} e^{-i\phi_1} \right)^* = \frac{|E_1|^2}{\mathcal{E}_1^2} - \frac{E_1 E_2^*}{\mathcal{E}_1 \mathcal{E}_2} e^{i\Delta\phi} - \frac{E_2 E_1^*}{\mathcal{E}_2 \mathcal{E}_1} e^{-i\Delta\phi} + \frac{|E_2|^2}{\mathcal{E}_2^2},$$

the crossed terms in blue corresponds to the same quantity but one is the complex conjugate of the other, so if we factor the minus sign we'll have the real part of that complex number, so, the left hand side is given by

$$\frac{|E_1|^2}{\mathcal{E}_1^2} + \frac{|E_2|^2}{\mathcal{E}_2^2} - 2 \frac{\text{Re} \{ E_1 E_2^* \}}{\mathcal{E}_1 \mathcal{E}_2} \cos \Delta\phi. \quad (3.38)$$

Combining the LHS 3.38 and the RHS 3.37 we will have the *polarisation ellipse*

$$\frac{|E_1|^2}{\mathcal{E}_1^2} + \frac{|E_2|^2}{\mathcal{E}_2^2} - 2 \frac{\text{Re} \{ E_1 E_2^* \}}{\mathcal{E}_1 \mathcal{E}_2} \cos \Delta\phi = \sin^2 \Delta\phi. \quad (3.39)$$

Then we've shown the electric field lives on an geometric space on $\mathcal{C}^2$ which is a real rotated ellipse, before going further we should notice something.

The Jones spinor 3.34 is not an unique thing of the electric field, also describes the magnetic field under these assumptions but also **describes any complex field with 2 degrees of freedom**, for example the gravitational waves have got a polarisation ellipse, 2D phonons have polarisation ellipse, and so on.

From a mathematical point of view the origin of the ellipse is obvious, the Jones spinor 3.34 corresponds to two unit complex circles which rotate asinchronically, so the general figure is an ellipse not a circle, but, there is a further a more fundamental question, and it is **why does the nature choose an ellipse as the fundamental description of the orientation of the electric field** The short answer is: *No, thenaturedoesnotchoosethat*, the long answer is going to be shown in the theory of coherence chapter.

3.5.1.1 How to build the polarisation ellipse: Phasors

Given any Jones spinor 3.34 how can we construct the polarisation ellipse? To answer this we're going to use phasors which is a formal word for a polar form of a complex number.

Because we've got a 2D vector we've got two components, two polar complex numbers $\mathcal{E}_m e^{i\phi_m}$ with $m = 1, 2$ so we can describe these two phasors as two circunferences as is shown in 3.2. The




Figure 3.2: The polarisation ellipse can be reconstructed taking all the possible rotations of the two phasors (which corresponds to the components of the Jones spinor).

x component or $1 - st$ component is a circumference with radius $\mathcal{E}_1$ and the initial point on this circumference is located at an angle ϕ_1 respect to the $x - axis$ (shown as the red circumference in the figure 3.2). The second phasor, corresponding to the y component of the jones spinor is another circumference but the radius is $\mathcal{E}_2$ and the initial point is located at an angle of ϕ_2 but respect to the $y - axis$.

To construct the polarisation ellipse, we start with the two initial points and draw two straight lines, for the horizontal phasor (the red one) the lines are horizontal, and for the vertical phasor (the blue one) the lines are vertical. In the intersection of the two rectes we draw a point, and we displace both initial points by the same angle, let's call it $\delta\phi$ until we arrive at the initial point. The set of all intersection points draw the polarisation ellipse, which their semi-axes are going to be $\mathcal{E}_1, \mathcal{E}_2$ and the orientation is going to depend on $\Delta\phi$ if $\Delta\phi = \frac{\pi}{2} + n\pi$ with $n \in \mathbb{Z}$ the ellipse isn't going to be orientated also $\Delta\phi$ conditionates if the electric field is going to rotate clockwise or counterclockwise.

3.5.1.2 How to build the polarisation ellipse: Ellipse angles




Figure 3.3: The general form of the polarisation ellipse can be described with only two angles; the orientation angle α and the ellipticity angle χ .

It is not very comfortable to work with phasor and even we can't see the ellipse immediately with the Jones spinor, so we would like a more visual or efficient way to interpretate the polarisation. First we must notice that the electric field amplitudes only change the size of the ellipse, but the relative phases ϕ_1, ϕ_2 are the ones who determine the form of the ellipse, so we can work out with a Jones spinor as an unit vector, this means $\vec{e} \rightarrow \hat{e}$ such that $\hat{e} \cdot \hat{e} = 1$.

The general polarisation ellipse can be seen in 3.3 which can be fully described using just two angles α and χ .

Definition 11 (*Characteristic polarisation angles*): We define the two characteristic polarisation angles as the orientation angle $-\pi \leq \alpha \leq \pi$ which describes how orientated is the ellipse respect to the horizontal axis, and the ellipticity angle $-\pi/2 \leq \chi \leq \pi/2$ which describes completely the form of the ellipse.

How can we translate the Jones spinor with characteristic angles? To do this, lets remind that without any loss of generality we can say the Jones spinor is unitary ($\hat{e} \cdot \hat{e} = 1$) and lets




Figure 3.4: The polarisation ellipse is completely unrotated due to the relative phases between the components of the electric field.

consider the following phases $\phi_1 = 0$ and $\phi_2 = \pi/2$ and we'll have a polarization ellipse as in shown in 3.4.

As we can see, the polarization vector can be describe using a polar parametrization with the angle χ , so the Jones spinor becomes

$$\hat{e}(\chi) = \begin{bmatrix} \cos \chi \\ i \sin \chi \end{bmatrix}, \quad (3.40)$$

the i term comes from the phase $\phi_2 = \pi/2$ where $\exp(\pi/2) = i$. Now, to construct the full polarization states shown in 3.3 we only have to rotate the Jones spinor 3.40 an angle α , so, the polarisation states is given by

$$\begin{aligned} \hat{e}(\alpha, \chi) &= \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \chi \\ i \sin \chi \end{bmatrix}, \\ \hat{e}(\alpha, \chi) &= \begin{bmatrix} \cos \alpha \cos \chi - i \sin \alpha \sin \chi \\ \sin \alpha \cos \chi + i \cos \alpha \sin \chi \end{bmatrix} \end{aligned}$$

Definition 12 (*Polarisation convention*): From now on in this lectures whenever we're working with polarisation states in terms of the angles (α, χ) instead of using $\hat{e}(\alpha, \chi)$ we're going to use the ket

notation, therefore $\hat{e}(\alpha, \chi) \mapsto |\alpha; \chi\rangle$ where we use the semicolon ; instead of the comma , because the angles are two parameters and does not represent a two level ket ($|\alpha, \chi\rangle$) (however polarisation is a two level system).

Having on count this convention, we've got that the general description of a polarisation state in terms of the characteristic angles is given by

$$|\alpha; \chi\rangle = \begin{bmatrix} \cos \alpha \cos \chi - i \sin \alpha \sin \chi \\ \sin \alpha \cos \chi + i \cos \alpha \sin \chi \end{bmatrix}. \quad (3.41)$$

Because the polarisation is a $2D$ vector in this formulation it can be written as the superposition of, at least, two linearly independent polarisation states. There is a convention, whenever we talk about polarisation we always write it as the superposition of two orthogonal polarisation states, so, how can we build this orthogonal states? We have got two answers, using the orientation angle or the ellipticity angle. Using the orientation angle is the most obvious and immediate way, because we only have to put the two ellipses orthogonal, this means taking the state $|\alpha; \chi\rangle$ and rotate it like $|\alpha + \frac{\pi}{2}; \chi\rangle$.

To think about orthogonal states using ellipticity we only have got to think about that given any two orthogonal polarisation states $|0; \chi\rangle, |0; \chi'\rangle$ the following two states $|\alpha; \chi\rangle$ and $|0; \chi'\rangle$ are also orthogonal because the rotation transformation preserves the relative angle between vectors. Having this on count, let's suppose that $\chi' = \chi + \theta$ and by orthogonality let's calculate the value of the angle θ that guarantees the orthogonality between polarisation states. Let's calculate the dot product

$$\begin{aligned} \langle 0; \chi | 0; \chi + \theta \rangle &= [\cos \chi, -i \sin \chi] \begin{bmatrix} \cos(\chi + \theta) \\ i \sin(\chi + \theta) \end{bmatrix} = 0, \\ \cos \chi \cos(\chi + \theta) + \sin \chi \sin(\chi + \theta) &= 0, \\ \cos(\chi + \theta - \chi) &= 0, \\ \cos \theta = 0 &\longrightarrow \theta = (2n - 1) \frac{\pi}{2} \quad n \in \mathbb{Z}, \end{aligned}$$

where we have used the cosine of the sum of two angles, so for the cosine of θ to vanish it must be an odd multiple of $\pi/2$, without any loss of generality let's take $n = 1$, so $\theta = \pi/2$ and the orthogonal state can be obtained just taking a displacement of $\pi/2$ on the ellipticity, then any polarization state can be written as

$$|\alpha; \chi\rangle = c_1 |\alpha'; \chi'\rangle + c_2 \left| \alpha'; \chi' + \frac{\pi}{2} \right\rangle. \quad (3.42)$$

3.5.1.3 The canonical polarisation states: horizontal, vertical, diagonal and circular

Equation 3.41 already contains the whole catalogue of polarisation states, so it is worth stopping here and finding explicitly the handful of states that we are going to meet again and again in the laboratory: the two linear states aligned with the coordinate axes, the two diagonal linear states, and the two circular states. For every one of them we want three things: the amplitudes and phases

$\mathcal{E}_1, \mathcal{E}_2, \phi_1, \phi_2$ of the original spinor 3.34, normalised so that $\mathcal{E}_1^2 + \mathcal{E}_2^2 = 1$, and the pair of characteristic angles (α, χ) that reproduce that spinor in 3.41.

Since $|\alpha; \chi\rangle$ has got a real first term in its combination and a purely imaginary second term whenever we compare it componentwise with a target spinor $\vec{e} = [e_1, e_2]^T$, it is convenient to split 3.41 into its real and imaginary parts once and for all,

$$\cos \alpha \cos \chi = \operatorname{Re}\{e_1\}, \quad -\sin \alpha \sin \chi = \operatorname{Im}\{e_1\}, \quad \sin \alpha \cos \chi = \operatorname{Re}\{e_2\}, \quad \cos \alpha \sin \chi = \operatorname{Im}\{e_2\}. \quad (3.43)$$

Every state below is obtained by feeding a particular $\vec{e}$ into the system 3.43 and solving for α, χ inside their allowed ranges $-\pi \leq \alpha \leq \pi$ and $-\pi/2 \leq \chi \leq \pi/2$.

Horizontal linear polarisation. The field only oscillates along the x axis, so the second component of the spinor must vanish, $\mathcal{E}_2 = 0, \mathcal{E}_1 = 1$. A common phase multiplying the whole spinor has got no physical meaning, since it is only a redefinition of the origin of time, so we are free to set $\phi_1 = 0$; the phase ϕ_2 plays no role at all because it multiplies a null amplitude. The normalised spinor is

$$\vec{e}_H = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \phi_1 = 0, \quad \mathcal{E}_2 = 0. \quad (3.44)$$

Plugging $e_1 = 1, e_2 = 0$ into 3.43 gives $\cos \alpha \cos \chi = 1$ together with $\sin \alpha \sin \chi = \sin \alpha \cos \chi = \cos \alpha \sin \chi = 0$, and because $\cos \alpha \cos \chi = 1$ is the maximum value that a product of two cosines can reach, it forces $\cos \alpha = \cos \chi = 1$, that is

$$\boxed{\alpha = 0, \quad \chi = 0} \quad \Longrightarrow \quad |0; 0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (3.45)$$

Vertical linear polarisation. By the same argument, now the field only oscillates along y , so $\mathcal{E}_1 = 0, \mathcal{E}_2 = 1$ and we set $\phi_2 = 0$ (again, ϕ_1 is irrelevant),

$$\vec{e}_V = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \phi_2 = 0, \quad \mathcal{E}_1 = 0. \quad (3.46)$$

Feeding $e_1 = 0, e_2 = 1$ into 3.43 gives $\sin \alpha \cos \chi = 1$, which forces $\sin \alpha = \cos \chi = 1$, so

$$\boxed{\alpha = \frac{\pi}{2}, \quad \chi = 0} \quad \Longrightarrow \quad |\pi/2; 0\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \quad (3.47)$$

Diagonal linear polarisation, at $\pi/4$ from the horizontal. Now both components carry the same amplitude, $\mathcal{E}_1 = \mathcal{E}_2 = 1/\sqrt{2}$, and, this being the new ingredient, they also share the same phase, $\phi_1 = \phi_2 = 0$, so that $\Delta\phi = \phi_1 - \phi_2 = 0$ and

$$\vec{e}_D = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \phi_1 = \phi_2 = 0. \quad (3.48)$$

Here 3.43 reads $\cos \alpha \cos \chi = \sin \alpha \cos \chi = 1/\sqrt{2}$ and $\sin \alpha \sin \chi = \cos \alpha \sin \chi = 0$. If $\sin \chi$ were different from zero the last two equations would force $\sin \alpha = \cos \alpha = 0$ at once, which is impossible, so we must have $\chi = 0$; the first two equations then reduce to $\cos \alpha = \sin \alpha = 1/\sqrt{2}$, giving

$$\boxed{\alpha = \frac{\pi}{4}, \quad \chi = 0}. \quad (3.49)$$

Antidiagonal linear polarisation, at $-\pi/4$ from the horizontal. The amplitudes are the same as in the diagonal case, but now the two components oscillate in opposition, $\phi_1 = 0, \phi_2 = \pi$, so that $e^{i\phi_2} = -1$, $\Delta\phi = -\pi$, and

$$\vec{e}_A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \phi_1 = 0, \quad \phi_2 = \pi. \quad (3.50)$$

The same reasoning used above forces again $\chi = 0$, and now $\cos \alpha = 1/\sqrt{2}, \sin \alpha = -1/\sqrt{2}$, so

$$\boxed{\alpha = -\frac{\pi}{4}, \quad \chi = 0}. \quad (3.51)$$

It is worth pausing on the four linear states 3.45, 3.47, 3.49 and 3.51: all of them share $\chi = 0$, and this is not a coincidence, since the ellipticity angle measures how far the ellipse has departed from a straight segment, and every linear state is nothing but a degenerate ellipse of zero ellipticity. The orientation angle α is the only one that changes from state to state, and it does exactly what its name promises, it tells us the tilt of that segment with respect to the x axis.

Circular polarisations. We now ask for the opposite extreme, the states in which the two orthogonal oscillations have got the same amplitude, $\mathcal{E}_1 = \mathcal{E}_2 = 1/\sqrt{2}$, but are shifted by a quarter of a cycle rather than being in phase or in opposition, $\phi_1 = 0, \phi_2 = \mp\pi/2$, so that $e^{i\phi_2} = \pm i$, $\Delta\phi = \pm\pi/2$, and

$$\vec{e}_{\pm} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ \pm i \end{bmatrix}, \quad \phi_1 = 0, \quad \phi_2 = \mp\frac{\pi}{2}. \quad (3.52)$$

Plugging $e_1 = 1/\sqrt{2}, e_2 = \pm i/\sqrt{2}$ into 3.43, the real part of the second line, $\sin \alpha \cos \chi = 0$, together with the imaginary part of the first line, $\sin \alpha \sin \chi = 0$, can only be satisfied inside the allowed ranges by $\alpha = 0$ (the case $\cos \chi = 0$ is discarded because it would force the left hand side of $\cos \alpha \cos \chi = 1/\sqrt{2}$ to vanish). With $\alpha = 0$ the remaining equations collapse to $\cos \chi = 1/\sqrt{2}$ and $\sin \chi = \pm 1/\sqrt{2}$, so

$$\boxed{\alpha = 0, \quad \chi = \pm\frac{\pi}{4}} \implies |0; \pm\pi/4\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ \pm i \end{bmatrix}. \quad (3.53)$$

Notice that the orientation angle α is completely undetermined by the shape of the state, and indeed it must be so, a circle has got no preferred direction to be tilted towards, so setting $\alpha = 0$ is only a matter of convenience. What really distinguishes the two circular states is the sign of χ , and this sign is directly inherited from the sign of $\Delta\phi = \phi_1 - \phi_2$, exactly as anticipated when we built the

ellipse from the two phasors: a positive $\Delta\phi$ produces $\chi < 0$ and a negative $\Delta\phi$ produces $\chi > 0$. To see the rotation itself, we restore the time dependence at a fixed point in space and take the real part of the spinor 3.52,

$$E_1(t) = \frac{1}{\sqrt{2}} \cos(\omega t), \quad E_2(t) = \frac{1}{\sqrt{2}} \cos\left(\omega t \mp \frac{\pi}{2}\right) = \pm \frac{1}{\sqrt{2}} \sin(\omega t),$$

so that at $\omega t = 0$ the vector (E_1, E_2) points along $+x$, and an instant later it has acquired a small positive component along $+y$ for the upper sign and along $-y$ for the lower sign. In other words, the state with $\chi = +\pi/4$ traces a counterclockwise circle and the state with $\chi = -\pi/4$ traces a clockwise circle in the x - y plane as time goes on. Whether counterclockwise is called left or right circular polarisation depends on whether the observer is defined to look along the propagation direction or against it, a convention we postpone fixing until the Poincaré sphere is introduced; for the moment we simply keep the two states labelled by the unambiguous quantity $\chi = \pm\pi/4$, the largest ellipticity angle allowed, exactly as $\chi = 0$ was the smallest.

Together, the six states 3.45, 3.47, 3.49, 3.51 and 3.53 already reveal the whole geometric role of the pair (α, χ) : α only rotates a state that has got some elongation, while χ interpolates continuously between a straight segment at $\chi = 0$ and a circle at $|\chi| = \pi/4$, with its sign fixing the sense in which the tip of the electric field turns; every other value of χ in between will describe, as we should expect by now, a genuine ellipse of intermediate eccentricity.

3.5.2 Time evolution of the polarisations: Stokes parameters and the Poincaré sphere

The mathematics developed so far describes a beam of light whose polarisation is perfectly defined at every instant, the situation formalised by the Jones spinor 3.34. Real light is seldom so cooperative: sunlight, the light of an incandescent lamp, or even laser light after it has bounced once off a rough surface, is generally a superposition of many independent wavetrains whose relative phase drifts randomly and far too fast to be tracked. Nothing in the deterministic polarisation ellipse built in the previous subsection can describe such a beam, and it was precisely this gap that George Gabriel Stokes set out to close in 1852, in a paper read before the Cambridge Philosophical Society and modestly titled *On the composition and resolution of streams of polarized light from different sources* (Stokes, 1852). Stokes, who a few years later became Lucasian Professor of Mathematics at Cambridge (the same chair later held by Dirac and, more recently, by Hawking), and whose name today is more readily attached to the Navier-Stokes equations of fluid motion, to Stokes' theorem of vector calculus and to the Stokes shift of fluorescence, proposed four real quantities, built from simple combinations of intensities measured behind a polariser and a quarter wave plate held at a handful of fixed orientations, that could characterise the polarisation of any beam whatsoever, fully polarised, fully unpolarised, or anything in between. Unlike the Jones spinor 3.34, which needs a single coherent phase to be written down at all, Stokes' four numbers are honest time averages, and they remain perfectly well defined even when no such phase exists.

It is a curious fact that this construction, so evidently useful and so close in spirit to the density matrix that quantum mechanics would introduce some seventy years later for exactly the same reason

(to describe mixtures that a single wavefunction cannot capture), attracted almost no attention for the rest of the nineteenth century. Stokes himself never returned to it: he went on to serve as secretary and later as president of the Royal Society, absorbed by the administrative life such offices demanded, and his scientific attention drifted to fluorescence, to the wave theory of light in the aether, and above all to hydrodynamics. Nor did the wider community pick it up. Optics in the second half of the nineteenth century was still occupied with problems that were, by construction, coherent and deterministic, interference, diffraction, the propagation of well defined wavefronts, and the statistical treatment of an ensemble of independent wavetrains had no natural home in that programme. Even the form in which Stokes left his result, four numbers without an accompanying algebra of matrices to act on them, made it look more like a table of measurements than like a theory. His four coefficients, not yet known by the name *Stokes parameters* that later authors would give them, waited in the pages of the *Transactions of the Cambridge Philosophical Society* for the better part of a century.

The first person to give this problem a genuinely new geometric life, although without building explicitly on Stokes' algebra, was Henri Poincaré. Poincaré (1854 to 1912) is often remembered as the last of the great universalist mathematicians, equally at home in celestial mechanics, where his study of the three body problem contained, without his fully realising it, the first seeds of chaos theory, in topology, where the conjecture that bears his name would only be settled a century later, and in the foundations of mathematical physics, where his work on the Lorentz group anticipated several of the kinematic ingredients of special relativity. In 1892, in the second volume of his lecture course *Théorie Mathématique de la Lumière*, Poincaré proposed representing every state of polarisation as a single point on the surface of a sphere (Poincaré, 1892). He built this sphere directly from the orientation and ellipticity angles of the polarisation ellipse, the very angles α and χ of the ket 3.41, so that the poles of the sphere fall on the two circular states, its equator is entirely covered by the linear states, exactly as anticipated in the historical overview at the beginning of this chapter, and every other point corresponds to one particular ellipse. It was only later, once the sphere and Stokes' four numbers were laid side by side, that it became clear the coordinates Poincaré had been plotting are, up to normalisation by the total intensity, precisely the three last Stokes parameters: two men, working four decades apart with entirely different tools, one through operational measurements of intensity and the other through the geometry of spherical trigonometry, had arrived at the same object from opposite directions. The sphere has carried Poincaré's name ever since, even though, as we shall see, its natural coordinates are Stokes'.

The parameters finally found the audience they deserved when a new generation of physicists needed to follow partially polarised light through processes that destroy coherence outright, above all multiple scattering in planetary atmospheres, in colloidal suspensions, and in the interstellar medium, a regime where the Jones calculus (Jones, 1941), powerful as it is, simply cannot be used, since it presupposes a single coherent amplitude that scattering does not leave behind. Soleillet, in 1929, and Francis Perrin, in a 1942 study of light scattered by opalescent media (Perrin, 1942), independently rebuilt essentially the same four component, matrix based description that the problem demanded, at first without fully recognising it as Stokes' own. The decisive act of recognition, and the paper usually credited with rescuing Stokes' parameters from a century of obscurity, restoring his original notation and giving the whole subject the name by which it is still taught, was published by Walter McMaster

in the *American Journal of Physics* in 1954 (McMaster, 1954). Subrahmanyan Chandrasekhar's monumental treatise on radiative transfer, appearing in the same decade, adopted the Stokes vector as the natural object to propagate through a scattering atmosphere and cemented its place in astrophysics (Chandrasekhar, 1950). From ellipsometry and remote sensing to the polarisation maps of the cosmic microwave background and to modern quantum optics, where the Stokes parameters reappear as the expectation values of the photon's Pauli operators, the formalism that Stokes wrote down in 1852, that the nineteenth century forgot, and that Poincaré unknowingly rediscovered forty years later in the language of spheres, is today one of the most used tools in the whole of optics.

Let us now build these four parameters explicitly, starting from nothing more than the polarisation ellipse 3.39 itself, and see why Stokes was forced to define them as time averages rather than simply reading them off the field.

The ellipse 3.39 is a fixed curve, but the physical field is not a curve, it is a point running around that curve once every optical period. To say this with equations, the real components E_1, E_2 that satisfy 3.39 at every instant are themselves functions of time,

$$E_1(t) = \mathcal{E}_1 \cos(\omega t - \phi_1), \quad E_2(t) = \mathcal{E}_2 \cos(\omega t - \phi_2), \quad (3.54)$$

the real part of the electric field 3.34 at a fixed point in space (its constant spatial phase $\vec{k} \cdot \vec{r}$ has simply been absorbed into ϕ_1, ϕ_2). Therefore the expression for the polarisation ellipse is given by

$$\frac{|E_1(t)|^2}{\mathcal{E}_1^2} + \frac{|E_2(t)|^2}{\mathcal{E}_2^2} - 2 \frac{\text{Re}\{E_1(t)E_2^*(t)\}}{\mathcal{E}_1\mathcal{E}_2} \cos \Delta\phi(t) = \sin^2 \Delta\phi(t) \quad (3.55)$$

One can check directly that eliminating t between the two equations 3.54, exactly as was done with the phasors 3.35 and 3.36, returns 3.39; the ellipse is nothing but the trace, frozen in space, left behind by the point $(E_1(t), E_2(t))$ as it oscillates.

A single period of visible light lasts a few femtoseconds, $T = \lambda/C \sim 2 \times 10^{-15}$ s for green light, while even the fastest photodetector available in a laboratory needs picoseconds to respond, and the human eye needs tens of milliseconds. In either case an enormous number of optical cycles, somewhere between 10^6 and 10^{13} of them depending on the instrument, go by before any detector can register a single reading; the point $(E_1(t), E_2(t))$ has swept around the ellipse an enormous number of times before any instrument manages to read anything at all. No detector, in other words, can ever measure the instantaneous field $E(t)$ itself, only some average of it taken over a great many periods. This is not a limitation that better technology could one day remove, it is what intensity means: the energy flux carried by a wave is proportional to the square of the field, so what a detector physically integrates is always $\langle E^2 \rangle$, never E . This is exactly the operation Stokes performed, building his four parameters out of quadratic, time averaged combinations of the oscillating field rather than out of the ellipse itself, and it is why they, unlike the Jones spinor, remain meaningful even when no single coherent phase exists to write down. Over one period $T = 2\pi/\omega$, define the time average of any function $f(t)$ as $\langle f \rangle \equiv \frac{1}{T} \int_0^T f(t) dt$. The first two averages we need are the mean square amplitudes,

$$\langle E_1^2 \rangle = \frac{\mathcal{E}_1^2}{T} \int_0^T \cos^2(\omega t - \phi_1) dt = \frac{\mathcal{E}_1^2}{2T} \int_0^T [1 + \cos(2\omega t - 2\phi_1)] dt = \frac{\mathcal{E}_1^2}{2T} [T + 0] = \frac{\mathcal{E}_1^2}{2},$$

where we used $\cos^2 \theta = \frac{1}{2}(1 + \cos 2\theta)$ and the fact that $\cos(2\omega t - 2\phi_1)$ completes exactly two full oscillations as t runs from 0 to T , so its integral over that stretch vanishes; the identical computation gives $\langle E_2^2 \rangle = \mathcal{E}_2^2/2$.

The remaining piece of information lives in how E_1 and E_2 oscillate together, and for that we need the cross average

$$\begin{aligned} \langle E_1 E_2 \rangle &= \frac{\mathcal{E}_1 \mathcal{E}_2}{T} \int_0^T \cos(\omega t - \phi_1) \cos(\omega t - \phi_2) dt \\ &= \frac{\mathcal{E}_1 \mathcal{E}_2}{2T} \int_0^T [\cos(\phi_1 - \phi_2) + \cos(2\omega t - \phi_1 - \phi_2)] dt = \frac{\mathcal{E}_1 \mathcal{E}_2}{2} \cos \Delta\phi, \end{aligned}$$

where we used $\cos A \cos B = \frac{1}{2}[\cos(A - B) + \cos(A + B)]$, the phase difference $\Delta\phi = \phi_1 - \phi_2$ already defined near 3.35, and, once again, that the piece oscillating at 2ω averages to zero over a full period.

This single average, however, only ever gives us $\cos \Delta\phi$, and a beam with $\Delta\phi$ and one with $-\Delta\phi$ (opposite handedness, exactly as when we built the circular states 3.53) would be indistinguishable from $\langle E_1 E_2 \rangle$ alone. To tell them apart we need a second, independent combination, and the natural one to try is E_2 sampled a quarter of a period earlier,

$$\tilde{E}_2(t) \equiv E_2\left(t - \frac{T}{4}\right) = \mathcal{E}_2 \cos\left(\omega t - \frac{\pi}{2} - \phi_2\right) = \mathcal{E}_2 \sin(\omega t - \phi_2),$$

using $\cos(x - \pi/2) = \sin x$ and $\omega T/4 = \pi/2$; physically, this quarter period delay is exactly what a birefringent plate cut to a quarter of a wavelength does to whichever component crosses it, so $\tilde{E}_2$ is not a mere mathematical trick but a genuinely measurable field. Its cross average with E_1 is

$$\begin{aligned} \langle E_1 \tilde{E}_2 \rangle &= \frac{\mathcal{E}_1 \mathcal{E}_2}{T} \int_0^T \cos(\omega t - \phi_1) \sin(\omega t - \phi_2) dt \\ &= \frac{\mathcal{E}_1 \mathcal{E}_2}{2T} \int_0^T [\sin(2\omega t - \phi_1 - \phi_2) + \sin \Delta\phi] dt = \frac{\mathcal{E}_1 \mathcal{E}_2}{2} \sin \Delta\phi, \end{aligned}$$

where now we used $\cos A \sin B = \frac{1}{2}[\sin(A + B) - \sin(A - B)]$ with $A - B = \phi_2 - \phi_1 = -\Delta\phi$, together with the vanishing, as always, of the term oscillating at 2ω .

Having these time averages the time-dependent polarisation ellipse 3.55 becomes

$$\frac{\mathcal{E}_1^2}{2\mathcal{E}_1^2} + \frac{\mathcal{E}_2^2}{2\mathcal{E}_2^2} - \frac{2\mathcal{E}_1 \mathcal{E}_2}{2\mathcal{E}_1 \mathcal{E}_2} \cos^2 \Delta\phi = \sin^2 \Delta\phi,$$

multiplying both side by $4\mathcal{E}_1^2 \mathcal{E}_2^2$ we'll have

$$2\mathcal{E}_1^2 \mathcal{E}_2^2 + 2\mathcal{E}_1^2 \mathcal{E}_2^2 - (2\mathcal{E}_1 \mathcal{E}_2 \cos \Delta\phi)^2 = (2\mathcal{E}_1 \mathcal{E}_2 \sin \Delta\phi)^2,$$

by mathematical convenience we're going to add zero as $0 = \mathcal{E}_1^4 + \mathcal{E}_2^4 - \mathcal{E}_1^4 - \mathcal{E}_2^4$ as follows

$$\begin{aligned} \mathcal{E}_1^4 + \mathcal{E}_2^4 + 2\mathcal{E}_1^2\mathcal{E}_2^2 - \mathcal{E}_1^4 - \mathcal{E}_2^4 + 2\mathcal{E}_1^2\mathcal{E}_2^2 - (2\mathcal{E}_1\mathcal{E}_2 \cos \Delta\phi)^2 &= (2\mathcal{E}_1\mathcal{E}_2 \sin \Delta\phi)^2, \\ (\mathcal{E}_1^2 + \mathcal{E}_2^2)^2 - (\mathcal{E}_1^2 - \mathcal{E}_2^2)^2 - (2\mathcal{E}_1\mathcal{E}_2 \cos \Delta\phi)^2 &= (2\mathcal{E}_1\mathcal{E}_2 \sin \Delta\phi)^2 \end{aligned}$$

We finally have everything we need. Stokes defined his four parameters, up to the overall factor of two needed to cancel the one half that every average above carries, as

$$S_0 \equiv 2\langle E_1^2 \rangle + 2\langle E_2^2 \rangle = \mathcal{E}_1^2 + \mathcal{E}_2^2, \quad (3.56)$$

$$S_1 \equiv 2\langle E_1^2 \rangle - 2\langle E_2^2 \rangle = \mathcal{E}_1^2 - \mathcal{E}_2^2, \quad (3.57)$$

$$S_2 \equiv 4\langle E_1 E_2 \rangle = 2\mathcal{E}_1\mathcal{E}_2 \cos \Delta\phi, \quad (3.58)$$

$$S_3 \equiv 4\langle E_1 \tilde{E}_2 \rangle = 2\mathcal{E}_1\mathcal{E}_2 \sin \Delta\phi. \quad (3.59)$$

Four numbers built out of only two amplitudes and one phase difference cannot be independent, and indeed squaring and adding 3.57 to 3.59 gives

$$\begin{aligned} S_1^2 + S_2^2 + S_3^2 &= (\mathcal{E}_1^2 - \mathcal{E}_2^2)^2 + 4\mathcal{E}_1^2\mathcal{E}_2^2 \cos^2 \Delta\phi + 4\mathcal{E}_1^2\mathcal{E}_2^2 \sin^2 \Delta\phi \\ &= \mathcal{E}_1^4 - 2\mathcal{E}_1^2\mathcal{E}_2^2 + \mathcal{E}_2^4 + 4\mathcal{E}_1^2\mathcal{E}_2^2 (\cos^2 \Delta\phi + \sin^2 \Delta\phi) \\ &= \mathcal{E}_1^4 + 2\mathcal{E}_1^2\mathcal{E}_2^2 + \mathcal{E}_2^4 = (\mathcal{E}_1^2 + \mathcal{E}_2^2)^2, \end{aligned}$$

where the whole phase difference $\Delta\phi$ has disappeared through the single identity $\cos^2 \Delta\phi + \sin^2 \Delta\phi = 1$, leaving, by 3.56,

$$S_1^2 + S_2^2 + S_3^2 = S_0^2. \quad (3.60)$$

This is the equality Stokes was really after: of the four numbers describing a fully polarised beam, only three are independent, exactly as only three are needed to fix a point on a sphere of given radius, a hint of the Poincaré sphere we will make precise shortly.

Every one of these four combinations is quadratic in the field, and this is precisely what lets us rewrite them, exactly, in terms of the complex components $e_1 \equiv \mathcal{E}_1 e^{i\phi_1}$ and $e_2 \equiv \mathcal{E}_2 e^{i\phi_2}$ of the spinor 3.34, without a single cosine or sine left in sight. Since $e_1 e_1^* = \mathcal{E}_1^2$, $e_2 e_2^* = \mathcal{E}_2^2$, and $e_1 e_2^* = \mathcal{E}_1 \mathcal{E}_2 e^{i(\phi_1 - \phi_2)} = \mathcal{E}_1 \mathcal{E}_2 (\cos \Delta\phi + i \sin \Delta\phi)$, comparing real and imaginary parts term by term with 3.58 and 3.59 shows that

$$\mathcal{E}_1^2 \pm \mathcal{E}_2^2 = e_1 e_1^* \pm e_2 e_2^*, \quad 2\mathcal{E}_1 \mathcal{E}_2 \cos \Delta\phi = e_1 e_2^* + e_2 e_1^*, \quad 2\mathcal{E}_1 \mathcal{E}_2 \sin \Delta\phi = i(e_1^* e_2 - e_1 e_2^*),$$

the last equality following from $e_1^* e_2 - e_1 e_2^* = \mathcal{E}_1 \mathcal{E}_2 (e^{-i\Delta\phi} - e^{i\Delta\phi}) = -2i \mathcal{E}_1 \mathcal{E}_2 \sin \Delta\phi$, so multiplying through by i removes the leftover $i^2 = -1$ and leaves a real number, exactly as a physical intensity must be.

Definition 13 (*Stokes parameters*): Given the Jones spinor $\vec{e} = [e_1, e_2]^T$ of a fully polarised,

monochromatic beam 3.34, its Stokes parameters are the four real numbers

$$S_0 = e_1 e_1^* + e_2 e_2^*, \quad (3.61)$$

$$S_1 = e_1 e_1^* - e_2 e_2^*, \quad (3.62)$$

$$S_2 = e_1 e_2^* + e_2 e_1^*, \quad (3.63)$$

$$S_3 = i(e_1^* e_2 - e_1 e_2^*). \quad (3.64)$$

Each of the four carries its own, immediate physical meaning. $S_0 = \mathcal{E}_1^2 + \mathcal{E}_2^2$ is simply the total intensity of the beam, the same quantity we normalised to one when we wrote the Jones ket 3.41 as a unit vector. $S_1 = \mathcal{E}_1^2 - \mathcal{E}_2^2$ measures how much the beam prefers the horizontal axis over the vertical one, positive for a beam closer to $|0; 0\rangle$ and negative for one closer to $|\pi/2; 0\rangle$. $S_2 = 2\mathcal{E}_1\mathcal{E}_2 \cos \Delta\phi$ measures the same kind of preference, only rotated by forty five degrees, between the two diagonal states 3.49 and 3.51. And $S_3 = 2\mathcal{E}_1\mathcal{E}_2 \sin \Delta\phi$ measures the preference between the two senses of circulation 3.53.

Together, (S_1, S_2, S_3) are nothing but three Cartesian coordinates. Writing the components of the general ket 3.41 as $e_1 = \cos \alpha \cos \chi - i \sin \alpha \sin \chi$ and $e_2 = \sin \alpha \cos \chi + i \cos \alpha \sin \chi$, for which $S_0 = 1$ since 3.41 is already a unit vector, and substituting into 3.62 to 3.64, a short calculation using $\cos^2 \theta - \sin^2 \theta = \cos 2\theta$ and $2 \sin \theta \cos \theta = \sin 2\theta$ on each term gives

$$S_1 = S_0 \cos 2\alpha \cos 2\chi, \quad S_2 = S_0 \sin 2\alpha \cos 2\chi, \quad S_3 = -S_0 \sin 2\chi, \quad (3.65)$$

so that $(S_1, S_2, S_3)/S_0$ is, quite literally, a point on a sphere of unit radius, parametrised by twice the orientation angle and twice the ellipticity angle: this is exactly the Poincaré (see the figure 3.5) sphere of the historical discussion above, and 3.65 is the dictionary between its Cartesian coordinates and the angles α, χ used throughout this chapter.

As a consistency check, squaring and adding the three equalities 3.65 reproduces the identity 3.60 once more, $S_1^2 + S_2^2 + S_3^2 = S_0^2 \cos^2 2\chi (\cos^2 2\alpha + \sin^2 2\alpha) + S_0^2 \sin^2 2\chi = S_0^2 (\cos^2 2\chi + \sin^2 2\chi) = S_0^2$, this time because every fully polarised beam corresponds to a single, definite pair of angles α, χ , that is, to a single point sitting exactly on the surface of the Poincaré sphere.

Real light, as we already remarked in the historical discussion, is rarely so cooperative: sunlight, or light that has scattered even once, is better described as a rapidly and randomly fluctuating superposition of many such states, with $\alpha(t)$ and $\chi(t)$ themselves wandering on a timescale far shorter than any detector can follow, so that what actually gets measured is not the Stokes vector of one ellipse but a further time average over many different ones. Averaging a family of unit vectors that point in different directions on the sphere never gives a vector as long as the individual ones were, so for such partially polarised light the equality 3.60 is generally replaced by the inequality $S_1^2 + S_2^2 + S_3^2 < S_0^2$, with the extreme case $S_1 = S_2 = S_3 = 0$ describing completely unpolarised light, a beam with a perfectly well defined intensity but no preferred point on the Poincaré sphere at all. We will not develop this idea further here, since it belongs properly to the theory of coherence taken up in 6, but it is worth keeping in mind already that the four numbers built in this subsection say everything there is to say about polarisation, deterministic or not, while the single ket 3.41 only ever describes the former.




Figure 3.5: In the Poincaré any polarisation state is a point on the surface on the sphere, however if the state is partially polarized it would be a point inside the Poincaré sphere whose distance from the center is $\sqrt{S_1^2 + S_2^2 + S_3^2}$.

In the Poincaré sphere 3.5 and the Stokes parameters 3.65 we see an interesting relation, given any state $|\alpha; \chi\rangle$ on the Poincaré sphere the characteristic angles aren't the same, these two are the double, so a full rotation on the Jones spinor becomes two full rotations of the Stokes vector $\vec{S} = [S_1, S_2, S_3]^T$. In fact, we've proven in 3.42 that two orthogonal polarisation states are separated an angle of $\pi/2$ but, on the Poincaré sphere this angle is the double, so two orthogonal states are separated π from the other, this means **any two orthogonal polarization states are diametrically opposite on the Poincaré sphere**.

3.5.2.1 Why the Stokes vector is not a vector

It is customary, and we shall keep the custom, to arrange the four numbers 3.61 to 3.64 into a column and to call the result the *Stokes vector*. The name is convenient and completely standard, but taken literally it is wrong, and the reason is not a pedantic one, since it is the very same doubling of angles that makes the Poincaré sphere work in the first place.

In physics an object does not earn the name of vector by being a list of numbers. It earns it by transforming in a definite way under the symmetry group of the space it is supposed to live in, so

that a rotation of the laboratory axes by an angle θ takes its components v_i into $R_{ij}(\theta)v_j$. The triple (S_1, S_2, S_3) fails this test at the first attempt. Turn the laboratory axes by θ about the propagation direction; the orientation of every ellipse is now measured from the new x axis, so $\alpha \rightarrow \alpha - \theta$, and 3.65 gives at once

$$S'_1 = S_1 \cos 2\theta + S_2 \sin 2\theta, \quad S'_2 = -S_1 \sin 2\theta + S_2 \cos 2\theta, \quad S'_3 = S_3. \quad (3.66)$$

The pair (S_1, S_2) does rotate, but by twice the angle through which the laboratory was turned. The cleanest way to see how fatal this is, is to turn the laboratory by half a turn, $\theta = \pi$. Every polarisation ellipse is centrally symmetric, so it is carried exactly onto itself, no state of light has changed, and every measurable quantity must come back unchanged; and indeed 3.66 returns $S'_i = S_i$, because $2\theta = 2\pi$. A genuine vector of the transverse plane would have come back as $-\vec{v}$ instead. So (S_1, S_2, S_3) is not a vector of the physical space in which the experiment sits, and there is no direction one could point at in the laboratory and call the direction of S_3 .

The second failure is algebraic rather than geometric. A vector lives in a vector space, and a vector space is closed under multiplication by any real number, in particular by -1 . But $-\vec{S}$ has got $S_0 < 0$, a beam of negative intensity, which is not a state of anything, so no physical Stokes vector possesses an additive inverse. What the physical Stokes vectors form is not a vector space but a convex cone, the region carved out by $S_0 \geq 0$ together with the inequality $S_0^2 \geq S_1^2 + S_2^2 + S_3^2$ discussed above. Even the addition that this cone does admit is not the one a physicist would naively expect: adding two Stokes vectors describes the superposition of two mutually *incoherent* beams, which was precisely Stokes' purpose, and it is emphatically not the coherent superposition of two fields, for which one must add the Jones spinors first and only afterwards form the four averages. The Stokes parameters are quadratic in the field, and quadratic quantities never add the way the field itself does.

If the four numbers are not the components of a vector, then components of what are they? Of an operator. It is enough to collect the very same products that appear in 3.61 to 3.64 into a single array.

Definition 14 (*Polarisation matrix*): The polarisation matrix, also called the coherence matrix, of a beam with Jones spinor $\vec{e} = [e_1, e_2]^T$ is the Hermitian, positive semidefinite matrix

$$\Phi \equiv \langle \vec{e} \vec{e}^\dagger \rangle = \begin{bmatrix} \langle e_1 e_1^* \rangle & \langle e_1 e_2^* \rangle \\ \langle e_2 e_1^* \rangle & \langle e_2 e_2^* \rangle \end{bmatrix}, \quad (3.67)$$

the averages being taken over the response time of the detector, exactly as in the construction of the Stokes parameters themselves.

Take now the four Hermitian matrices

$$\sigma_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \sigma_1 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}, \quad (3.68)$$

the identity together with the three Pauli matrices, ordered and signed to match the conventions we have been using. Any Hermitian 2×2 matrix can be expanded in this set, and carrying out the four traces term by term against 3.67 reproduces exactly the definitions 3.61 to 3.64,

$$S_\mu = \text{tr}(\Phi \sigma_\mu), \quad \Phi = \frac{1}{2} \sum_{\mu=0}^3 S_\mu \sigma_\mu. \quad (3.69)$$

The S_μ are therefore the coordinates of a Hermitian *operator* in the basis $\{\sigma_\mu\}$, which is a tensorial object living in an internal, abstract space, and not a vector of the laboratory. This reading also explains, in one line, the inequality that partially polarised light forced upon us, because writing 3.69 out gives

$$\Phi = \frac{1}{2} \begin{bmatrix} S_0 + S_1 & S_2 + iS_3 \\ S_2 - iS_3 & S_0 - S_1 \end{bmatrix}, \quad \det \Phi = \frac{1}{4} (S_0^2 - S_1^2 - S_2^2 - S_3^2) \geq 0,$$

the determinant being non negative precisely because Φ is positive semidefinite, that is, because no measurement of intensity may ever return a negative number. That single physical demand is the entire content of $S_1^2 + S_2^2 + S_3^2 \leq S_0^2$, with equality, a vanishing determinant and a rank one matrix, describing fully polarised light.

There is, finally, one honest sense in which the word vector can be rescued. The combination $S_0^2 - S_1^2 - S_2^2 - S_3^2$ that has just appeared is exactly a Minkowski interval, with S_0 in the role of the time component and (S_1, S_2, S_3) in that of the spatial one, and the physical states fill the forward light cone of an abstract four dimensional space: fully polarised light sits on the cone itself, where the interval vanishes, partially polarised light strictly inside it, and completely unpolarised light on the time axis, where the three spatial components vanish together. The real matrices that describe the action of optical elements on this four tuple, the Mueller matrices to which we will come later, then turn out to belong to a group closely related to the Lorentz group, an observation already made by McMaster when he rescued the formalism (McMaster, 1954). The Stokes four tuple is a four vector of that abstract space, never a three vector of the laboratory, and keeping the traditional name is harmless as long as we remember which space it is a vector of.

3.5.3 The bridge between Jones and Stokes: the stereographic projection

We now own two different languages for one and the same physical object. The Jones formalism describes a polarisation state with two complex numbers 3.34, and the Stokes formalism describes it with four real ones, 3.61 to 3.64, which for fully polarised light are constrained by 3.60 to live on the surface of the Poincaré sphere 3.5. Two languages that speak about the same thing must be translatable into each other, and the translation here is not a mere change of symbols: it is a piece of geometry, and a very old and very beautiful one, the stereographic projection. What follows is a deliberately phenomenological tour of that bridge, close in spirit to the way Azzam and Bashara present it in their treatise on polarised light (Azzam and Bashara, 1977), where the reader will also find all the fine print we happily skip here.

3.5.3.1 What the stereographic projection is

Stereographic projection is the oldest recipe known for drawing a sphere on a flat sheet of paper. Take a sphere resting on a plane, single out the point N of the sphere diametrically opposite to the point of contact, and call it the centre of projection. To find where a point P of the sphere lands on the paper, draw the straight line joining N to P and follow it until it pierces the plane; the piercing point Q is the image of P . That is the whole construction, and figure 3.6 shows it at a glance.

Definition 15 (*Stereographic projection*): Let S^2 be a sphere resting on a plane $\mathbb{R}^2$, and let $N \in S^2$ be the point diametrically opposite to the point of contact. The stereographic projection from N assigns to every point $P \in S^2$, with $P \neq N$, the unique point $Q \in \mathbb{R}^2$ at which the straight line through N and P meets the plane. The map is one to one, it sends circles drawn on the sphere to circles or straight lines drawn on the plane, and it preserves the angle at which any two curves cross. The centre N alone has no image, so the plane is completed with a single extra point, the point at infinity, declared to be the image of N .




Figure 3.6: The stereographic projection. A ray leaving the centre of projection N and passing through a point P of the sphere S^2 meets the plane $\mathbb{R}^2$ at a single point Q . Points near the contact point fall near the origin, and points near N fly off towards infinity.

The last clause of the definition is the one that carries all the physics. The plane by itself is not enough to hold the whole sphere, since one point of the sphere always escapes; but as soon as we agree that all the ways of running off to infinity on the plane are one and the same destination, the accounting becomes exact and the sphere and the completed plane become the same object seen twice. This completed plane, $\mathbb{C} \cup \{\infty\}$, is what mathematicians call the extended complex plane, and the sphere carrying it is what they call the Riemann sphere. The construction is far older than that name: Hipparchus and Ptolemy already used it to draw star charts, and the medieval astrolabe is nothing but a stereographic map of the celestial sphere engraved on a brass disc, which works precisely because the projection turns circles into circles and keeps angles honest.

3.5.3.2 Why a polarisation state is a single complex number

Nothing in the previous paragraph mentions light, so we must now ask what a sphere and a plane have to do with polarisation. The answer starts with a simple count. The Jones spinor 3.34 carries two complex numbers, that is, four real ones, the two amplitudes $\mathcal{E}_1, \mathcal{E}_2$ and the two phases ϕ_1, ϕ_2 ; but two of those four say nothing whatsoever about polarisation. The overall size of the spinor is the intensity of the beam, and dimming the light does not change the shape of its ellipse; the overall phase is only a choice of the origin of time, and starting the clock a little earlier does not change that shape either. Multiplying the whole spinor by any complex number therefore leaves the polarisation state untouched, and what survives such a multiplication is only the *ratio* of the two components.

Definition 16 (*Complex polarisation ratio*): Given the Jones spinor $\vec{e} = [e_1, e_2]^T$ of a fully polarised beam 3.34, its complex polarisation ratio is the number

$$w \equiv \frac{e_2}{e_1} = \frac{\mathcal{E}_2}{\mathcal{E}_1} e^{i(\phi_2 - \phi_1)} = \frac{\mathcal{E}_2}{\mathcal{E}_1} e^{-i\Delta\phi} \in \mathbb{C} \cup \{\infty\}, \quad (3.70)$$

with the value $w = \infty$ reserved for the case $e_1 = 0$. Two Jones spinors describe the same polarisation state if and only if they share the same w .

Both pieces of w are quantities we have already been using throughout this chapter, only now packed into one symbol. Its modulus $|w| = \mathcal{E}_2/\mathcal{E}_1$ is the ratio of the two amplitudes, which decides how elongated the ellipse is and towards which axis it leans, and its argument $\arg w = -\Delta\phi$ is the phase lag between the two oscillations, which decides how open the ellipse is and in which sense the field turns. Feeding the six canonical states of the previous subsection into 3.70 gives, immediately,

$$w_H = 0, \quad w_V = \infty, \quad w_D = 1, \quad w_A = -1, \quad w_{\pm} = \pm i, \quad (3.71)$$

read off from the spinors 3.44, 3.46, 3.48, 3.50 and 3.52. Notice that the list already forces the point at infinity upon us: vertical polarisation has got no finite ratio at all, since its first component vanishes, and it is a perfectly ordinary state of light that no honest map is allowed to leave out. So the natural home of polarisation is not the plane $\mathbb{C}$ but the completed plane $\mathbb{C} \cup \{\infty\}$, and by the previous subsection that home is a sphere.

3.5.3.3 The Poincaré sphere is that sphere

The sphere we have just been handed is not a new one. Rest the Poincaré sphere on the complex plane of w , so that it touches the plane at the origin, and project from the point N on top, as in figure 3.7. Every reading of the resulting picture is a statement about light.

The point of contact carries $w = 0$, and by 3.71 that is horizontal polarisation; the centre of projection is sent to infinity, and that is vertical polarisation. The two states that our coordinate axes single out therefore sit at the two poles of the picture, diametrically opposite each other, exactly as they should, since they are orthogonal states and we already knew that orthogonal states are




Figure 3.7: The Poincaré sphere resting on the complex plane of the ratio w . Horizontal polarisation sits at the point of contact, $w = 0$, vertical polarisation sits at the centre of projection and is sent to the point at infinity, and the equator, where the two amplitudes are equal, is mapped onto the unit circle $|w| = 1$. The angles 2α and 2χ that locate a state P on the sphere are twice the orientation and ellipticity angles of its ellipse.

antipodal on the Poincaré sphere. Halfway between them lies the equator, the set of states for which the two amplitudes are equal, and it is mapped onto the unit circle $|w| = 1$. On that circle sit the diagonal state at $w = 1$, the antidiagonal at $w = -1$, and the two circular states at $w = \pm i$, each pair again antipodal, again orthogonal. Moving radially outwards from the origin means pouring amplitude into the second component and turning a horizontal state gradually into a vertical one; going around at fixed radius means sliding the phase difference $\Delta\phi$ and opening or closing the ellipse without touching its amplitudes.

The antipodal rule survives the trip down to the plane in a form worth recording. The state orthogonal to $\vec{e} = [e_1, e_2]^T$ is $[-e_2^*, e_1^*]^T$, whose ratio is $e_1^*/(-e_2^*)$, so on the plane the antipodal map reads

$$w \mapsto -\frac{1}{w^*}, \quad (3.72)$$

which indeed exchanges 0 with ∞ , 1 with -1 , and $+i$ with $-i$, that is, horizontal with vertical, diagonal with antidiagonal, and each circular state with the other one. A statement that on the sphere was a plain geometric fact, going to the other side, becomes on the plane a short algebraic rule.

The dictionary between the two formalisms is now explicit. Dividing the Stokes parameters 3.61 to 3.64 by S_0 and writing them in terms of the ratio 3.70 gives

$$\frac{S_1}{S_0} = \frac{1 - |w|^2}{1 + |w|^2}, \quad \frac{S_2}{S_0} = \frac{2 \operatorname{Re}\{w\}}{1 + |w|^2}, \quad \frac{S_3}{S_0} = \frac{-2 \operatorname{Im}\{w\}}{1 + |w|^2}, \quad (3.73)$$

and these three expressions are, letter for letter, the classical formulae of the inverse stereographic projection, the ones that lift a point of the plane back up to the sphere. The sign carried by S_3 is only the mirror convention of the drawing, and says which of the two circular states we agree to place on the upper half of the imaginary axis. Nothing else in 3.73 is a matter of convention: the Jones ratio and the Stokes vector are the same information, once written on the plane and once written on the sphere.

3.5.3.4 A flat atlas of every polarisation state

Because the projection loses nothing, we may throw the sphere away and keep only the map. Figure 3.8 does exactly that: it is the complex plane of w with the polarisation ellipse of each state drawn at the point that represents it, a complete atlas of polarised light on a single sheet.

Three features of the atlas deserve to be read out loud. First, the whole real axis is filled with linear states: a real w means $\Delta\phi = 0$ or $\Delta\phi = \pi$, the two components oscillate in step or in opposition, and the ellipse collapses to a segment, whose tilt sweeps through every possible orientation as w runs from one end of the axis to the other. Second, the imaginary axis carries the states whose ellipse is aligned with the coordinate axes, growing from the flat horizontal segment at the origin, through wider and wider ellipses, to a perfect circle at $w = \pm i$, and then closing again into the vertical segment far away. Third, and most striking, the edges of the atlas in every direction, $+\infty$, $-\infty$, $+i\infty$ and $-i\infty$ alike, all carry the same vertical polarisation, because on the sphere all those escape routes converge on the single point N . The plane looks infinite, but it is not: it closes up on itself, and the object it really depicts is compact, bounded and without edges, a sphere.

This is the sense in which the Jones and the Stokes formalisms are one theory. A polarisation state is a point; call it w and you are doing Jones calculus on the plane, call it $(S_1, S_2, S_3)/S_0$ and you are doing Stokes calculus on the sphere, and 3.73 carries you back and forth at will. The choice between them is a choice of convenience, and each has got its own: the sphere makes orthogonality and the doubling of angles visible at a glance, while the plane reduces a state to a single number that can simply be divided or multiplied. The next subsection exploits precisely this, because once a state is a single complex number, what an optical element does to light becomes a transformation of the plane onto itself, and the geometry of the sphere tells us in advance which transformations are allowed.




Figure 3.8: The flattened Poincaré sphere. Each point of the complex plane of w carries the ellipse of the state it represents. The origin is horizontal polarisation, the real axis carries every linear state, the imaginary axis carries the ellipses aligned with the coordinate axes together with the two circular states, and every direction of escape to infinity leads to one and the same state, the vertical one.

3.5.4 Transformations of Polarisation: Jones matrices

Every polarisation state built so far is a fixed vector in $\mathbb{C}^2$, but light does not stay the same forever: it crosses lenses, reflects off mirrors, is dimmed by tinted glass, or travels through crystals whose very atomic arrangement singles out a preferred direction in space. Every one of these processes, as long as it acts linearly and does not depend on where or when the light arrives (that is, as long as it is a genuine, homogeneous optical element and not, say, a nonlinear medium or a time varying modulator), can only do one thing to the Jones spinor 3.34: multiply it by a fixed matrix.

Definition 17 (*Jones matrix*): A linear, homogeneous optical element acting on a fully polarised, monochromatic beam is represented by a matrix $J \in \mathbb{C}^{2 \times 2}$, its Jones matrix, such that the outgoing spinor 3.34 is related to the incoming one by

$$\vec{e}' = J \vec{e}.$$

Almost every optical element used in a real laboratory falls into one of three families, according to what kind of matrix J it is: retarders, rotators and dichroics. We build all three from the same simple idea, that of writing J in the basis of its own eigenstates, and we let the physics of each device tell us what those eigenvalues should be.

3.5.4.1 Retarders

Physically, a retarder, also called a wave plate, is cut from a birefringent crystal, quartz, calcite or mica, whose internal atomic arrangement is not the same along every direction, so that the refractive index it presents to light depends on how that light is polarised. Every birefringent plate has, built into it, two mutually orthogonal privileged directions, its fast axis and its slow axis, and light polarised exactly along one of them propagates through unchanged in form, picking up only an ordinary propagation phase, but at a different rate along the fast axis, index n_f , than along the slow one, index n_s . After a thickness d , the component travelling along the slow axis has fallen behind the one along the fast axis by a phase

$$\delta = \frac{2\pi d}{\lambda} (n_s - n_f), \quad (3.74)$$

called the retardance, while any other input, being a superposition of the two axes, comes out with its ellipse reshaped, because the two halves of the superposition have rotated relative to each other in the complex plane.

Definition 18 (*Retarder*): Let $|\psi_1\rangle = |\alpha; \chi\rangle$ and $|\psi_2\rangle = |\alpha + \pi/2; -\chi\rangle$ be a pair of orthogonal Jones kets 3.41⁴. A retarder of eigenstates $|\psi_1\rangle, |\psi_2\rangle$ and retardance δ is the Jones matrix

$$W(\alpha, \chi; \delta) = e^{i\delta/2} |\psi_1\rangle \langle \psi_1| + e^{-i\delta/2} |\psi_2\rangle \langle \psi_2|. \quad (3.75)$$

Equation 3.75 is exactly the spectral decomposition of W : $|\psi_1\rangle, |\psi_2\rangle$ are, by construction, its eigenvectors, with eigenvalues $e^{\pm i\delta/2}$ of unit modulus, so W is unitary and preserves the total intensity $\vec{e}^\dagger \vec{e}$ of any beam it acts on, exactly as an element that only dephases, and never absorbs, should.

The case $\chi = 0$ is by far the most common, since it is what an ordinary birefringent plate, cut with its fast axis in the plane of the crystal, produces; we call it a linear retarder, because both eigenstates $|\alpha; 0\rangle = [\cos \alpha, \sin \alpha]^\top$ and $|\alpha + \pi/2; 0\rangle = [-\sin \alpha, \cos \alpha]^\top$ are linearly polarised. Substituting them into 3.75 and writing $e^{\pm i\delta/2} = \cos(\delta/2) \pm i \sin(\delta/2)$, the four entries work out to

$$\begin{aligned} W_{11} &= \cos^2 \alpha e^{i\delta/2} + \sin^2 \alpha e^{-i\delta/2} = \cos \frac{\delta}{2} + i \sin \frac{\delta}{2} \cos 2\alpha, \\ W_{22} &= \sin^2 \alpha e^{i\delta/2} + \cos^2 \alpha e^{-i\delta/2} = \cos \frac{\delta}{2} - i \sin \frac{\delta}{2} \cos 2\alpha, \\ W_{12} &= W_{21} = \sin \alpha \cos \alpha (e^{i\delta/2} - e^{-i\delta/2}) = i \sin \frac{\delta}{2} \sin 2\alpha, \end{aligned}$$

⁴Writing out the components of $|\alpha; \chi\rangle$ and $|\alpha + \pi/2; -\chi\rangle$ and multiplying term by term shows that $\langle \psi_1 | \psi_2 \rangle = 0$ identically, for every α and every χ , so this pair is always orthogonal.

so that the linear retarder of fast axis at α and retardance δ is

$$W(\alpha, 0; \delta) = \begin{bmatrix} \cos \frac{\delta}{2} + i \sin \frac{\delta}{2} \cos 2\alpha & i \sin \frac{\delta}{2} \sin 2\alpha \\ i \sin \frac{\delta}{2} \sin 2\alpha & \cos \frac{\delta}{2} - i \sin \frac{\delta}{2} \cos 2\alpha \end{bmatrix}. \quad (3.76)$$




Figure 3.9: A quarter wave plate retarder is a retarder with $\delta = \pi/2$ and when it has got its fast axis oriented with the horizontal one it transforms the polarization in a very specific way. The $\pm 45^\circ$ diagonal states transform into circular polarization states, and the circular polarization states transform into $\pm 45^\circ$ diagonal states.

Two particular values of δ are so useful that they have earned their own names. When $\delta = \pi/2$, a quarter of a full cycle, we speak of a quarter wave plate (see figure 3.9); taking the fast axis horizontal, $\alpha = 0$, equation 3.76 becomes, up to the irrelevant global phase $e^{i\pi/4}$,

$$W_{\lambda/4} = \begin{bmatrix} 1 & 0 \\ 0 & -i \end{bmatrix}, \quad (3.77)$$

whose action on the diagonal state 3.49 is immediate to check: $|\pi/4; 0\rangle = \frac{1}{\sqrt{2}}[1, 1]^T$ is sent to $\frac{1}{\sqrt{2}}[1, -i]^T = |0; -\pi/4\rangle$, one of the circular states 3.53. This is exactly why a quarter wave plate is the standard tool for turning linear polarisation into circular, and back.

When instead $\delta = \pi$, half a full cycle, we get the half wave plate,

$$W(\alpha, 0; \pi) = i \begin{bmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{bmatrix}, \quad (3.78)$$

and, up to the global phase i , this matrix is a reflection: acting on a linear state it sends $|\theta; 0\rangle$ to $|2\alpha - \theta; 0\rangle$, reflecting its orientation about the fast axis, so a half wave plate rotates any linear input by twice the angle between the input and the fast axis, and it is the tool of choice whenever an arbitrary rotation of a linear polarisation is needed. Acting instead on a circular state, a short calculation with 3.78 shows that $|0; \pi/4\rangle$ is sent to $|0; -\pi/4\rangle$ for every value of α , so a half wave plate always reverses the handedness of a circular input, whatever its own orientation happens to be.

Nothing prevents us from choosing genuinely elliptical eigenstates, $0 < |\chi| < \pi/4$: the resulting elliptical retarder is a real, if less common, device, built from a birefringent material whose axes have additionally been twisted by a small optical activity, so that its privileged states are ellipses rather than straight lines, and it is worth constructing its Jones matrix properly, both because it is the fully general case of 3.75 and because doing so will let us obtain the rotator, further on, in a single line.

Write the first eigenstate $|\psi_1\rangle = |\alpha; \chi\rangle$ of 3.41 as $|\psi_1\rangle = [a, b]^T$ with

$$a = \cos \alpha \cos \chi - i \sin \alpha \sin \chi, \quad b = \sin \alpha \cos \chi + i \cos \alpha \sin \chi,$$

so that $|a|^2 + |b|^2 = 1$, since $|\psi_1\rangle$ is a unit vector. Comparing component by component with $|\psi_2\rangle = |\alpha + \pi/2; -\chi\rangle$ shows the compact relation $|\psi_2\rangle = [-b^*, a^*]^T$, so that the matrix built from the two orthonormal eigenstates as columns,

$$U(\alpha, \chi) = [|\psi_1\rangle \quad |\psi_2\rangle] = \begin{bmatrix} a & -b^* \\ b & a^* \end{bmatrix}, \quad (3.79)$$

is unitary, $U^\dagger U = U U^\dagger = I$, precisely because $|\psi_1\rangle, |\psi_2\rangle$ are orthonormal. Equation 3.79 is nothing but the change of basis matrix from the eigenbasis $\{|\psi_1\rangle, |\psi_2\rangle\}$, in which the retarder is diagonal by definition, to the standard basis in which we write the Jones spinor 3.34. Writing $\Lambda(\delta) = \text{diag}(e^{i\delta/2}, e^{-i\delta/2})$ for the retarder expressed in its own eigenbasis, this change of basis gives a second, completely explicit way of writing 3.75,

$$W(\alpha, \chi; \delta) = U(\alpha, \chi) \Lambda(\delta) U(\alpha, \chi)^\dagger, \quad (3.80)$$

and carrying out the product explicitly,

$$U \Lambda U^\dagger = \begin{bmatrix} a & -b^* \\ b & a^* \end{bmatrix} \begin{bmatrix} e^{i\delta/2} & 0 \\ 0 & e^{-i\delta/2} \end{bmatrix} \begin{bmatrix} a^* & b^* \\ -b & a \end{bmatrix} = \begin{bmatrix} |a|^2 e^{i\delta/2} + |b|^2 e^{-i\delta/2} & ab^* (e^{i\delta/2} - e^{-i\delta/2}) \\ a^* b (e^{i\delta/2} - e^{-i\delta/2}) & |b|^2 e^{i\delta/2} + |a|^2 e^{-i\delta/2} \end{bmatrix}.$$

Squaring and multiplying the two expressions for a, b above gives $|a|^2 - |b|^2 = \cos 2\alpha \cos 2\chi$, $|a|^2 + |b|^2 = 1$ and $ab^* = \frac{1}{2} \sin 2\alpha \cos 2\chi - \frac{i}{2} \sin 2\chi$, exactly the same three combinations already met while building the linear retarder; using them, together with $e^{\pm i\delta/2} = \cos(\delta/2) \pm i \sin(\delta/2)$, the elliptical retarder takes the closed form

$$W(\alpha, \chi; \delta) = \begin{bmatrix} \cos \frac{\delta}{2} + i \sin \frac{\delta}{2} \cos 2\alpha \cos 2\chi & \sin \frac{\delta}{2} \sin 2\chi + i \sin \frac{\delta}{2} \sin 2\alpha \cos 2\chi \\ -\sin \frac{\delta}{2} \sin 2\chi + i \sin \frac{\delta}{2} \sin 2\alpha \cos 2\chi & \cos \frac{\delta}{2} - i \sin \frac{\delta}{2} \cos 2\alpha \cos 2\chi \end{bmatrix}. \quad (3.81)$$

Setting $\chi = 0$ in 3.81 gives $\cos 2\chi = 1$, $\sin 2\chi = 0$ and collapses it exactly back to the linear retarder 3.76, as it must: the fully general elliptical case was, after all, only ever hiding the extra angle χ inside the change of basis matrix 3.79.

3.5.4.2 Rotators

Let us push the general elliptical retarder 3.81 to the opposite extreme from the linear case and choose circular eigenstates, $\alpha = 0, \chi = \pi/4$, that is $|\psi_1\rangle = |0; \pi/4\rangle$ and $|\psi_2\rangle = |0; -\pi/4\rangle$ of 3.53. Since $\cos 2\chi = \cos(\pi/2) = 0$ and $\sin 2\chi = \sin(\pi/2) = 1$, every term in 3.81 carrying a factor of $\cos 2\chi$ vanishes identically and we are left with

$$W(0, \pi/4; \delta) = \begin{bmatrix} \cos \frac{\delta}{2} & \sin \frac{\delta}{2} \\ -\sin \frac{\delta}{2} & \cos \frac{\delta}{2} \end{bmatrix}, \quad (3.82)$$

and the right hand side is exactly the rotation matrix already used in 3.41 to build every ellipse out of the unrotated form 3.40. A retarder of circular eigenstates, in other words, is nothing but a rotation of the whole Jones plane: it sends $|\alpha; \chi\rangle$ to $|\alpha - \delta/2; \chi\rangle$ for every α and every χ , leaving the ellipticity untouched and shifting only the orientation.

Definition 19 (*Rotator*): A rotator by an angle θ is the Jones matrix

$$R(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}, \quad (3.83)$$

equivalently the circular retarder 3.82 of retardance $\delta = -2\theta$, so that $R(\theta) |\alpha; \chi\rangle = |\alpha + \theta; \chi\rangle$ for every α, χ .

Physically, this is exactly what happens inside an optically active medium: a solution of chiral sugar molecules, or a slab of quartz cut along its optic axis, where it is not the two linear directions that see different refractive indices, but the two circular ones. The phenomenon is called circular birefringence, and a beam of any polarisation crossing a thickness d of such a medium has its ellipse rotated, rigidly and without distortion, by an angle proportional to d ; the same net effect, a pure rotation with no change of ellipticity, is produced by a Faraday rotator, a piece of glass or crystal placed in a strong magnetic field along the beam, in which it is the magnetic field rather than molecular chirality that splits the two circular eigenstates. Notice that $R(\theta)$, unlike a generic retarder, is a real matrix: rotators are the only non-trivial polarisation transformations whose Jones matrix has got no imaginary part at all, a reflection of the fact that neither optical activity nor the Faraday effect privileges any particular linear direction in space, only a sense of circulation.

3.5.4.3 Dichroics and polarisers

Every element considered so far is unitary: it reshapes the polarisation ellipse but never changes the total intensity. A dichroic element breaks this on purpose. Physically, it is again built from an anisotropic material, but now one whose two privileged directions are absorbed differently rather than merely retarded differently, because the microscopic charges responsible for the absorption,

free electrons in a metal wire, iodine chains in a stretched polymer, the delocalised electrons of an elongated organic molecule, can oscillate much more easily along one direction than along the perpendicular one. Mathematically we only have to repeat the construction 3.75 once more, replacing the two phase factors of unit modulus by two real, non-negative amplitude transmittances.

Definition 20 (*Dichroic*): A dichroic element of eigenstates $|\psi_1\rangle, |\psi_2\rangle$ (orthogonal, as in the retarder) and amplitude transmittances $t_1, t_2 \in \mathbb{R}$, $t_1, t_2 \geq 0$, is the Jones matrix

$$D = t_1 |\psi_1\rangle \langle \psi_1| + t_2 |\psi_2\rangle \langle \psi_2|. \quad (3.84)$$

Because t_1, t_2 are real, D in 3.84 is Hermitian rather than unitary, unless $t_1 = t_2$, the trivial case in which nothing happens to the polarisation, only to the overall brightness: light along $|\psi_1\rangle$ comes out multiplied by t_1 and light along $|\psi_2\rangle$ by t_2 , and any state in between comes out reshaped towards whichever eigenstate survives best.

Taking $\chi = 0$ once again gives the linear dichroic, and repeating the same short calculation that led to 3.76, now with t_1, t_2 in place of $e^{\pm i\delta/2}$,

$$D(\alpha, 0; t_1, t_2) = \begin{bmatrix} t_1 \cos^2 \alpha + t_2 \sin^2 \alpha & (t_1 - t_2) \sin \alpha \cos \alpha \\ (t_1 - t_2) \sin \alpha \cos \alpha & t_1 \sin^2 \alpha + t_2 \cos^2 \alpha \end{bmatrix}. \quad (3.85)$$

The most important member of this family is obtained by pushing the contrast between the two eigenstates to its limit, $t_1 = 1$ and $t_2 = 0$: the surviving axis is transmitted perfectly and the other is extinguished completely. Equation 3.85 then collapses to

$$P(\alpha) = \begin{bmatrix} \cos^2 \alpha & \sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha & \sin^2 \alpha \end{bmatrix} = |\alpha; 0\rangle \langle \alpha; 0|, \quad (3.86)$$

which we recognise, on the right, as the orthogonal projector onto the linear state $|\alpha; 0\rangle$.

Definition 21 (*Ideal linear polariser*): An ideal linear polariser of transmission axis at angle α is the limiting linear dichroic 3.85 with $t_1 = 1, t_2 = 0$, that is, the orthogonal projector $P(\alpha) = |\alpha; 0\rangle \langle \alpha; 0|$ of 3.86.

This is the deepest sense in which a polariser is a simple device: it does not really transform a polarisation state, it projects it, discarding every component along $|\alpha + \pi/2; 0\rangle$ and keeping only the shadow that the incoming ellipse casts onto its own transmission axis. Every partial linear dichroic 3.85 with $t_2 \neq 0$ is only an imperfect version of this same projection, letting some of the unwanted component leak through.

Circular dichroism is built the same way, with $|\psi_1\rangle = |0; \pi/4\rangle$ and $|\psi_2\rangle = |0; -\pi/4\rangle$ in 3.84. Specialising a, b from the change of basis matrix 3.79 to $\alpha = 0, \chi = \pi/4$ gives $a = 1/\sqrt{2}$, $b = i/\sqrt{2}$, so




Figure 3.10: A linear polariser with its polariser axis oriented parallel to the horizontal axis projects all the incident states to horizontal ones, however the transmitted intensity is reduced by the relative angle between the incident state $|\alpha; \chi\rangle$ and the polariser axis α_{pol} in this case $\alpha_{pol} = 0^\circ$.

that $|a|^2 = |b|^2 = 1/2$ and $ab^* = -i/2$; repeating, with these values, the same matrix multiplication that led to 3.81, but with the real transmittances t_1, t_2 in place of the phases $e^{\pm i\delta/2}$,

$$D_{\odot}(t_1, t_2) = \begin{bmatrix} \frac{t_1 + t_2}{2} & -i \frac{t_1 - t_2}{2} \\ i \frac{t_1 - t_2}{2} & \frac{t_1 + t_2}{2} \end{bmatrix}, \quad (3.87)$$

describing a material that absorbs its two circular eigenstates differently. Circular dichroism is a much smaller effect than linear dichroism in most materials, but it is an extremely sensitive probe of molecular chirality, proteins, sugars and DNA are all optically active and mildly circularly dichroic, which is why circular dichroism spectroscopy is a standard tool in structural biology and chemistry.

Stepping back from the algebra, what actually makes a material dichroic is always some built in anisotropy at the microscopic level: an elongated molecule, a stretched polymer chain, or a crystal lattice, along which charges are freer to respond to the oscillating electric field of light in one direction than in the perpendicular one, so the material absorbs, rather than merely slows down, one polarisation component more strongly than the other. A polariser, more generally, is simply any device, dichroic or not, designed to output light of a well controlled polarisation state from an input that may not have one at all; dichroism is only one of the physical mechanisms that nature and engineering have found to build one.

Three real polarisers illustrate how different those mechanisms can be. Tourmaline, the mineral in which dichroism was first noticed, is naturally dichroic, but the polariser found today in almost every pair of sunglasses, camera filter and liquid crystal display is Edwin Land's Polaroid sheet, a film of polyvinyl alcohol stretched in one direction to align its long polymer chains and then doped with iodine, whose mobile electrons run freely along the chains and absorb the field component parallel to them, the linear dichroic 3.85 realised in a plastic film. A wire grid polariser applies the same physical idea to a metal instead of a dye: an array of parallel conducting wires, spaced much closer together than the wavelength of the light, lets the free electrons in the wires oscillate and reflect the field component parallel to them, while the perpendicular component, unable to drive

a current across the gaps between wires, passes through almost unaffected; because it works by reflection rather than absorption, it survives far higher optical powers than a dyed polymer sheet, and it is the polariser of choice in the infrared and in microwave optics. Finally, the Glan-Thompson prism takes a completely different route, using no absorption at all: two prisms of calcite, a strongly birefringent crystal, are cut and cemented together so that, at the internal interface, the ordinary ray strikes beyond its critical angle and is totally internally reflected out of the beam, while the extraordinary ray, seeing a different refractive index, is transmitted straight through. Being lossless for the transmitted beam and free of any absorbing dye, this kind of polariser can handle the intense, focused light of a laser without being damaged, which the simple dichroic sheet cannot.

3.5.5 Transformations of Polarisation: Mueller matrices

The Jones calculus of the previous subsection is exact, elegant and, for a large class of problems, entirely sufficient. It is also, in a very precise sense, too narrow. A Jones matrix acts on the spinor 3.34, and that spinor exists only when the beam has got a single, well defined phase relation between its two components, that is, only when the light is fully polarised. The moment we point our instrument at sunlight, at a cloud, at a sheet of paper, at a biological tissue or at any surface rough on the scale of a wavelength, the object we must transform is no longer a spinor but the four Stokes parameters 3.61 to 3.64, and no 2×2 complex matrix will do the job.

There is a second and deeper failure, and it is worth stating before we build anything. A Jones matrix can never *depolarise*. Whatever it does to a fully polarised beam, the outgoing beam is again fully polarised, so a formalism built on Jones matrices alone cannot describe the single most common thing that real matter does to light. We shall prove this in a moment, and the proof will also tell us exactly what has to be added.

The formalism that repairs both defects was assembled by Hans Mueller at the Massachusetts Institute of Technology in the early nineteen forties, building on the earlier scattering work of Soleillet and of Perrin (Perrin, 1942), and it was the same wave of activity that rescued the Stokes parameters from the obscurity described at the beginning of this section. Its later systematisation owes a great deal to McMaster (McMaster, 1954) and to Chandrasekhar's treatment of radiative transfer (Chandrasekhar, 1950), where polarised light must be propagated through a scattering atmosphere and where Jones spinors are simply not available.

3.5.5.1 From the Jones matrix to the coherence matrix

The bridge between the two calculi is the polarisation matrix 3.67, and building it is a one line calculation whose consequences occupy the rest of this subsection. Let a linear, homogeneous element with Jones matrix J act on the beam, so that $\vec{e}' = J\vec{e}$. The outgoing polarisation matrix is, by definition,

$$\Phi' = \langle \vec{e}' \vec{e}'^\dagger \rangle = \langle J\vec{e} (J\vec{e})^\dagger \rangle = \langle J\vec{e} \vec{e}^\dagger J^\dagger \rangle = J \langle \vec{e} \vec{e}^\dagger \rangle J^\dagger = J \Phi J^\dagger. \quad (3.88)$$

Every step is elementary, but the third one is the whole point and deserves to be said out loud: the matrix J could be taken outside the time average because the *element* is deterministic even

when the *light* is not. The waveplate does not flicker; only the field does. This is precisely why 3.88 remains valid for partially polarised and even for completely unpolarised light, where the spinor $\vec{e}$ has got no meaning at all and only the averages survive.

Notice also the shape of the transformation. The field transforms linearly, $\vec{e} \rightarrow J\vec{e}$, but the polarisation matrix transforms by a **congruence**, $\Phi \rightarrow J\Phi J^\dagger$, with J appearing twice. Since the Stokes parameters are themselves quadratic in the field, this is exactly what we should have expected, and it is the algebraic reason why the Mueller matrix we are about to build is quadratic in the Jones matrix.

3.5.5.2 The Mueller matrix

Because the Stokes parameters are linear coordinates of Φ through 3.69, and because 3.88 is linear in Φ , the map induced on the four Stokes parameters must be linear as well.

Definition 22 (*Mueller matrix*): A linear, homogeneous optical element acting on a beam of arbitrary degree of polarisation is represented by a real 4×4 matrix M , its *Mueller matrix*, such that the outgoing Stokes vector is related to the incoming one by

$$S'_\mu = \sum_{\nu=0}^3 M_{\mu\nu} S_\nu, \quad \text{that is} \quad \vec{S}' = M\vec{S}. \quad (3.89)$$

When the element also possesses a Jones matrix, the two descriptions must agree, and forcing them to agree hands us the Mueller matrix explicitly. Insert 3.88 into the first of 3.69 and then expand the incoming Φ with the second,

$$S'_\mu = \text{tr}(\Phi' \sigma_\mu) = \text{tr}(J\Phi J^\dagger \sigma_\mu) = \text{tr}\left(J \left[\frac{1}{2} \sum_{\nu=0}^3 S_\nu \sigma_\nu \right] J^\dagger \sigma_\mu\right) = \sum_{\nu=0}^3 \left[\frac{1}{2} \text{tr}(\sigma_\mu J \sigma_\nu J^\dagger) \right] S_\nu,$$

where the trace has been taken through the finite sum and the cyclic property $\text{tr}(AB) = \text{tr}(BA)$ has been used to bring σ_μ to the front. Comparing with 3.89 we can read off the coefficients at once.

Definition 23 (*Mueller matrix of a Jones element*): An element whose action on fully polarised light is given by the Jones matrix J has got the *Mueller matrix*

$$M_{\mu\nu} = \frac{1}{2} \text{tr}(\sigma_\mu J \sigma_\nu J^\dagger) \quad (3.90)$$

with σ_μ the four matrices 3.68. A Mueller matrix that can be written in this form is called **non depolarising**, or a **Mueller-Jones matrix**.

Formula 3.90 is the dictionary we were after, and three of its properties are worth extracting immediately.

Taking the complex conjugate of 3.90 and using $\overline{\text{tr } A} = \text{tr } A^\dagger$ together with the hermiticity $\sigma_\mu^\dagger = \sigma_\mu$,

$$\overline{M_{\mu\nu}} = \frac{1}{2} \text{tr} \left(\left[\sigma_\mu J \sigma_\nu J^\dagger \right]^\dagger \right) = \frac{1}{2} \text{tr} \left(J \sigma_\nu J^\dagger \sigma_\mu \right) = M_{\mu\nu},$$

the last step being once more the cyclic property. So M is a real matrix, as it must be, since it maps four measurable intensities into four measurable intensities.

Replacing J by $e^{i\theta} J$ leaves 3.90 untouched, because the phase from J cancels against the conjugate phase from $J^\dagger$. The map $J \mapsto M$ is therefore not injective: all Jones matrices differing by an overall phase collapse onto the same Mueller matrix. This is not a defect but an improvement, since that phase was never measurable in the first place, and we shall see in a moment that it is the same two to one correspondence that produces the doubling of angles on the Poincaré sphere.

If light crosses first the element J_1 and then the element J_2 , the combined Jones matrix is $J_2 J_1$, and substituting it into 3.90 while inserting the resolution $\frac{1}{2} \sum_\lambda \sigma_\lambda \text{tr}(\sigma_\lambda \cdot)$ between the two factors shows that

$$M(J_2 J_1) = M(J_2) M(J_1), \quad (3.91)$$

so that 3.90 is a group homomorphism and an optical bench is described, exactly as in the Jones calculus, by the ordered product of the matrices of its elements.

3.5.5.3 Retarders are rotations of the Poincaré sphere

Let us now compute a Mueller matrix explicitly, and let us choose the simplest retarder, the linear one with its fast axis along x , whose Jones matrix is 3.76 with $\alpha = 0$, that is $W = \text{diag}(e^{i\delta/2}, e^{-i\delta/2})$. We need the four objects $W \sigma_\nu W^\dagger$. The first two are immediate, since W is unitary and diagonal and therefore commutes with σ_0 and σ_1 ,

$$W \sigma_0 W^\dagger = W W^\dagger = \sigma_0, \quad W \sigma_1 W^\dagger = \sigma_1.$$

The interesting ones are the other two. Multiplying out,

$$W \sigma_2 W^\dagger = \begin{bmatrix} 0 & e^{i\delta} \\ e^{-i\delta} & 0 \end{bmatrix} = \cos \delta \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \sin \delta \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix} = \cos \delta \sigma_2 + \sin \delta \sigma_3,$$

where we simply split $e^{\pm i\delta} = \cos \delta \pm i \sin \delta$ and recognised the two Pauli matrices, and in exactly the same way

$$W \sigma_3 W^\dagger = \begin{bmatrix} 0 & i e^{i\delta} \\ -i e^{-i\delta} & 0 \end{bmatrix} = -\sin \delta \sigma_2 + \cos \delta \sigma_3.$$

Since $\frac{1}{2} \text{tr}(\sigma_\mu \sigma_\nu) = \delta_{\mu\nu}$, the coefficients just found are literally the columns of M , and we obtain

$$M_W(\delta) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \cos \delta & -\sin \delta \\ 0 & 0 & \sin \delta & \cos \delta \end{bmatrix}. \quad (3.92)$$

Read this matrix slowly, because it is the central geometrical fact of the whole subsection. The first row and the first column are those of the identity, so S_0 is untouched: a retarder absorbs nothing, exactly as its unitarity promised. The remaining 3×3 block leaves S_1 alone and rotates the pair (S_2, S_3) by the angle δ . In other words, on the Poincaré sphere the element performs a **rigid rotation by the retardance δ about the S_1 axis**, and the S_1 axis is precisely the diameter joining the two eigenstates of this retarder, the horizontal and vertical states 3.45 and 3.47. The eigenstates are the fixed points of the motion, as fixed points of a rotation must be, and every other state is carried around them on a circle of constant latitude.

Nothing in this conclusion depends on our having chosen $\alpha = 0$. Repeating the computation with the general elliptical retarder 3.81 is a longer but entirely mechanical exercise, and it yields the following statement, which we record as the geometrical heart of the Mueller calculus.

Theorem 1 (*Retarders as rotations*): *The Mueller matrix of a retarder of retardance δ whose eigenstates are $|\alpha; \chi\rangle$ and its orthogonal partner is*

$$M_W = \begin{bmatrix} 1 & \vec{0}^\top \\ \vec{0} & R(\hat{n}, \delta) \end{bmatrix}, \quad (3.93)$$

where $R(\hat{n}, \delta)$ is the ordinary 3×3 rotation by the angle δ about the unit vector

$$\hat{n} = (\cos 2\alpha \cos 2\chi, \sin 2\alpha \cos 2\chi, -\sin 2\chi), \quad (3.94)$$

which by 3.65 is exactly the point of the Poincaré sphere representing the eigenstate $|\alpha; \chi\rangle$.

Every waveplate on an optical bench is therefore a rotation of the Poincaré sphere (see figure 3.11), and the two numbers that specify it have got transparent meanings: the orientation of its axis fixes *about which diameter* the sphere turns, and its retardance fixes *by how much*. Three familiar devices become obvious.

A **quarter wave plate**, $\delta = \pi/2$, turns the sphere by a quarter of a full turn. Starting from a linear state on the equator and choosing the axis at $\pi/4$ from it, a quarter turn lands exactly on a pole, which is the statement that a quarter wave plate converts linear light into circular light, as we found by direct matrix multiplication in the previous subsection.

A **half wave plate**, $\delta = \pi$, turns the sphere by half a full turn about an equatorial axis. A rotation by π about the axis $\hat{n}$ sends any vector $\vec{v}$ to $2(\hat{n} \cdot \vec{v})\hat{n} - \vec{v}$, so a linear state at azimuth 2θ on the equator, acted on by a plate whose axis sits at azimuth 2α , is carried to azimuth $4\alpha - 2\theta$. Dividing by two, the physical polarisation direction goes from θ to $2\alpha - \theta$: **a half wave plate reflects the plane of polarisation about its own fast axis**. Applying the same rule to the poles, $\vec{v} = (0, 0, \pm 1)$ with $\hat{n}$ equatorial gives $\vec{v} \rightarrow -\vec{v}$, so the two circular states are exchanged and the handedness of any beam is reversed.

A **rotator** 3.83 is, as we proved, a circular retarder, so its axis 3.94 is the polar axis $\hat{n} = (0, 0, -1)$ and it spins the sphere about the line joining the two circular states. A rotator that turns the




Figure 3.11: An initial state $|\alpha; \chi\rangle$ under the effect of a rotated retarder with retard δ and eigenstate r is rotated an angle of δ around the r -axis to a final state $|\alpha'; \chi'\rangle$.

physical polarisation by an angle θ has got retardance $\delta = 2\theta$, so it rotates the Poincaré sphere by 2θ . This is the doubling of angles once more, and we can now see its true origin: the map 3.90 restricted to unitary Jones matrices is the classical two to one homomorphism from $SU(2)$ onto the rotation group $SO(3)$. Turning a waveplate through a full physical revolution, $\theta : 0 \rightarrow \pi$, returns the Jones matrix not to $+I$ but to $-I$, while the Mueller matrix has already completed a full 2π rotation and returned to the identity. The polarisation spinor is a spinor in exactly the sense that an electron wavefunction is, and the Poincaré sphere is its Bloch sphere.

3.5.5.4 Polarisers, dichroics and the Malus law

Not every element is unitary, and the non unitary ones move the sphere in a completely different way. Take first the ideal linear polariser 3.86, whose Jones matrix is the projector $J = |\psi\rangle\langle\psi|$ onto the transmitted state. Feeding it into 3.90 and writing $\hat{n}$ for the Stokes vector of that state, the result is the strikingly simple

$$M_P = \frac{1}{2} \begin{bmatrix} 1 & \hat{n}^T \\ \hat{n} & \hat{n}\hat{n}^T \end{bmatrix}. \quad (3.95)$$

This matrix has got rank one in a strong sense: whatever comes in, the outgoing Stokes vector is proportional to $(1, \hat{n})$, that is, the entire Poincaré sphere is crushed onto the single point $\hat{n}$. The map is catastrophically non invertible, which is the geometric expression of the obvious physical fact that a polariser destroys information and cannot be undone.

The Malus law now drops out with no effort at all. Send in linear light at an angle θ , whose Stokes vector is $\vec{S} = (1, \cos 2\theta, \sin 2\theta, 0)$, through a polariser whose axis is at α , so that $\hat{n} = (\cos 2\alpha, \sin 2\alpha, 0)$. The transmitted intensity is the zeroth component of 3.95 acting on $\vec{S}$,

$$S'_0 = \frac{1}{2} \left(1 + \hat{n} \cdot \vec{S}_{123} \right) = \frac{1}{2} [1 + \cos 2\alpha \cos 2\theta + \sin 2\alpha \sin 2\theta] = \frac{1}{2} [1 + \cos (2\theta - 2\alpha)] = \cos^2 (\theta - \alpha),$$

which is the Malus law, obtained here as a statement about the angle between two vectors on the Poincaré sphere rather than as a statement about a projection in the laboratory. Feeding unpolarised light $\vec{S} = (1, 0, 0, 0)$ into the same formula gives $S'_0 = 1/2$ together with a fully polarised output, which is the equally familiar fact that a polariser transmits exactly half of natural light.

The general linear dichroic 3.85, with real transmittances t_1 and t_2 along its two eigenstates, interpolates between the retarder and the polariser and is more interesting still. Taking $J = \text{diag}(t_1, t_2)$ in 3.90 and computing the four congruences as before,

$$M_D = \begin{bmatrix} a & b & 0 & 0 \\ b & a & 0 & 0 \\ 0 & 0 & c & 0 \\ 0 & 0 & 0 & c \end{bmatrix}, \quad a = \frac{t_1^2 + t_2^2}{2}, \quad b = \frac{t_1^2 - t_2^2}{2}, \quad c = t_1 t_2, \quad (3.96)$$

and a one line computation shows that these three numbers are not independent,

$$a^2 - b^2 = \frac{(t_1^2 + t_2^2)^2 - (t_1^2 - t_2^2)^2}{4} = t_1^2 t_2^2 = c^2.$$

The relation $a^2 - b^2 = c^2$ is the signature of a hyperbolic rotation. Writing $a = c \cosh \zeta$ and $b = c \sinh \zeta$, which the constraint permits, 3.96 becomes

$$M_D = t_1 t_2 \begin{bmatrix} \cosh \zeta & \sinh \zeta & 0 & 0 \\ \sinh \zeta & \cosh \zeta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \tanh \zeta = \frac{t_1^2 - t_2^2}{t_1^2 + t_2^2} \equiv D, \quad (3.97)$$

an overall attenuation multiplying a **Lorentz boost** along the S_1 axis, with rapidity ζ and with the quantity D , called the diattenuation, playing the part of the velocity β .

This is the promise made at the end of the discussion of why the Stokes vector is not a vector, now delivered. There we found that the physical states fill the forward light cone of an abstract Minkowski space, with the interval $S_0^2 - S_1^2 - S_2^2 - S_3^2$ measuring the departure from full polarisation. We can now name the transformations: **retarders are the rotations of that Minkowski space and dichroics are its boosts**, and together they generate the proper orthochronous Lorentz group.

The geometry is faithful in every detail. A boost along $\hat{n}$ drags points of the sphere toward $\hat{n}$, crowding them ever closer as the rapidity grows, and this is exactly what a partial polariser does: it does not turn a state, it pulls it toward its own transmission axis. The ideal polariser 3.95 is the limit $t_2 \rightarrow 0$, that is $D \rightarrow 1$ and $\zeta \rightarrow \infty$, the infinite boost that carries every point to the same destination, which is why the sphere collapses.

3.5.5.5 What no Jones matrix can do: depolarisation

We can now prove the claim made at the very beginning. Suppose the incoming light is fully polarised, so that by the discussion of the Stokes identity its polarisation matrix has $\det \Phi = 0$, equivalently $\Phi = \vec{e}\vec{e}^\dagger$ has rank one. Then by 3.88 and the multiplicative property of the determinant,

$$\det \Phi' = \det (J\Phi J^\dagger) = |\det J|^2 \det \Phi = 0, \quad (3.98)$$

so the outgoing light is fully polarised as well. **A Jones element maps pure states to pure states and can never reduce the degree of polarisation.** The Poincaré sphere may be rotated, boosted or crushed, but a point on its surface never moves into its interior.

Yet the interior is where most of the light in the universe lives. The resolution is that a depolarising sample is not one Jones matrix but a *statistical ensemble* of them, fluctuating faster than the detector can follow or varying from point to point across the illuminated spot: the rough surface presents a different microfacet to each part of the beam, the tissue offers a different scattering history to each photon, the turbulent atmosphere a different birefringence at each instant. Averaging 3.88 over that ensemble and using the linearity of 3.90 in the pair $(J, J^\dagger)$,

$$M = \langle M(J) \rangle = \sum_k p_k M(J_k), \quad p_k \geq 0, \quad \sum_k p_k = 1, \quad (3.99)$$

a convex combination of non depolarising matrices. Convex sums of rotations are not rotations, and this is where the degree of polarisation is lost, in exactly the way that averaging unit vectors pointing in different directions was found to shorten the Stokes vector. The archetype is the ideal depolariser

$$M_\Delta = \text{diag}(1, d_1, d_2, d_3), \quad |d_j| \leq 1, \quad (3.100)$$

which shrinks the sphere into an ellipsoid with semiaxes d_1, d_2, d_3 centred on the origin, and which for $d_1 = d_2 = d_3 = 0$ collapses everything onto the centre, that is, onto completely unpolarised light of unchanged intensity.

A parameter count makes the size of the gap plain. A general Mueller matrix carries 16 real numbers. A Jones matrix carries 8, of which one is the unmeasurable global phase, so the non depolarising Mueller matrices form a family of only 7 parameters sitting inside a 16 dimensional space. The overwhelming majority of physically admissible Mueller matrices therefore have got no Jones counterpart whatsoever, and this is the precise sense in which the Mueller calculus is not a reformulation of the Jones calculus but a genuine enlargement of it.

However, we'll see in the chapter of they of coherence that our work in the GOTS can use the Jones matrices to explain with great precision the depolarisation phenomena.

3.5.5.6 Why this matters

It is worth closing with the reason this formalism is used far outside the optics laboratory. A Mueller matrix is 16 real numbers, and every one of them is measurable: illuminate the sample with four known input states, analyse each output with four known projections, and the resulting 16 intensities invert to give M . The technique is called Mueller polarimetry, and what it delivers is a complete linear optical fingerprint of the sample, containing its birefringence, its dichroism and, crucially, its depolarising power, all of them separable from one another by decomposing M into a product of a retarder, a dichroic and a depolariser.

The depolarising part is often the most informative, precisely because it is the part Jones could not describe. Healthy and cancerous tissue differ markedly in how strongly they depolarise, since the difference in the arrangement of collagen fibres changes the statistics of the ensemble in 3.99, and polarimetric imaging is being developed as a label free diagnostic on exactly this basis. In remote sensing the depolarisation of scattered sunlight distinguishes spherical droplets from irregular ice crystals in a cloud, which is why the Stokes formalism entered atmospheric physics through the radiative transfer of Chandrasekhar (Chandrasekhar, 1950). In semiconductor metrology, ellipsometry measures the Mueller matrix of a wafer to determine film thicknesses to sub nanometre precision (Azzam and Bashara, 1977). And in astronomy the polarisation of starlight, scrambled by interstellar dust, maps the galactic magnetic field.

There is finally a structural lesson that reaches well beyond optics. The polarisation matrix Φ is a 2×2 Hermitian, positive semidefinite matrix of unit trace once normalised, which is precisely the definition of the density matrix of a two level quantum system; the Poincaré sphere is its Bloch sphere; a Jones matrix is a unitary evolution; and a general Mueller matrix is the classical counterpart of a quantum channel, a completely positive map, with 3.99 playing the role of the ensemble decomposition of that channel. Depolarisation is decoherence. Everything we have built in this subsection, the rotations, the boosts, the convex mixing that destroys purity, reappears unchanged in the theory of open quantum systems, and a bench with a laser and two waveplates is one of the cheapest and most transparent laboratories in which those ideas can be watched happening.

3.5.6 The Jones theorems

Jones did not merely propose a convenient notation. In the series of papers that runs through the *Journal of the Optical Society of America* between 1941 and 1956, and above all in the second of them, written jointly with Henry Hurwitz (Hurwitz and Jones, 1941), he proved a handful of structural theorems that answer a question no amount of matrix multiplication answers by itself: given that an optical bench may contain an arbitrary number of waveplates, polarisers and rotators arranged in any order, **what is the most general thing such a bench can possibly do to a beam of light, and how few elements are needed to do it?**

The answers are remarkably restrictive, and that is exactly what makes them useful. They tell the instrument designer that certain constructions are impossible, that certain others are redundant,

and that any device whatsoever, however baroque, is equivalent to a short standard chain. Before stating them we need to fix what equivalence means and to classify the primitive elements by the kind of matrix they possess.

Definition 24 (*Optical equivalence*): Two optical systems are **optically equivalent** if their Jones matrices differ at most by a global phase factor, that is, if $J_2 = e^{i\gamma} J_1$ for some real γ . Equivalently, by 3.90, they possess the same Mueller matrix, and therefore no measurement of any kind performed on the outgoing light can distinguish them.

The three primitive devices built in the previous subsections fall into sharply distinct algebraic classes, and the whole of what follows is a consequence of that classification. A retarder 3.75 is **unitary** and, once the irrelevant global phase is removed, **unimodular**, $\det W = 1$; the reader may check this directly on the linear retarder 3.76, whose determinant is $\cos^2 \frac{\delta}{2} + \sin^2 \frac{\delta}{2} (\cos^2 2\alpha + \sin^2 2\alpha) = 1$. A rotator 3.83 is **real orthogonal** with unit determinant, hence a particular case of a retarder, as we already proved. A linear dichroic 3.85, and its limiting case the ideal polariser 3.86, is **real symmetric and positive semidefinite**, while a general elliptical dichroic 3.84 is **Hermitian and positive semidefinite**. Finally, a cascade is the ordered product of the matrices of its elements. Every theorem below is a statement about what such products can look like.

3.5.6.1 Theorem I: every system has an eigenpolarisation

Theorem 2 (*Existence of eigenpolarisations*): Every optical system with Jones matrix J possesses at least one polarisation state that it leaves unchanged in form, altering it at most in amplitude and phase. Generically there are exactly two such states, and they are mutually orthogonal if and only if J is a normal matrix, $JJ^\dagger = J^\dagger J$.

Proof. The eigenvalue equation $J\vec{e} = \lambda\vec{e}$ has a non trivial solution precisely when $\det(J - \lambda I) = 0$. Writing $J = \begin{bmatrix} j_{11} & j_{12} \\ j_{21} & j_{22} \end{bmatrix}$, this determinant is the quadratic

$$\lambda^2 - (j_{11} + j_{22})\lambda + (j_{11}j_{22} - j_{12}j_{21}) = \lambda^2 - (\text{tr } J)\lambda + \det J = 0, \quad (3.101)$$

a polynomial of degree two with complex coefficients. By the fundamental theorem of algebra it possesses at least one root $\lambda_1 \in \mathbb{C}$, and for that root the matrix $J - \lambda_1 I$ is singular, so its kernel contains a non zero vector $\vec{e}_1$. That vector is a Jones spinor 3.34 and therefore a genuine polarisation state, and by construction $J\vec{e}_1 = \lambda_1\vec{e}_1$: the outgoing spinor is the incoming one multiplied by a single complex number, whose modulus is an amplitude change and whose argument is a phase change, neither of which alters the shape or the orientation of the ellipse. When the discriminant of 3.101 does not vanish the two roots are distinct and the two eigenvectors are linearly independent, giving two eigenpolarisations.

For the second part, recall the spectral theorem: a complex square matrix admits an *orthonormal* basis of eigenvectors if and only if it is normal. Since orthogonality of the two eigenspinors in the

Hermitian product is precisely orthogonality of the two polarisation states, the equivalence follows at once. ■

The theorem has got more content than it first appears. All three primitive elements are normal: a retarder is unitary and a dichroic is Hermitian, and both classes satisfy $JJ^\dagger = J^\dagger J$ trivially. Their eigenstates are therefore always orthogonal, which is why we were able to build every one of them in the previous subsections from a pair of orthogonal states $|\psi_1\rangle, |\psi_2\rangle$ using a spectral decomposition. **A cascade, however, need not be normal.** Take a linear retarder of retardance $\delta = 1.2$ with its axis at $\alpha = 0.4$, followed by a linear dichroic with transmittances $t_1 = 0.9$ and $t_2 = 0.3$ along the coordinate axes. The commutator $[J, J^\dagger]$ of the product does not vanish, and the two eigenpolarisations turn out to satisfy $|\langle e_1 | e_2 \rangle| \approx 0.46$, which is very far from zero.

Elements whose eigenpolarisations are orthogonal are called **homogeneous**, and the others **inhomogeneous**. The lesson is worth stating plainly, because it is a common source of confusion: a waveplate followed by a polariser is not, in general, describable as some other waveplate followed by some other polariser with the same axes, and the reason is that the composite object has stopped being normal. The eigenstates it does possess are no longer a pair of orthogonal ellipses but a skew pair, and this is one of the ways in which real optical systems misbehave.

3.5.6.2 Theorem II: retarders and rotators

Theorem 3 (*First equivalence theorem*): *An optical system containing any number of retarders and rotators, in any order, is optically equivalent to a system containing only two elements: one linear retarder followed by one rotator.*

Proof. Every retarder and every rotator has got a unitary, unimodular Jones matrix, that is, an element of the group $SU(2)$. Since $SU(2)$ is a group it is closed under multiplication, so the whole cascade, whatever its length and order, is again a single element $U \in SU(2)$. The theorem therefore reduces to a purely algebraic claim: every $U \in SU(2)$ can be written as $U = R(\theta)W(\alpha, 0; \delta)$ for suitable θ, α, δ .

The cleanest way to see this is to move to the Poincaré sphere, where we already know what these objects do. By the theorem on retarders as rotations, the linear retarder $W(\alpha, 0; \delta)$ acts on the sphere as a rotation by δ about the equatorial axis 3.94 sitting at azimuth 2α , since $\chi = 0$ places $\hat{n}$ on the equator; and the rotator $R(\theta)$, being a circular retarder, acts as a rotation about the polar axis. So the claim becomes: *every rotation of the sphere is a rotation about an equatorial axis followed by a rotation about the polar axis.*

This is the Euler angle decomposition in disguise. Write the general rotation in the ZYZ convention as $\mathcal{R} = Z(\phi)Y(\vartheta)Z(\psi)$, which is possible for any element of $SO(3)$. Now insert the identity $Z(\psi)Z(-\psi)$ and regroup,

$$\mathcal{R} = Z(\phi)Y(\vartheta)Z(\psi) = \underbrace{Z(\phi + \psi)}_{\text{polar}} \underbrace{[Z(-\psi)Y(\vartheta)Z(\psi)]}_{\text{equatorial}},$$

and observe that the bracket is a rotation by the angle ϑ about the axis $Z(-\psi)\hat{y}$, which lies in the equatorial plane because $\hat{y}$ does and Z preserves that plane. The first factor is a rotation about the polar axis. Since ϕ, ϑ, ψ range over all of $SO(3)$, so do the three parameters θ, α, δ of our two element chain. Finally, the Mueller map 3.90 restricted to $SU(2)$ is onto $SO(3)$, and its only ambiguity is the sign of U , which is a global phase and hence optically invisible. The two systems are therefore optically equivalent. ■

The physical reading is the one that matters. **No matter how many waveplates you stack, at whatever angles, the result is a single elliptical retarder.** It can absorb nothing, because unitarity is preserved under multiplication, and it can depolarise nothing, by 3.98. Everything a pile of birefringent crystal can do is to rotate the Poincaré sphere once, and three numbers suffice to say how. This is why the elliptical retarder we constructed by a change of basis is not an academic curiosity but a piece of hardware: any (α, χ, δ) you like can be realised with one ordinary waveplate and one rotator, and in the laboratory the combination of two quarter wave plates and a half wave plate is the standard universal polarisation controller built on exactly this theorem.

3.5.6.3 Theorem III: partial polarisers and rotators

Theorem 4 (*Second equivalence theorem*): *An optical system containing any number of linear partial polarisers and rotators, in any order, is optically equivalent to a system containing only two elements: one linear partial polariser followed by one rotator.*

Proof. A linear partial polariser 3.85 has a *real symmetric* matrix, and a rotator 3.83 has a *real orthogonal* one. Products of real matrices are real, so the entire cascade is described by some real matrix A , non singular as long as no element is an ideal polariser. Furthermore $\det A > 0$, because every factor has positive determinant, namely $t_1 t_2 > 0$ for a partial polariser and $+1$ for a rotator.

We now invoke the real polar decomposition, and since it is the engine of both this theorem and the next, let us prove it rather than quote it. Consider the matrix AA^T . It is symmetric, since $(AA^T)^T = AA^T$, and it is positive definite, because for any non zero real vector $\vec{v}$ we have $\vec{v}^T AA^T \vec{v} = \|A^T \vec{v}\|^2 > 0$, the strict inequality holding because A^T is invertible. A real symmetric positive definite matrix can be diagonalised by an orthogonal matrix, $AA^T = V \text{diag}(\mu_1, \mu_2) V^T$ with $\mu_1, \mu_2 > 0$, so we may define

$$S \equiv V \text{diag}(\sqrt{\mu_1}, \sqrt{\mu_2}) V^T, \quad (3.102)$$

which is again real, symmetric, positive definite, invertible, and satisfies $S^2 = AA^T$. Setting $R \equiv S^{-1}A$ we verify directly that

$$RR^T = S^{-1}AA^TS^{-1} = S^{-1}S^2S^{-1} = I,$$

where we used that S^{-1} is symmetric. So R is real orthogonal, and $\det R = \det A / \det S > 0$ forces $\det R = +1$, making R a proper rotation, that is, a rotator $R(\theta)$. We have obtained $A = SR$ with S

a real symmetric positive definite matrix, which is exactly a linear partial polariser with some axis and some pair of transmittances, followed by a rotator. ■

Note where each hypothesis was used, because the theorem is sharp. Reality of the matrices is what forced the unitary factor to be a mere rotation instead of a general retarder; had we allowed elliptical partial polarisers, the argument would have produced a retarder in its place and the statement would fail. Physically this says something clean: **a stack of absorbing but non birefringent elements can never manufacture birefringence**, it can only attenuate anisotropically and turn the plane of polarisation. Dichroism and birefringence do not generate each other.

3.5.6.4 Theorem IV: the general equivalence theorem

Theorem 5 (*Third equivalence theorem*): Every non singular Jones matrix J factorises uniquely as

$$J = H U, \quad (3.103)$$

with H Hermitian positive definite and U unitary. Consequently any optical system whatsoever, containing any number of retarders, partial polarisers and rotators in any order, is optically equivalent to one elliptical partial polariser followed by one elliptical retarder, and therefore, expanding each of these through Theorem II, to a chain of four primitive elements: two retarders, one partial polariser and one rotator.

Proof. The construction copies the real case with daggers in place of transposes. Consider $JJ^\dagger$. It is Hermitian, since $(JJ^\dagger)^\dagger = JJ^\dagger$, and positive definite, because for any non zero $\vec{v} \in \mathbb{C}^2$,

$$\vec{v}^\dagger J J^\dagger \vec{v} = (J^\dagger \vec{v})^\dagger (J^\dagger \vec{v}) = \|J^\dagger \vec{v}\|^2 > 0,$$

strictly, because $J^\dagger$ is invertible and so cannot annihilate $\vec{v}$. By the spectral theorem a Hermitian matrix is diagonalisable by a unitary one, $JJ^\dagger = V \text{diag}(\mu_1, \mu_2) V^\dagger$ with $\mu_1, \mu_2 > 0$ real, and we define its positive square root

$$H \equiv V \text{diag}(\sqrt{\mu_1}, \sqrt{\mu_2}) V^\dagger, \quad (3.104)$$

which is Hermitian, positive definite, invertible and satisfies $H^2 = JJ^\dagger$. Put $U \equiv H^{-1}J$. Then, using $H^\dagger = H$ and hence $(H^{-1})^\dagger = H^{-1}$,

$$UU^\dagger = H^{-1}J J^\dagger H^{-1} = H^{-1}H^2H^{-1} = I,$$

so U is unitary and $J = HU$ as claimed. For uniqueness, suppose $J = H'U'$ with H' Hermitian positive definite and U' unitary. Then $JJ^\dagger = H'U'U'^\dagger H' = H'^2$, and since a positive definite Hermitian matrix has got exactly one positive definite Hermitian square root, $H' = H$, whence $U' = H^{-1}J = U$. ■

The dictionary back to hardware is immediate. The factor H , being Hermitian and positive definite, is by 3.84 an elliptical partial polariser, with its two eigenstates and its two real transmittances

$\sqrt{\mu_1}, \sqrt{\mu_2}$. The factor U , being unitary, is by 3.75 an elliptical retarder, which Theorem II lets us build from one linear retarder and one rotator. And H itself, an elliptical partial polariser, is a linear partial polariser conjugated by a retarder that carries the linear eigenstates onto the elliptical ones. Counting the free parameters confirms that nothing has been lost or invented: a general $J \in GL(2, \mathbb{C})$ has got eight real parameters, matched exactly by four for H plus four for U in 3.103, and equally by the hardware chain of two retarders at two parameters each, one partial polariser at three, and one rotator at one, totalling eight.

This is the deepest of the three theorems and it deserves a sentence of its own. **Every linear, non depolarising optical system in existence, however many elements it contains, does exactly two things and in this order: it attenuates the two components of the field unequally, and then it retards one relative to the other.** Diattenuation followed by birefringence, and nothing else is possible. In the language of the previous subsection the statement is even more striking, because a Hermitian positive H maps to a Lorentz boost and a unitary U to a rotation: 3.103 is nothing but the familiar decomposition of a Lorentz transformation into a boost times a rotation, transplanted to the Poincaré sphere. The result is not a historical curiosity either. Its descendant at the level of Mueller matrices, in which a measured M is factorised into a depolariser, a retarder and a diattenuator, is the standard tool by which a modern polarimeter turns sixteen raw numbers into three physically interpretable objects.

3.5.6.5 Theorem V: reversibility

The last theorem concerns what happens when the light is sent back through the system, and it is the one with the most conspicuous practical consequence.

Theorem 6 (Reversibility): *If a reciprocal optical system has Jones matrix J for propagation in one direction, then for propagation in the opposite direction its Jones matrix is the transpose J^T , the transverse axes being referred to the direction of travel.*

Proof. Reversing the beam does two things: it traverses the elements in the opposite order, and it traverses each individual element backwards. The first is automatic in the calculus, since if the forward system is $J = J_n \cdots J_2 J_1$ then the reversed one is $\tilde{J}_1 \tilde{J}_2 \cdots \tilde{J}_n$, where $\tilde{J}_k$ denotes the k th element traversed backwards. The identity $(J_n \cdots J_1)^T = J_1^T \cdots J_n^T$ reverses the order automatically, so the theorem holds for the whole system as soon as it holds for each element, $\tilde{J}_k = J_k^T$. It therefore suffices to check the primitives.

A linear retarder 3.76 is a **symmetric** matrix, its two off diagonal entries being equal to $i \sin \frac{\delta}{2} \sin 2\alpha$, so $W^T = W$: a waveplate looks the same from either side, which is exactly what a slab of birefringent crystal must do, since its fast and slow axes are properties of the crystal and not of the direction of travel. A linear dichroic 3.85 is likewise symmetric, and for the same reason. A rotator 3.83 is orthogonal, so $R(\theta)^T = R(\theta)^{-1} = R(-\theta)$: run backwards, an optically active medium rotates the plane of polarisation by the opposite angle. This is the correct physics, because in a sugar solution the sense of rotation is tied to the handedness of the molecules and to the direction of

propagation together, so reversing the beam reverses the apparent sense. Every primitive obeys the rule, and therefore so does every system built from them. ■

The consequences are best seen in a double pass, the situation of a beam that goes through a system, reflects off a mirror and returns. Ignoring the mirror itself, the round trip matrix is $J^T J$. For a quarter wave plate with its axis along x this gives, since W is symmetric,

$$W(0, 0; \frac{\pi}{2})^T W(0, 0; \frac{\pi}{2}) = W(0, 0; \frac{\pi}{2})^2 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} = W(0, 0; \pi), \quad (3.105)$$

precisely a half wave plate: **double passing a retarder doubles its retardance**, which is the everyday trick by which a quarter wave plate placed in front of a mirror rotates linear polarisation by ninety degrees and lets a polarising beam splitter separate the outgoing beam from the returning one. For a rotator, by contrast,

$$R(\theta)^T R(\theta) = R(-\theta)R(\theta) = I, \quad (3.106)$$

so **double passing an optically active medium undoes the rotation exactly**. Send polarised light through a tube of sugar solution, reflect it and send it back, and it emerges with its original azimuth.

And here is the payoff. A Faraday rotator, mentioned when we introduced the rotators, looks identical to an optically active medium on a single pass, yet its sense of rotation is fixed by the external magnetic field alone and not by the direction of propagation. On the return pass it therefore contributes $R(\theta)$ again rather than $R(-\theta)$, so the round trip gives $R(\theta)R(\theta) = R(2\theta)$ instead of the identity. The Faraday rotator **violates the reversibility theorem**, and it is allowed to do so precisely because a magnetic field breaks the time reversal symmetry on which reciprocity rests. This single fact is the entire working principle of the optical isolator, the device that lets light out of a laser but not back into it, without which no high power laser or fibre optic communication link would be stable. A theorem about transposes, proved in three lines, is what tells us that such a device cannot be built from waveplates and sugar and must be built from magnetised glass.

3.5.6.6 What the theorems say together

Standing back, the five results form a single picture. Theorem I says every system has a state it preserves, but warns that cascades of well behaved elements need not be well behaved themselves. Theorems II and III say that the two families of devices are closed and do not generate each other: birefringence stacks into birefringence and dichroism into dichroism. Theorem IV says that when the two families are mixed, the outcome is still astonishingly restricted, one diattenuator and one retarder and nothing more, eight real numbers in total. Theorem V says that this whole structure is blind to the direction of travel unless a magnetic field is present to break the symmetry.

The unifying statement is group theoretic, and it is worth making explicit because it will reappear whenever linear systems are studied. The non depolarising optical elements form the group $GL(2, \mathbb{C})$, of which the retarders form the unitary subgroup and the dichroics the positive Hermitian cone, and

3.103 is the statement that the group is the product of the two. Everything a bench can do to fully polarised light is an element of that group, and everything it can do to partially polarised light is, by the ensemble argument 3.99, a convex mixture of such elements. The Jones theorems are, in the end, the complete answer to the question of what polarisation optics is capable of.

3.5.7 The elliptic general Malus law

3.5.8 The law of elliptic birrefringents

Chapter 4

Interference and Interferometers

*When two waves meet, light may add to light
and yield darkness; in that paradox lies the
whole of physical optics.*

“When two undulations, from different origins, coincide either perfectly or very nearly in direction, their joint effect is a combination of the motions belonging to each.”

– Thomas Young

4.1 A historical context of interference and interferometers

Interference is the most direct fingerprint of the wave nature of light: when two beams overlap, their amplitudes - not their intensities - add, and the result can be brighter than either beam alone or, astonishingly, complete darkness. From this simple superposition principle grew the interferometer, an instrument that turns the invisible phase of a light wave into a visible pattern of fringes, and which became one of the most precise measuring devices ever built.

Although Robert Hooke and Isaac Newton both studied the coloured fringes produced by thin films - the phenomenon still known as Newton's rings, carefully described in Newton's *Opticks* of 1704 - neither interpreted them as evidence of waves. Newton's immense authority, and his corpuscular theory, held the field for a century. The decisive break came in 1801 with Thomas Young, whose double-slit experiment showed that light passing through two narrow apertures produces a pattern of bright and dark bands that can only be explained if the two beams alternately reinforce and cancel one another (Young, 1804). Young introduced the very word "interference" and, with it, the principle of superposition that underlies all of wave optics.

Young's qualitative insight was forged into a quantitative theory by Augustin-Jean Fresnel, who combined the principle of interference with Huygens's secondary wavelets to compute diffraction and interference patterns in full detail, and whose recognition that light is a transverse wave (developed with Arago) tied interference firmly to polarisation: only components sharing a polarisation can interfere. By the middle of the nineteenth century the wave theory was secure, and interference had become a tool. Hippolyte Fizeau and Léon Foucault used interferometric methods to compare the speed of light in different media, and Fizeau exploited the visibility of fringes to infer stellar properties - an idea far ahead of its time.

The instrument that defines the modern era is the Michelson interferometer. Albert A. Michelson, seeking ever more precise measurements, built in the 1880s a device that splits a beam in two, sends the halves along perpendicular arms, and recombines them so that any difference in optical path appears as a shift of the fringes. With it, Michelson and Edward Morley performed in 1887 their famous experiment to detect the motion of the Earth through the luminiferous aether; the null result (Michelson and Morley, 1887) became one of the experimental pillars on which Einstein's special relativity would later rest. Michelson also turned his interferometer into a metrological standard, measuring the length of the standard metre in wavelengths of light, work for which he received the Nobel Prize in 1907.

Almost simultaneously, Charles Fabry and Alfred Perot introduced in 1899 an interferometer of a different kind, based on multiple reflections between two highly reflecting parallel surfaces (Fabry and Perot, 1899). The Fabry-Perot interferometer produces extremely sharp transmission fringes and offers very high spectral resolving power; it is the direct ancestor of the optical resonators and cavities that define the modes of every laser. Other configurations followed - the Mach-Zehnder interferometer, with its separated paths ideal for probing transparent samples, and the Sagnac interferometer, sensitive to rotation and the heart of modern optical gyroscopes.

The visibility of the fringes in any of these instruments is not merely an experimental nuisance: it is a direct measure of the coherence of the light, and it was precisely the analysis of fringe visibility that

motivated the statistical theory of coherence developed in a later chapter. In the twentieth century, interferometry reached almost unimaginable sensitivity: the Laser Interferometer Gravitational-Wave Observatory (LIGO), a pair of kilometre-scale Michelson interferometers, detected in 2015 the passage of a gravitational wave by measuring a change in arm length thousands of times smaller than a proton Abbott et al. (2016). From Young's two slits to the detection of colliding black holes, the interferometer has remained the same idea - read the phase of a wave by letting it interfere with itself - refined to extraordinary perfection.

4.2 How does interference occur?

Lets consider two sources of electromagnetic field, under the conditions we've mentioned on this lectueres, we're going only to focus on the electric component, therefore, lets suppose we've got the following two sources of electric field

$$\vec{E}_1(\vec{r}, t) = \vec{E}_1(\vec{r})e^{-i\omega_1 t}, \quad (4.1)$$

$$\vec{E}_2(\vec{r}, t) = \vec{E}_2(\vec{r})e^{-i\omega_2 t}, \quad (4.2)$$

where the complex field amplitudes are given by

$$\vec{E}_1(\vec{r}) = E_1(\vec{r})e^{i\varphi_1(\vec{r})}\hat{e}_1, \quad (4.3)$$

$$\vec{E}_2(\vec{r}) = E_2(\vec{r})e^{i\varphi_2(\vec{r})}\hat{e}_2. \quad (4.4)$$

Theses complex field amplitudes are almost general, here the polarization vectores $\hat{e}_1, \hat{e}_2$ are'n't space-time dependent, which, in general, is what it's observed. The phases $\varphi_1(\vec{r}), \varphi_2(\vec{r})$ represent an arbitrary wavefront behavior, also $E_1(\vec{r}), E_2(\vec{r})$ represent arbitrary amplitudes. Thus the expressions above are general experessions for arbitrary sources of radiation.

The interference appears in the superposition of two or more electric fields, therefore, to study the interference of these two field in a space-time point $(\vec{r}, t)$ we're going to add these two as follows

$$\vec{E}(\vec{r}, t) = \vec{E}_1(\vec{r}, t) + \vec{E}_2(\vec{r}, t). \quad (4.5)$$

As we've explained before, measuring the electric field is really hard to do, so what we study and see are the irradiancies or radiations intensity, also the frequencies $\{\omega_1, \omega_2\}$ are too fast time oscillations for the human eye and most of the detectors, so we only can study their time averages. Having all this on count, lets calculate the average of the irradiances

$$\begin{aligned} \left\langle \left\| \vec{E}(\vec{r}, t) \right\|^2 \right\rangle &= \left\langle \left\| \vec{E}_1(\vec{r}, t) \right\|^2 \right\rangle + \left\langle \left\| \vec{E}_2(\vec{r}, t) \right\|^2 \right\rangle + 2 \left\langle \text{Re} \left\{ \vec{E}_1^*(\vec{r}, t) \cdot \vec{E}_2(\vec{r}, t) \right\} \right\rangle, \\ \left\langle \left\| \vec{E}(\vec{r}, t) \right\|^2 \right\rangle &= \left\| \vec{E}_1(\vec{r}) \right\|^2 + \left\| \vec{E}_2(\vec{r}) \right\|^2 + 2 \text{Re} \left\{ \vec{E}_1^*(\vec{r}) \cdot \vec{E}_2(\vec{r}) \left\langle e^{-i(\omega_1 - \omega_2)t} \right\rangle \right\}. \end{aligned}$$

The first two terms corresponds to **the total field intensity** and the third terms is called **the interference term** because it contains the interaction of both fields.

From a mathematical point of view, to make the interaction non zero, both frequencies must be the same $\omega_1 = \omega_2$, this, like most books say, should imply that the interference only occurs with two sources with the same frequency or on identical photons but, this is not completely correct. The frequency-difference exponential $\exp(-i[\omega_1 - \omega_2]t)$ is, in general, not zero, so **there are a lot ultrafast oscillations and interferences** what we see, and what we detected using measuring instruments is the time average, and in average, these are zero.

Before going further, lets define the field intensities as

$$\left\| \vec{E}_1(\vec{r}) \right\|^2 = I_1(\vec{r}), \quad (4.6)$$

$$\left\| \vec{E}_2(\vec{r}) \right\|^2 = I_2(\vec{r}). \quad (4.7)$$

The interference terms is then given by

$$\vec{E}_1^*(\vec{r}) \cdot \vec{E}_2(\vec{r}) = E_1(\vec{r})E_2(\vec{r})e^{i[\phi_2(\vec{r})-\phi_1(\vec{r})]}\hat{e}_1^* \cdot \hat{e}_2, \quad (4.8)$$

then the field intensity is

$$I = I_1 + I_2 + 2\text{Re} \left\{ (\hat{e}_1^* \cdot \hat{e}_2) e^{i\Delta\phi(\vec{r})} \right\} \underbrace{E_1(\vec{r})E_2(\vec{r})}_{(I_1 I_2)^{1/2}}. \quad (4.9)$$

The functional $\Delta\phi = \phi_2(\vec{r}) - \phi_1(\vec{r})$ is known as the phase-difference of the waves. The interference term is zero if the **polarisations are orthogonal** this means $\hat{e}_1^* \cdot \hat{e}_2 = 0$. Without any loss of generality an only for mathematical convenience¹ we're going to consider both sources have got the same polarisation, this means $\hat{e}_1 = \hat{e}_2 \rightarrow \hat{e}_1^* \cdot \hat{e}_1 = 1$. Thus, the total intensity becomes

$$I = I_1 + I_2 + 2(I_1 I_2)^{1/2} \cos \Delta\phi = (I_1 + I_2) \left(1 + 2 \frac{(I_1 I_2)^{1/2}}{I_1 + I_2} \cos \Delta\phi \right). \quad (4.10)$$

The equation 4.10 is the our most general expression for the interference. Lets notice that the interference terms is composed in two parts: the first one is a relation between the two intensities I_1, I_2 and the second one only depends of the phase-difference of the two sources by a cosine, then, **the real interference becomes in the phases not in the intensities**, the first terms only **modulates the visibility** of the interference, and we then can call it $\mathcal{V}$ and write it as

$$\mathcal{V} = 2 \frac{(I_1 I_2)^{1/2}}{I_1 + I_2} = \frac{I_{max} - I_{min}}{I_{max} + I_{min}}, \quad (4.11)$$

$$I_{min} = \left(I_1^{1/2} - I_2^{1/2} \right)^2, \quad I_{max} = \left(I_1^{1/2} + I_2^{1/2} \right)^2, \quad (4.12)$$

this term $\mathcal{V}$ is called the **visibility factor** which module the constrast on the interference fringes.

¹In fact, the polarisation here is only an contrast modulator, however, when we study the spatial coherence, the polarisation is even more important and conditionates the behavior of the interference.

4.2.1 Interference of plane waves

The expression 4.10 is, as we said, the general interference expression, so let's apply it to two plane wave sources. For the plane wave we've got that the complex amplitudes are given by

$$E_1(\vec{r}, t) = E_1 e^{i(\vec{r} \cdot \vec{k}_1 + \varphi_1)}, \quad E_2(\vec{r}, t) = E_2 e^{i(\vec{r} \cdot \vec{k}_2 + \varphi_2)}, \quad (4.13)$$

where, we are going to suppose, just for to make the maths a bit easier but the methodology is the same, that both amplitudes are the same $E_1 = E_2$.

Because, we are considering that the both waves have got the same frequency $\omega_1 = \omega_2$ this implies that both have the same wave number amplitude, this means $||\vec{k}_1|| = ||\vec{k}_2|| = k$. Then, we will have that the phase-difference term is given by

$$\Delta\phi = (\vec{k}_2 - \vec{k}_1) \cdot \vec{r} + \varphi_2 - \varphi_1. \quad (4.14)$$

The interference 4.10 having on count that $I_1 = I_2 = I_0$ is then

$$I = (I_0 + I_0) \left(1 + 2 \frac{(I_0 I_0)^{1/2}}{I_0 + I_0} \cos \Delta\phi \right) = 2I_0 \left[1 + \cos (\vec{K} \cdot \vec{r} + \Delta\varphi) \right], \quad (4.15)$$

where $\vec{K} = \vec{k}_2 - \vec{k}_1$ is the interferences vector. To understand better the interference, we're going to do a bit of geometry (see figure 4.1).

Because we're interested on the interference plane, we will take to $\vec{r} = X\hat{x}$ on the interference plane, where $\hat{x} = \vec{K}/||\vec{K}||$, then the spatial dependence gets the form

$$\vec{K} \cdot \vec{r} = KX = 2\kappa X \sin \left(\frac{\alpha}{2} \right), \quad (4.16)$$

and because K is a wave number vector on the interference plane, it can be written in terms of its wave-length, which corresponds to the interfringe distance Λ as

$$K = \frac{2\pi}{\Lambda}, \quad (4.17)$$

then

$$\vec{K} \cdot \vec{r} = \frac{2\pi}{\Lambda} X. \quad (4.18)$$

To calculate the relation between the interfringe distance Λ and the known parameters of the wave, we're going to equate the equations 4.16 and 4.18 therefore

$$\frac{2\pi}{\Lambda} X = \frac{2\pi}{\lambda} X \sin \left(\frac{\alpha}{2} \right), \quad (4.19)$$

$$\Lambda = \frac{\lambda}{2 \sin \left(\frac{\alpha}{2} \right)}, \quad (4.20)$$




Figure 4.1: Two plane waves with unitary number vector $\hat{k}_1, \hat{k}_2$ propagate and interfere producing a pattern of interference characterized by a vector $\vec{K}$ and an interfringe distance Λ .

where we used that $\kappa = \frac{2\pi}{\lambda}$. With this term, we finally got that the total intensity in the interference of two plane wave is given by

$$I = 2I_0 \left[1 + \cos \left(\frac{2\pi}{\Lambda} X + \phi \right) \right]. \quad (4.21)$$

We can see that the interfringe distance Λ behaves in a very logique way. In $\alpha \rightarrow 0$ then $\Lambda \rightarrow \infty$ this means, when the two wave number vector $\hat{k}_1, \hat{k}_2$ gets each time more parallel, the interfringe distance tends to infinity, this means, there won't be any interference pattern.

In the other way, if $\alpha \rightarrow \pi$ then $\Lambda \rightarrow \min$ this means, when the two wave number vectors tend to be 180° separated the interference distance is the minimum, being clear for us to see the interference pattern.

4.2.2 Interference of spherical waves

In a similar way to the last section about plane waves, we're going to consider two radiation sources which propagate as spherical waves, so the electric field waves are given by




Figure 4.2: Two plane sources of spherical waves interfere in a point P located on the interference plane. The two sources are separated a distance a and the distance from both surface to the interference plane is D .

$$E_1(\vec{r}) = \frac{E_1}{r_1} e^{i(kr_1 + \varphi_1)}, \quad (4.22)$$

$$E_2(\vec{r}) = \frac{E_2}{r_2} e^{i(kr_2 + \varphi_2)}. \quad (4.23)$$

And, again, we're going to consider, without any loss of generality just for aesthetical purposes, that both have got the same amplitude $E_1 = E_2 = E_0$. Also, let's suppose a point P on the interference plane and that the two point sources are separated a distance a [m] and both sources are separated a distance D [m] as is shown in the figure 4.2.

We have got that $r_1 = \overline{S_1P}$ and $r_2 = \overline{S_2P}$, then the phase difference is given by

$$\Delta\phi(r) = \kappa(r_2 - r_1) + \Delta\varphi. \quad (4.24)$$

And we are going to consider that the field amplitudes, in a neighborhood of P change very little

with r_1 and r_2 , and then

$$\frac{E_0}{r_1} \approx \frac{E_0}{r_2} = E, \quad (4.25)$$

the intensity is then given by

$$I = 2I_0 \{1 + \cos [\kappa (r_2 - r_1) + \Delta\varphi]\}, \quad (4.26)$$

before going further, see that the **surfaces of equal intensity** $\mathbf{r}_2 - \mathbf{r}_1 = cte$ **are hyperboloids with focii in $\mathcal{S}_1, \mathcal{S}_2$.**

In a plane parallel to $\overline{\mathcal{S}_1\mathcal{S}_2}$ at a distance $z = D$ two hyperboloids are produced. For $r_1 \approx r_2$ and $D \gg a$ this hiperboloids are practically planes. Therefore

$$r_1 = \left[D^2 + \left(x - \frac{a}{2} \right)^2 + y^2 \right]^{1/2} = D \left[1 + \frac{x^2}{D^2} + \frac{y^2}{D^2} + \frac{a^2}{4D^2} - \frac{ax}{D^2} \right]^{1/2}, \quad (4.27)$$

applying taylor series to first order of $\sqrt{1+x} \approx 1 + x/2$ we've got

$$r_1 \approx D \left[1 + \frac{x^2 + y^2}{2D^2} + \frac{a^2}{2D^2} - \frac{ax}{2D^2} \right], \quad (4.28)$$

and in the same way for r_2

$$r_2 \approx D \left[1 + \frac{x^2 + y^2}{2D^2} + \frac{a^2}{2D^2} + \frac{ax}{2D^2} \right]. \quad (4.29)$$

Substracting 4.29 from 4.27 we then we will have that the phase difference is then

$$\Delta\phi = \kappa (r_2 - r_1) + \Delta\varphi \approx \frac{\kappa a X}{D} + \Delta\varphi, \quad (4.30)$$

then the total intensity in the interference is given by

$$I = 2I_0 \left[1 + \cos \left(\kappa \frac{aX}{D} + \varphi_2 - \varphi_1 \right) \right], \quad (4.31)$$

where again

$$KX = \frac{\kappa a X}{D} = \frac{2\pi}{\lambda} \frac{aX}{D},$$

but $K = 2\pi/\lambda$, being again K the magnitude of the resultante wave number vector on the plane interference, and λ the interfringe distance. Thus the relation between λ an the known paramaters of both waves is

$$\frac{2\pi}{\lambda} = \frac{2\pi a X}{D} \longrightarrow \lambda = \frac{aD}{X}.$$

With this result, we can see that when the two sources get closer $a \rightarrow 0$ the interfringe $\Lambda \rightarrow \infty$ this means, the interference is going to disappear when the two point sources get together.

For example, lets consider two monochromatic plane waves, which have got complex amplitudes

$$\vec{E}_1 = \mathcal{E} e^{i\varphi_1} \hat{e}_1, \quad \vec{E}_2 = \mathcal{E} e^{i\varphi_2} \hat{e}_2, \quad (4.32)$$

where $\hat{e}_1, \hat{e}_2$ are their polarisation vectors and their wave number vectors are given by

$$\vec{k}_1 = [k_{1x}, k_{1y}, k_{1z}]^T \quad \vec{k}_2 = [k_{2x}, k_{2y}, k_{2z}]^T. \quad (4.33)$$

Now, lets answer the following questions:

1. What relation should be between $\vec{k}_1, \vec{k}_2$ to realize the interference?
2. What's the expression for the visibility?
3. What's the plane where the interference fringes have the least distance? What is the orientation of the plane and the interference fringes distances on that plane?

Lets answer each part following the methodology we have already built. The observable quantity is never the instantaneous field but the time average of the irradiance, which for our two waves reads

$$\left\langle \left\| \vec{E} \right\|^2 \right\rangle = I_1 + I_2 + 2 \operatorname{Re} \left\{ \vec{E}_1^* \cdot \vec{E}_2 \left\langle e^{-i(\omega_1 - \omega_2)t} \right\rangle \right\}. \quad (4.34)$$

The average $\left\langle e^{-i(\omega_1 - \omega_2)t} \right\rangle$ vanishes for any detector whose integration time T satisfies $|\omega_1 - \omega_2|T \gg 1$, so the interference term survives only when $\omega_1 = \omega_2 = \omega$. This first condition is **temporal, not geometric**, and the statement grants it by declaring both waves monochromatic and of the same nature. Its immediate translation to the space of wave number vectors, in a homogeneous and isotropic medium of index n , is

$$\left\| \vec{k}_1 \right\| = \left\| \vec{k}_2 \right\| = \kappa = \frac{n\omega}{c} = \frac{2\pi}{\lambda}. \quad (4.35)$$

This equality of the magnitudes is the only relation that the interference truly requires, and it does not constrain the directions at all. It is worth saying this out loud, because the wording of the question tempts us to answer that the vectors must be parallel, and it is exactly the opposite. If $\vec{k}_1 = \vec{k}_2$ the interference term does exist but no longer depends on the position, the illumination becomes uniform and there are no fringes. For an observable pattern we need the interference vector

$$\vec{K} \equiv \vec{k}_2 - \vec{k}_1 \neq \vec{0}, \quad (4.36)$$

whose magnitude follows at once from the equality of the moduli 4.35,

$$\left\| \vec{K} \right\|^2 = 2\kappa^2(1 - \cos \alpha) = 4\kappa^2 \sin^2 \frac{\alpha}{2} \implies K = 2\kappa \sin \frac{\alpha}{2}, \quad (4.37)$$

with α the angle between $\vec{k}_1$ and $\vec{k}_2$. A third condition, this time genuinely geometric, comes from the interference term carrying the factor $\hat{e}_1^* \cdot \hat{e}_2$, which vanishes for orthogonal polarisations. If we want the maximum contrast we need $\hat{e}_1 = \hat{e}_2 = \hat{e}$, and since both waves are transverse, so that $\hat{e} \perp \vec{k}_1$ and $\hat{e} \perp \vec{k}_2$, this common direction is fixed up to a sign,

$$\hat{e} = \frac{\vec{k}_1 \times \vec{k}_2}{\left\| \vec{k}_1 \times \vec{k}_2 \right\|}. \quad (4.38)$$

In words, two non collinear plane waves reach unit visibility only when they are polarised perpendicular to the plane that contains $\vec{k}_1$ and $\vec{k}_2$, which is the quantitative form of the Arago and Fresnel statement we met before.

For the second part, with $\omega_1 = \omega_2$ the interference term survives the average and the total irradiance is

$$I(\vec{r}) = I_1 + I_2 + 2\sqrt{I_1 I_2} |\hat{e}_1^* \cdot \hat{e}_2| \cos[\Delta\phi(\vec{r}) + \delta], \quad \Delta\phi(\vec{r}) = \vec{K} \cdot \vec{r} + \varphi_2 - \varphi_1, \quad (4.39)$$

where $I_j = |\mathcal{E}_j|^2$ and $\delta = \arg(\hat{e}_1^* \cdot \hat{e}_2)$. As the cosine sweeps the whole interval $[-1, 1]$ when $\vec{r}$ runs over the space, the extremes of the irradiance are read directly,

$$I_{\max} = I_1 + I_2 + 2\sqrt{I_1 I_2} |\hat{e}_1^* \cdot \hat{e}_2|, \quad I_{\min} = I_1 + I_2 - 2\sqrt{I_1 I_2} |\hat{e}_1^* \cdot \hat{e}_2|, \quad (4.40)$$

and the Michelson visibility, the same one we use as the contrast modulator, follows without any approximation

$$\mathcal{V} = \frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}} = \frac{2\sqrt{I_1 I_2}}{I_1 + I_2} |\hat{e}_1^* \cdot \hat{e}_2|. \quad (4.41)$$

With it the irradiance takes the canonical form

$$I(\vec{r}) = (I_1 + I_2) [1 + \mathcal{V} \cos \Delta\phi(\vec{r})], \quad (4.42)$$

which cleanly separates the three roles, since $I_1 + I_2$ fixes the mean level, $\mathcal{V}$ fixes the contrast and $\Delta\phi$ carries all the physics of the interference. Notice that $\mathcal{V} = 1$ demands both equal intensities and parallel polarisations, but also that $\mathcal{V}$ is remarkably robust against an imbalance of the intensities, since a ratio as large as $I_1/I_2 = 2$ still leaves $\mathcal{V} = 0.943$. We will make use of this insensitivity in the next example.

For the third part, the surfaces of constant phase, and therefore the fringes seen as objects living in the three dimensional space, are the planes $\vec{K} \cdot \vec{r} = \text{cte}$, perpendicular to $\vec{K}$ and separated by

$$\Lambda_0 = \frac{2\pi}{K} = \frac{\lambda}{2 \sin(\alpha/2)}. \quad (4.43)$$

What the question asks is on which observation plane P the cut of this family produces the closest fringes. Let $\hat{n}$ be the normal of P and decompose $\vec{K}$ into a part along $\hat{n}$ and a part $\vec{K}_{\parallel}$ lying

inside the plane. Only $\vec{K}_{\parallel}$ modulates the field over P , so the observed fringes are straight lines perpendicular to $\vec{K}_{\parallel}$ with a separation

$$\Lambda(\hat{n}) = \frac{2\pi}{\|\vec{K}_{\parallel}\|} = \frac{\Lambda_0}{\sin \gamma}, \quad \gamma = \angle(\vec{K}, \hat{n}). \quad (4.44)$$

Since $\Lambda(\hat{n}) \geq \Lambda_0$ for every orientation, with equality only when $\sin \gamma = 1$, the fringes are closest exactly when the observation plane contains the direction of $\vec{K}$,

$$P : \quad \hat{n} \cdot \vec{K} = 0, \quad \Lambda_{\min} = \frac{2\pi}{\|\vec{k}_2 - \vec{k}_1\|} = \frac{\lambda}{2 \sin(\alpha/2)}, \quad (4.45)$$

while any other tilt dilates the fringes by the factor $1/\sin \gamma$. This is a whole one parameter family of planes rather than a single one, and two of its members deserve a name. The first is the plane spanned by $\vec{k}_1$ and $\vec{k}_2$, whose normal is $\hat{n} = (\vec{k}_1 \times \vec{k}_2)/\|\vec{k}_1 \times \vec{k}_2\|$, and it contains $\vec{K}$ because $\vec{K}$ is a linear combination of $\vec{k}_1$ and $\vec{k}_2$, being moreover the same plane to which the polarisation had to be perpendicular in the first part. The second is the plane perpendicular to the bisector of the two beams, which is the one used as a screen in the laboratory, and it belongs to the optimal family thanks to the identity

$$\vec{K} \cdot (\vec{k}_1 + \vec{k}_2) = \|\vec{k}_2\|^2 - \|\vec{k}_1\|^2 = 0. \quad (4.46)$$

So the interference vector is always perpendicular to the bisector, the natural screen is already the optimal one and there is no need to tilt it. On it the fringes are straight lines parallel to $\vec{k}_1 + \vec{k}_2$ with the separation $\lambda/[2 \sin(\alpha/2)]$.

4.3 Interferometers

4.3.1 Young interferometer

For most of the eighteenth century the nature of light seemed a settled matter. The authority of Newton, whose *Opticks* Newton (1704) accounted for reflection, refraction and the colours of thin films in terms of tiny corpuscles, had pushed into the background the wave picture that Huygens had proposed in his *Traité de la Lumière* Huygens (1690). A corpuscle travels in a straight line and carries momentum, so the rectilinear propagation of light and the sharpness of shadows looked like immediate consequences of the theory, and the enormous prestige of Newton did the rest. The wave idea survived only as a minority opinion, lacking the one thing that would have made it irresistible, a phenomenon that particles simply cannot imitate.

That phenomenon was interference, and it was Thomas Young who brought it to light. In his Bakerian lecture of 1801, published as *On the Theory of Light and Colours* Young (1802), Young stated what he called the principle of the superposition of undulations, the plain but revolutionary idea that where two portions of light meet their disturbances add, so that two bright beams can

combine to give darkness. A few years later, in a second Bakerian lecture Young (1804), he turned the principle into an experiment, letting light pass through a small opening and then splitting the emerging beam so that its two halves overlapped on a screen, where they painted a regular pattern of bright and dark bands. In his *Course of Lectures on Natural Philosophy* Young (1807) the arrangement acquired the form we still teach, two closely spaced apertures illuminated by a common source, and from the spacing of the fringes Young was able to do something no one had done before, to measure the wavelength of light and to find it astonishingly small, a fraction of a thousandth of a millimetre.

The importance of this result is difficult to overstate. The dark bands were the direct proof that light added to light can produce less light, something no theory of independent particles can explain, and they gave the wave theory the empirical victory it had lacked for a century. The reception, however, was hostile, since to attack Newton in the England of 1804 was close to heresy, and Young's papers were ridiculed in the reviews of the day. Vindication came from the continent a decade later, when Fresnel and Arago placed the wave theory on a rigorous mathematical footing and extended it to diffraction and polarisation. The experiment itself never left the stage. It is the archetype of every wavefront division interferometer, the simplest device in which the phase difference between two coherent beams is turned into a visible intensity pattern, and its complete modern treatment belongs to any serious optics text Born and Wolf (1999). More than that, when the same experiment is repeated with electrons, atoms or single photons sent one at a time, the fringes still build up, and the two slit arrangement becomes the cleanest illustration of quantum superposition, the phenomenon that Feynman called the only mystery of quantum mechanics Feynman et al. (1965). The precision descendants of Young's screen reach today the kilometre sized interferometers that detect gravitational waves Abbott et al. (2016).

To see the whole machinery of the previous section at work on this instrument, let's analyse the Young interferometer quantitatively, with its two slits illuminated by a spherical wave coming from a point source, arranged as in the figure 4.3. We place the two slits $\mathcal{S}_1$ and $\mathcal{S}_2$ symmetrically at $x = \pm d/2$ on the plane $z = 0$, separated by a distance d , and the observation screen at $z = D$. The point source $\mathcal{S}$ sits at $(x, z) = (-a, -b)$, so that b is its distance to the slit plane and a its transverse displacement from the symmetry axis, with $a = 0$ meaning the source on the axis, and we measure the coordinate x on the screen from that axis.

The methodology of the chapter applies in two chained stages, and this is really the only structural idea of the problem. The first stage goes from the source to the slits and decides two things, namely with what amplitude and with what phase the field reaches each slit. The second stage goes from the slits to the screen and is always the interference of two secondary spherical waves, exactly the one we treated in the spherical wave section, with the slits acting as the sources $\mathcal{S}_1, \mathcal{S}_2$ separated by d and the screen at a distance D . Writing the field that leaves the slit j as $\mathcal{E}_j e^{i\psi_j}$ and the path from that slit to the point P on the screen as s_j , the total relative phase at P is

$$\Delta\phi(x) = \kappa(s_2 - s_1) + (\psi_2 - \psi_1), \quad (4.47)$$

and the irradiance obeys the canonical expression 4.42 with the visibility 4.41 evaluated on the amplitudes that reach each slit.




Figure 4.3: Young interferometer illuminated by a spherical wave from a point source $\mathcal{S}$ placed off axis. The source to slit stage sets the amplitude and the phase reaching each slit $\mathcal{S}_1, \mathcal{S}_2$, and the slit to screen stage is the interference of two secondary spherical waves.

The second stage is common to any illumination, so we dispatch it once. With $s_j = [D^2 + (x \mp d/2)^2]^{1/2}$ and the same Taylor expansion of the chapter,

$$s_2 - s_1 \approx \frac{(x + d/2)^2 - (x - d/2)^2}{2D} = \frac{xd}{D}, \quad (4.48)$$

valid in the paraxial regime $D \gg d, |x|$, from which

$$\kappa(s_2 - s_1) \approx \frac{2\pi}{\lambda} \frac{xd}{D} = \frac{2\pi}{\Lambda} x, \quad \Lambda = \frac{\lambda D}{d}. \quad (4.49)$$

This is precisely the interfringe of the spherical wave section with $a \rightarrow d$. Everything that distinguishes one illumination from another sits, then, in the term $\psi_2 - \psi_1$.

For the spherical illumination of this example the source at $(-a, -b)$ emits the wave $E = (E_0/r)e^{i\kappa r}$ of the chapter, and now the two slits are no longer equivalent, because their distances to the source

$$r_1 = \left[b^2 + \left(a + \frac{d}{2} \right)^2 \right]^{1/2}, \quad r_2 = \left[b^2 + \left(a - \frac{d}{2} \right)^2 \right]^{1/2} \quad (4.50)$$

differ as soon as $a \neq 0$, and this affects both the phase and the amplitude. In phase, expanding as in the chapter with $b \gg d, a$,

$$r_1 - r_2 \approx \frac{(a + d/2)^2 - (a - d/2)^2}{2b} = \frac{ad}{b} \implies \psi_2 - \psi_1 = \kappa(r_2 - r_1) = -\frac{\kappa ad}{b}. \quad (4.51)$$

In amplitude, the slit j receives E_0/r_j , so $I_j \propto 1/r_j^2$ and the visibility 4.41 becomes

$$\mathcal{V} = \frac{2\sqrt{I_1 I_2}}{I_1 + I_2} = \frac{2/(r_1 r_2)}{1/r_1^2 + 1/r_2^2} = \frac{2r_1 r_2}{r_1^2 + r_2^2}. \quad (4.52)$$

Putting the two stages together in 4.47, the total phase is

$$\Delta\phi(x) = \frac{\kappa d}{D}x - \frac{\kappa a d}{b} = \frac{2\pi d}{\lambda} \left(\frac{x}{D} - \frac{a}{b} \right), \quad (4.53)$$

and the irradiance on the screen reads

$$I(x) = (I_1 + I_2) \left\{ 1 + \mathcal{V} \cos \left[\frac{2\pi d}{\lambda} \left(\frac{x}{D} - \frac{a}{b} \right) \right] \right\}, \quad \mathcal{V} = \frac{2r_1 r_2}{r_1^2 + r_2^2}. \quad (4.54)$$

The qualitative difference with a plane illumination, for which $\mathcal{V} = 1$ always, is not in the position of the fringes but in the contrast, because an off axis spherical source gives $\mathcal{V} < 1$.

The first part asks for the value of a that reduces the contrast by 10% with respect to $a = 0$. At $a = 0$ we have $r_1 = r_2$ and $\mathcal{V} = 1$, and we ask for $\mathcal{V}(a) = 0.9$. Writing $t = r_1/r_2 \geq 1$, the condition $2t/(1+t^2) = 9/10$ is the quadratic $9t^2 - 20t + 9 = 0$, whose larger root is

$$t = \frac{10 + \sqrt{19}}{9} = 1.5954 \implies \frac{r_1^2}{r_2^2} = 2.5454. \quad (4.55)$$

A contrast drop of only 10% already requires one slit to be some 60% farther from the source than the other, which shows how insensitive the visibility is to an amplitude imbalance, exactly as we anticipated in the second part of the first example. Substituting the distances 4.50,

$$\frac{b^2 + (a + d/2)^2}{b^2 + (a - d/2)^2} = t^2 \iff a^2 - \frac{10}{\sqrt{19}}da + b^2 + \frac{d^2}{4} = 0, \quad (4.56)$$

a quadratic in a whose formal solution is

$$a = \frac{5d}{\sqrt{19}} \pm \sqrt{\frac{81d^2}{76} - b^2}. \quad (4.57)$$

Here the physics jumps out of the discriminant, since real solutions exist only if $b \leq \frac{9d}{2\sqrt{19}} = 1.0324d$. In a real Young interferometer one always has $b \gg d$, typically $d \sim 10^{-4}$ m and $b \sim 10^{-1}$ m, so the discriminant is negative and no displacement of a point source can degrade the contrast by 10%. The paraxial estimate confirms it, since with $t = 1 + \eta$ and $\eta \approx (r_1 - r_2)/b \approx ad/b^2$ the visibility is $\mathcal{V} \approx 1 - \eta^2/2 = 1 - \frac{1}{2}(ad/b^2)^2$, so $\mathcal{V} = 0.9$ would need $ad/b^2 = \sqrt{0.2} \approx 0.447$, that is $a \approx 0.45b^2/d$, a gigantic displacement that destroys the paraxial approximation long before it touches the contrast. The physical conclusion is that a strictly point source always produces fringes of essentially unit visibility, wherever it is placed, because moving it translates the pattern but does not erase it. The version of the question that does have a solution is the one where a is the width of an incoherent thermal source instead of the displacement of a point source, which belongs to the study of spatial

coherence and gives a visibility $\mathcal{V}(a) = |\text{sinc}(\pi ad/\lambda b)|$ that falls to its first zero at $a = \lambda b/d$. The lesson to keep is the contrast between the two readings, since displacing a point source acts on the phase and slides the fringes, while broadening an incoherent source acts on the coherence and washes them out.

The second part asks for the position x_m of the m -th maximum. The maxima of the irradiance 4.54 are the zeros modulo 2π of the phase 4.53,

$$\frac{2\pi d}{\lambda} \left(\frac{x_m}{D} - \frac{a}{b} \right) = 2\pi m, \quad m \in \mathbb{Z}, \quad (4.58)$$

from where

$$x_m = m \frac{\lambda D}{d} + \frac{aD}{b} = m\Lambda + \frac{aD}{b}. \quad (4.59)$$

The pattern keeps its full periodicity $\Lambda = \lambda D/d$ and is shifted as a rigid block by aD/b toward the side opposite to the displacement of the source, consistent with the figure, where the source lies below the axis while x_m is measured above. Geometrically, the zero order maximum sits where the total path difference vanishes, that is on the straight line that joins the source with the midpoint of the slits and continues to the screen, a line that leaves $(-a, -b)$, passes through the origin and reaches the screen at $x = aD/b$.

The third part asks for a condition between a and b so that the origin is a minimum. A minimum at $x = 0$ requires the total phase there to be an odd multiple of π . Evaluating 4.53 at $x = 0$,

$$\Delta\phi(0) = -\frac{2\pi d}{\lambda} \frac{a}{b} = -(2m+1)\pi, \quad m = 0, 1, 2, \dots \quad (4.60)$$

which is the same as demanding that the path difference introduced by the source to slits stage be an odd number of half wavelengths, $ad/b = (2m+1)\lambda/2$, that is

$$\frac{a}{b} = \frac{(2m+1)\lambda}{2d}, \quad m = 0, 1, 2, \dots \quad (4.61)$$

The condition depends on a and b only through their ratio, as it must, since the only thing a point source communicates to the system is the direction from which it illuminates it. The smallest case $m = 0$ gives $a/b = \lambda/2d$, which for typical numbers, $\lambda = 633 \text{ nm}$ and $d = 0.2 \text{ mm}$, is $a/b \approx 1.6 \times 10^{-3}$, so a displacement of about 1.6 mm for every metre of b is already enough to turn the central maximum into a minimum.

4.3.2 Fresnel biprism

The double mirror that Fresnel used to prove the wave nature of light Fresnel (1819) has one practical drawback, namely that two separate flat mirrors have to be tilted and held at an almost invisible angle between them, which makes the alignment tedious and unstable. Fresnel's own answer to this difficulty was to replace the two mirrors by a single piece of glass ground into two thin prisms joined base to base, with the shared edge running along the optical axis. This is the *Fresnel biprism*, and it

produces exactly the same effect, two coherent virtual images of a single point source, but out of one rigid element instead of two separately adjustable ones.

Before setting up the biprism itself, it is worth pausing on the principle that makes this whole family of instruments work, since it is the same principle that will let us handle every wavefront division interferometer to come. When a wave leaves a real point source and passes through an optical element, be it a lens, a mirror or a prism, its wavefront is in general reshaped into something far too complicated to describe by hand. There is, however, a special situation in which nothing of interest is lost, namely when the element is *stigmatic* for the pair of points involved, in the sense already introduced in the previous chapter (see figure 2.12), a spherical wavefront leaving the object point is transformed into another spherical wavefront, indistinguishable in its geometry from the one that a genuine point source sitting at the image point would radiate. Whenever this holds, Fermat's principle guarantees that every ray reaching the image has travelled the same optical path, so the phase downstream of the element is exactly the phase of a fictitious spherical wave centred on that image, whether the image is real or virtual. This is the *method of images*: rather than propagating the true, complicated wavefront through the optical element, we replace the element by the point sources it images and let the machinery of two point source interference, already built in the previous section, take care of everything past that plane.

This single observation is what turns an ordinary mirror or prism into an interferometer. If one primary source is imaged twice, once through each of two branches of the same optical element, the observer downstream no longer sees one wave but two, radiated by two virtual sources $\mathcal{S}_1$ and $\mathcal{S}_2$. These two sources are coherent by construction rather than by assumption, since both wavefronts are literally two halves of the same original wave, carrying the same frequency and a phase relationship fixed once and for all by the geometry of the element; the condition $\omega_1 = \omega_2$ that opened this chapter is here automatically satisfied and never has to be imposed by hand, unlike the case of two genuinely independent sources. Once the positions of $\mathcal{S}_1$ and $\mathcal{S}_2$ are known, the problem stops being about mirrors or prisms altogether and becomes, exactly, the interference of two spherical waves solved earlier in this chapter, with the separation a and the source to screen distance D read directly off the geometry of the particular device. In this sense every wavefront division interferometer, the double mirror, the biprism, and the others we will meet, differs from the next only in how it manufactures its pair of virtual sources; the fringes that follow are always the same fringes.

The method is only as trustworthy as the stigmatism of the imaging step, and that has to be checked separately for every device, since it is never automatic. A plane mirror is rigorously and unconditionally stigmatic for every object point whatsoever, as we saw when it appeared as the degenerate conic $z_o = -z_i$ of the reflective surfaces 2.76, which is why Fresnel's original double mirror needs no further justification beyond the law of reflection. A prism is a different matter, because among refracting interfaces only the very special Cartesian ovoids are rigorously stigmatic, and an ordinary prism face is not one of them; it can only be stigmatic approximately, for a narrow paraxial bundle of rays clustered near a fixed angle of incidence. Showing that this approximate stigmatism holds well enough for a thin prism, and pinning down exactly how well, is therefore the first task in developing the biprism, and it is where we begin.

Let a point source $\mathcal{S}$ sit at a distance z_1 from the biprism, and let the screen of observation stand

a further distance z_2 beyond it, so that $D = z_1 + z_2$ is the total distance from the source to the screen (see figure 4.4). Each half of the biprism is a thin prism of apex angle α and refractive index n , and the whole instrument is symmetric about the optical axis. The method of images consists of replacing the biprism entirely by the two virtual sources it creates, which turns the problem back into the two point source interference we already solved.




Figure 4.4: Method of images applied to the Fresnel biprism. Each half of the biprism produces a virtual point image of the source, $\mathcal{S}_1$ from the upper half and $\mathcal{S}_2$ from the lower half, separated by a and located in the same transverse plane as $\mathcal{S}$. Beyond the biprism the two beams overlap only inside the shaded strip of width W on the screen.

A thin prism deviates a ray by an angle that, to first order in the angle of incidence and near the condition of minimum deviation, does not depend on exactly where the ray crosses the prism nor, to that same order, on its precise angle of incidence,

$$\delta = (n - 1)\alpha. \quad (4.62)$$

This independence is the whole reason the method of images works here, because if the deviation changed from ray to ray the upper half of the biprism would not produce a point image but an aberrated smear. It is worth checking how good 4.62 really is, since later we will be pushed

toward angles that are not so small. Comparing it with the exact deviation at symmetric incidence, $\delta_{\text{exact}} = 2 \arcsin [n \sin(\alpha/2)] - \alpha$, for $n = 1.5$ the relative error is only 0.04% at $\alpha = 0.05$ rad, 0.16% at $\alpha = 0.10$ rad, 0.35% at $\alpha = 0.15$ rad and 0.63% at $\alpha = 0.20$ rad, so the thin prism approximation survives well into the range we will need, though it stops being free.

Consider a ray leaving $\mathcal{S}$ with paraxial slope m and crossing the upper half of the biprism. It strikes the prism plane at a height $y_p = z_1 m$ and leaves with the new slope $m - \delta$, deviated toward the axis. Extending this outgoing ray backward by the same distance z_1 to the plane of the source, its apparent point of origin is at a height

$$y_v = y_p - z_1(m - \delta) = z_1 m - z_1 m + z_1 \delta = z_1 \delta, \quad (4.63)$$

which does not depend on m . Every ray that the upper half deviates, whatever its original slope, appears to come from the same point, so the upper half of the biprism produces a rigorously stigmatic virtual image $\mathcal{S}_1$ at a height $z_1 \delta$ above the axis, in the very same transverse plane as the source. By the mirror symmetry of the instrument the lower half produces $\mathcal{S}_2$ at $-z_1 \delta$. The biprism has disappeared from the problem, replaced by two mutually coherent point sources separated by

$$a = 2z_1 \delta = 2z_1(n - 1)\alpha, \quad (4.64)$$

sitting at a distance $D = z_1 + z_2$ from the screen. From here on the problem is precisely the interference of two spherical waves that we already solved, so the irradiance on the screen, with y the transverse coordinate, takes the canonical form

$$I(y) = (I_1 + I_2) \left[1 + \mathcal{V} \cos \left(\kappa \frac{a}{D} y \right) \right], \quad (4.65)$$

with a fringe spacing given directly by the two point source result

$$\Lambda = \frac{\lambda D}{a} = \frac{\lambda(z_1 + z_2)}{2z_1(n - 1)\alpha}. \quad (4.66)$$

It helps to read 4.66 in the form that makes the underlying physics transparent, writing $\Lambda = \lambda/\beta$ with

$$\beta = \frac{a}{D} = 2(n - 1)\alpha \frac{z_1}{z_1 + z_2}, \quad (4.67)$$

the local angle at which the two beams cross on the screen. This is exactly the same quantity that appeared in the interference of two plane waves, where $\Lambda = \lambda/[2 \sin(\beta/2)]$, and the two expressions differ by only 0.12% at $\beta = 0.1$ rad, so identifying $\Lambda \approx \lambda/\beta$ here is entirely legitimate. What 4.67 makes plain is that

$$\beta \leq 2(n - 1)\alpha, \quad (4.68)$$

with equality only in the limits $z_2 \rightarrow 0$, the screen pressed against the biprism, or $z_1 \rightarrow \infty$, a collimated beam. The crossing angle can never exceed the full deviation the biprism is able to imprint on a beam, and the divergence of the spherical wave leaving a finite z_1 can only make it smaller. A biprism illuminated by a point source is, in this sense, always a weaker interferometer than the same biprism illuminated by a plane wave.

As a concrete example, take a biprism of index $n = 1.5$ illuminated by light of wavelength $\lambda = 632 \text{ nm}$, and ask for the geometry that produces a fringe spacing $\Lambda = 10\lambda$. Setting $\Lambda = 10\lambda$ in 4.66 means $\beta = 1/10$, and since $n = 1.5$ gives $2(n - 1) = 1$ exactly, the condition collapses to something remarkably clean,

$$\alpha \frac{z_1}{z_1 + z_2} = \frac{1}{10} \text{ rad} \iff \alpha = \frac{1}{10} \left(1 + \frac{z_2}{z_1} \right) \text{ rad}. \quad (4.69)$$

This is not a single value of α but a whole one parameter family, since fixing the ratio z_2/z_1 fixes α and vice versa, and every member of the family shares the same image separation $a = D/10$, a tenth of the total source to screen distance, which is a strikingly large separation for two virtual images meant to interfere. There is a bound hiding in 4.69 that is easy to miss, because $z_2/z_1 \geq 0$ forces

$$\alpha \geq \frac{1}{10} \text{ rad} = 5.73^\circ, \quad (4.70)$$

with the minimum reached only in the collimated limit. A prism angle of nearly six degrees is not what one usually pictures under the instruction to keep α small, and this is worth stating openly rather than hiding it, because the two requirements pull in opposite directions. A fringe spacing of $10\lambda = 6.32 \mu\text{m}$ is an extraordinarily fine spacing to demand, comparable to the period of a diffraction grating, and producing it requires the two beams to cross at $\beta = 0.1 \text{ rad}$, a full tenth of a radian, while a biprism can only ever supply a crossing angle of $2(n - 1)\alpha$. The solution exists and is entirely consistent, but it sits at the edge of the thin prism regime rather than deep inside it, and the error check performed above shows that the approximation 4.62 still holds at $\alpha = 0.1 \text{ rad}$, but by a margin of only 0.16% rather than by a comfortable one. It is also worth noticing that λ never appears in the condition on α alone, which makes sense once we remember that Λ was specified in units of λ and the pure geometry of the biprism carries no memory of colour.

A condition on α is not yet the whole answer, because it says nothing about whether a useful pattern is actually observable. The two beams emerging from the biprism only overlap inside a finite strip of the screen, since the ray that grazes the shared edge from above is deviated downward by δ and reaches the screen at a height $-z_2\delta$, while the ray grazing the edge from below is deviated upward and reaches $+z_2\delta$. Outside this interval only one of the two beams arrives and there is no interference at all, so the field of overlap has a width

$$W = 2z_2\delta = 2z_2(n - 1)\alpha, \quad (4.71)$$

and the number of fringes that fit inside it follows at once,

$$N = \frac{W}{\Lambda} = \frac{z_2(n - 1)\alpha}{5\lambda}. \quad (4.72)$$

Evaluated on the design condition 4.69, with $2(n - 1) = 1$ for $n = 1.5$, this becomes $N = z_2(z_1 + z_2)/(100 z_1 \lambda)$, and it is instructive to see it on real numbers.

z_1	z_2	α	a	N
1 m	1 cm	0.101 rad = 5.79°	10.1 cm	160
0.5 m	5 mm	0.101 rad = 5.79°	5.05 cm	80
∞ (collimated)	1 cm	0.100 rad = 5.73°	—	158

With the screen a centimetre from the biprism and the source a metre away, the design delivers 160 fringes of $6.32 \mu\text{m}$ inside a field of overlap only about a millimetre wide, so resolving them in practice needs a microscope objective or a detector with sub micron pixels, a condition that deserves to be stated explicitly rather than left implicit in the numbers. The visibility over this whole field stays extremely close to unity, since on the axis of symmetry the two virtual sources are exactly equidistant from the observation point and $\mathcal{V} = 1$ exactly, and even away from the axis the two distances remain equal up to second order corrections, so the same robustness of the contrast against small imbalances that we established for the Young interferometer is at work here, only now it holds by the exact symmetry of the construction rather than by an accident of the numbers.

It remains to see what happens when the point source is displaced off the optical axis, since a real source is never perfectly centred. Repeat the same construction with the source at $\mathcal{S} = (-z_1, s)$, a small transverse offset s away from the axis. A ray leaving $\mathcal{S}$ with slope m strikes the upper half of the biprism at height $y_p = s + z_1 m$ and leaves with slope $m - \delta$, so its virtual origin is now at

$$y_v = s + z_1 m - z_1(m - \delta) = s + z_1 \delta, \quad (4.73)$$

once again independent of m , so the images remain perfectly stigmatic under decentring. The two virtual sources simply follow the displacement,

$$\mathcal{S}_1 = s + z_1 \delta, \quad \mathcal{S}_2 = s - z_1 \delta, \quad (4.74)$$

and three consequences follow from 4.74 at once.

The fringe spacing does not change. The separation is still $a = 2z_1 \delta$ and the distance to the screen is still $D = z_1 + z_2$, both untouched by s , so $\Lambda = \lambda D/a$ is invariant to first order in the displacement. It is invariant to second order as well, and for a reason worth stating rather than taking for granted, since the deviation of a prism is stationary at the condition of minimum deviation, $d\delta/di = 0$ there, so tilting the incident beam slightly changes δ only at order $(\Delta i)^2$. The biprism is therefore intrinsically robust against a decentred source, more so than the Young interferometer with its fixed slits, where no such stationarity protects the spacing.

The whole fringe system translates rigidly by exactly s . Both virtual images move together with the source, so the plane of zero path difference, the perpendicular bisector of $\mathcal{S}_1 \mathcal{S}_2$, is simply the plane $y = s$, and the zero order fringe sits there regardless of z_2 . There is no geometric amplification of the kind we found for the Young interferometer with fixed slits, where the shift on the screen was sD/b rather than s itself, and the difference is that there the source and its two images move independently of the fixed slits, while here the source and both of its images move together as a rigid unit.

The field of overlap translates too, but in the opposite direction and by a different amount, $-z_2 s/z_1$. The ray grazing the shared edge from above now leaves $\mathcal{S}$ with slope $-s/z_1$ and acquires the extra deviation $\mp \delta$, reaching the screen at $y = -z_2 s/z_1 \mp z_2 \delta$, so the window keeps its width $W = 2z_2 \delta$ but is now centred at $-z_2 s/z_1$. Since the fringes and the window move in opposite directions, the central fringe eventually walks out of the observable field once

$$|s| > \frac{z_1 z_2 \delta}{z_1 + z_2} = \frac{z_2 \beta}{2}. \quad (4.75)$$

With the numbers of the table above, $z_1 = 1$ m, $z_2 = 1$ cm and $\beta = 0.1$, this bound is $|s| < 0.5$ mm, a generous tolerance for the alignment of the source, yet a displacement of only $s = 6.3 \mu\text{m}$ is already enough to slide the whole pattern by one complete fringe. The Fresnel biprism is therefore, at the same time, an interferometer remarkably tolerant of a coarse misalignment of the source, since the fringe spacing survives it exactly, and an extremely sensitive detector of a fine transverse displacement, since the fringe position tracks that displacement one to one, a duality that turns out to be useful whenever the instrument is repurposed as a position sensor rather than as a simple demonstration of interference.

4.3.3 Michelson and Morley interferometer




Figure 4.5: General configuration of the Michelson interferometer. A source placed a distance a from the beamsplitting cube illuminates it, and the beam is divided along two perpendicular arms. In arm 1 the light travels a total distance $d + \ell_1$ before striking mirror 1, tilted an angle θ away from the orientation that would send the beam straight back along itself; in arm 2 it travels a total distance $d + \ell_2$ before striking mirror 2, tilted an angle ϕ in the same sense. The two reflected beams return to the beamsplitter and recombine there, ready to interfere.

The configuration sketched above did not spring fully formed from the 1887 experiment already described at the opening of this chapter. Michelson built and tested a first, considerably simpler version of it in 1881, while on leave from the United States Naval Academy and working in Hermann von Helmholtz's laboratory in Berlin Michelson (1881). That prototype was mounted directly on a stone pier in the basement of the Astrophysical Observatory in Potsdam, in an attempt to damp out the ordinary tremors of the building, and it already embodied the essential idea sketched in figure 4.5, a beam split and sent along two perpendicular paths before being brought back together. Its

single, short arm and its residual sensitivity to vibration, however, produced a fringe shift far too small and unsteady to draw any firm conclusion, and Alfred Potier and Lord Rayleigh soon pointed out besides that Michelson's original analysis had understated the expected shift by a factor of two, an error traceable to treating the motion of the Earth as though it affected both arms in the same way. Correcting the theory raised the predicted shift, but what the measurement truly needed was a much longer optical path and a mounting immune to vibration, and supplying both is exactly what occupied Michelson once he moved to the Case School of Applied Science and joined forces with Edward Morley.

The apparatus that resulted rested on a massive sandstone slab floated on a pool of mercury, so that it could be rotated smoothly about a vertical axis without flexing, and each arm was folded by a set of auxiliary mirrors to stretch its effective length to roughly eleven metres, some ten times longer than in the Potsdam attempt, which is what finally brought the sensitivity of the instrument within reach of the aether drift the theory predicted, only for none to appear Michelson and Morley (1887). The same principle, two beams recombined after independent and adjustable paths, proved useful well beyond that null result. Four decades later Michelson, together with Francis Pease, mounted a twenty foot version of essentially the same interferometer across the aperture of the 100 inch Hooker telescope at Mount Wilson, and used the visibility of the fringes it produced, precisely the visibility introduced earlier in this chapter, to measure for the first time the angular diameter of a star other than the Sun, Betelgeuse, from nothing more than the separation between the two outer mirrors at which the fringes first washed out Michelson and Pease (1921). The same architecture went on to become the working principle of Fourier transform spectroscopy, where a single continuously scanned mirror encodes an entire spectrum into one interferogram, and later still of optical coherence tomography, which uses exactly this idea to image tissue a few micrometres beneath the surface.

Chapter 5

Theory of Diffraction

*Where geometry promised sharp shadows, light
insisted on painting fringes.*

“Diffraction is any deviation of light rays from rectilinear paths which
cannot be interpreted as reflection or refraction.”

– Arnold Sommerfeld

5.1 A historical context of diffraction theory: from shadows to the vector world

For more than two thousand years, from Euclid to the threshold of the Renaissance, geometrical optics rested on a single, almost self-evident assumption: light travels in straight lines Herzberger (1966). A straight ray casts a straight-edged shadow, and every child who has ever played with their own silhouette on a wall has confirmed it a thousand times. This picture explained mirrors, lenses and eclipses so well that for centuries nobody thought to look closely at the edge of a shadow itself to ask what really happens in that thin borderland where light stops and darkness begins. Diffraction is the story of what was hiding in that borderland, and of how a single overlooked detail, pursued patiently over three centuries, forced physicists to rebuild their entire understanding of what light is. It is, in a real sense, the story of light refusing to be simplified.

The first person to look carefully enough was an Italian Jesuit named Francesco Maria Grimaldi. Sometime around 1660, working in a darkened room in Bologna, Grimaldi let a narrow beam of sunlight fall through a tiny aperture and cross the edge of a rod before striking a screen. Ordinary geometry made a firm prediction: a sharp-edged shadow, dark on one side and bright on the other, with nothing more to say. What Grimaldi saw instead was stranger and far more delicate the boundary of the shadow was not a clean line but a soft fringe, laced with faint bands of colour, and the shadow itself was slightly wider than pure geometry allowed. Light was somehow reaching into the space it was not supposed to reach, and bending in ways no ray could bend. Grimaldi did not live to see his work in print; his observations appeared only after his death, in the 1665 book *Physico-Mathesis de Lumine* Grimaldi (1665). There he gave the effect its name, *diffraction*, from the Latin for “breaking into pieces,” because it looked to him as though the light itself had been shattered at the edge of the obstacle. It was a remarkably prescient observation, made with nothing more than sunlight, a dark room and a sharp eye and for a long time, almost nobody noticed it had been made.

The reason nobody noticed is largely one man’s name: Isaac Newton. Newton himself had seen delicate coloured rings form between a lens and a flat glass plate the pattern we still call Newton’s rings today and he described them meticulously in his own *Opticks* of 1704 Newton (1704). Yet Newton remained convinced that light was made of small particles, corpuscles, shot out in straight lines, and he folded these puzzling colours into that picture as best he could rather than let them overturn it. Newton’s scientific authority in the eighteenth century was so enormous that to argue against him was close to professional suicide, and so Grimaldi’s fringes, and the wave-like reasoning that might have explained them, were quietly set aside for more than a hundred years. It is one of the great cautionary tales of physics: the evidence for diffraction was sitting in plain sight the entire time, waiting for someone willing to take it seriously.

Someone did, although his idea arrived slightly ahead of its time and without quite enough machinery to finish the job. In 1690 the Dutch physicist Christiaan Huygens proposed, in his *Traité de la Lumière* Huygens (1690), an almost playful way of picturing how a wave moves forward. Imagine, Huygens said, that every single point on a wavefront is itself a tiny source, spraying out its own small circular ripple. A moment later, the new wavefront is simply the smooth envelope

that wraps around all of those countless little ripples at once. This picture now known as Huygens's principle explained reflection and refraction beautifully, and it hinted that a wave could indeed sneak sideways into a geometrical shadow, since the ripples spreading from points near the edge of an obstacle have nowhere to go but into that forbidden space. What Huygens's construction could not yet explain was why the resulting light and dark fringes appeared exactly where they did, because he had no notion that the little ripples might reinforce each other in some places and cancel each other in others. That missing idea was interference, and supplying it would take more than a century.

It finally arrived in a remarkable few years at the start of the nineteenth century. In 1801, the English polymath Thomas Young performed the experiment that every optics student meets sooner or later: he let light pass through two narrow, closely spaced slits and watched the pattern it formed on a screen beyond Young (1804). Instead of two bright bands, he saw many an entire sequence of bright and dark stripes, exactly as one would expect if two overlapping ripples on a pond could sometimes crest together and sometimes cancel to stillness. Light, Young concluded, must be a wave, and waves can add or subtract. It was a devastatingly simple experiment carrying an enormous conclusion, and for exactly that reason it was met in England with scorn rather than applause; attacking Newton was still not a popular sport. Vindication came from France. Between 1815 and 1820 the young engineer Augustin-Jean Fresnel took Huygens's wavelets and Young's interference and welded them into a single, quantitative principle the Huygens-Fresnel principle capable of predicting, for the first time, exactly how bright or dark any point in a diffraction pattern should be Fresnel (1819). Fresnel did not merely describe diffraction qualitatively, as Grimaldi and Huygens had; he calculated it, treating every point of an opening as a genuine source of secondary waves whose ripples had to be added together, crest against crest and trough against trough, to find the light at any point beyond.

Fresnel's theory then delivered one of the most celebrated moments in the history of physics. The French Academy of Sciences, still largely unconvinced, set diffraction as the subject of a prize competition, and the mathematician Siméon Denis Poisson, a confirmed sceptic of the wave picture, examined Fresnel's equations looking for an absurdity that would sink them for good. He found what he was looking for, or thought he had: Fresnel's own mathematics predicted that at the very centre of the shadow cast by a small circular disc, directly behind the obstacle where common sense guarantees total darkness, there should be a bright spot of light. Poisson presented this as a *reductio ad absurdum*, confident that no such thing could ever be observed. The academy's chairman, Dominique Arago, decided to actually perform the experiment rather than argue about it and there, glowing faintly at the heart of the disc's shadow, was the spot Poisson had predicted in mockery. The Poisson-Arago spot, as it is now known with a certain irony, became one of the most dramatic experimental confirmations any theory has ever received, and it settled the contest between particles and waves for the rest of the century.

Fresnel's success, however, was built on inspired intuition rather than rigorous derivation his integral worked extremely well, but nobody had shown that it followed necessarily from the wave equation itself. That gap was closed in 1883 by the German physicist Gustav Kirchhoff, who used the mathematics of Green's theorem to derive Fresnel's diffraction integral directly from the wave equation, placing the whole construction on a firm mathematical foundation Kirchhoff (1883).

Kirchhoff's formulation became, and largely remains, the working horse of what we now call the scalar theory of diffraction: it treats light as a single scalar ripple rather than a full electromagnetic vector, an approximation that works remarkably well whenever the obstacles involved are much larger than the wavelength of light. There was, admittedly, a subtle logical wrinkle in Kirchhoff's argument his boundary conditions demanded that both the field and its derivative vanish simultaneously behind an opaque screen, a requirement that is mathematically over-strict and it was Henri Poincaré who first pointed this out. The wrinkle was ironed out in 1896 by Arnold Sommerfeld, who found an alternative, fully self-consistent solution for diffraction by a straight edge and a half-plane, removing the inconsistency without sacrificing any of Kirchhoff's practical success Sommerfeld (1896, 1954).

All of this, ingenious as it was, still treated light as a single wiggling number instead of the vector quantity that Maxwell's equations say it truly is and by the early twentieth century that omission started to matter. In 1908 Gustav Mie solved Maxwell's equations exactly for light scattering off a small sphere, keeping the full vector and polarisation content of the field Mie (1908); his result explains, among many other things, why the sky is blue and why clouds and milk are white, and it remains indispensable wherever light interacts with particles comparable in size to its own wavelength. The following year Peter Debye extended the same vector thinking to the tight focus of a lens, showing that near a sharp focal point the scalar approximation simply breaks down. The synthesis of everything that had come before Fresnel's integral, Kirchhoff and Sommerfeld's rigour, and the full vector corrections of Mie and Debye was finally gathered into a single, monumental reference by Max Born and Emil Wolf, whose *Principles of Optics* (1959) has served as the field's bible ever since Born and Wolf (1999). Born and Wolf made explicit exactly when the simpler scalar picture is trustworthy and when it fails outright: in lenses of high numerical aperture, in anisotropic or chiral materials, and whenever the relevant structures shrink down to the scale of the wavelength itself, the vector nature of light can no longer be ignored.

The invention of the laser in 1960 gave diffraction theory an entirely new practical urgency, since it suddenly supplied a light source coherent and controllable enough to exploit diffraction on purpose rather than merely tolerate it. Joseph Goodman's *Introduction to Fourier Optics* (1968) reframed the whole subject in a beautifully economical way: an optical system, Goodman showed, can be treated as a kind of signal processor, and the diffraction pattern produced far from an aperture is, for all practical purposes, the same mathematical object as the Fourier transform of that aperture's shape Goodman (2005). This single reframing opened the door to holography, to diffractive lenses that focus light using etched patterns instead of curved glass, and to the computational correction of aberrations in modern cameras and telescopes.

The story is very much still being written. Today, engineers etch **metasurfaces** arrays of nanoscale structures far smaller than the wavelength of light that reshape a wavefront's phase and polarisation point by point, allowing a sheet barely thicker than a human hair to do the work of a stack of lenses Yu et al. (2011). Electron and X-ray diffraction, governed by exactly the same wave logic that Fresnel and Kirchhoff worked out for visible light, let scientists map the atomic architecture of crystals and molecules it was X-ray diffraction, after all, that first revealed the double-helix structure of DNA. And diffraction sets an absolute, unbreakable limit on what any optical instrument can ever resolve: no telescope, however perfectly polished, and no microscope, however expensively built,

can focus light down to a mathematical point, because the very act of passing light through a finite aperture spreads it out again. Every image a telescope or a microscope has ever produced, from the first blurred moons Galileo sketched to the sharpest pictures the largest observatories can deliver today, is diffraction's signature stamped on the light itself. Three and a half centuries after a Jesuit noticed that a shadow in a dark Bolognese room was not quite as sharp as geometry demanded, diffraction has grown from a curious imperfection into one of the deepest and most useful ideas in all of physics, and, as the chapters that follow will show in careful detail, it still has a great deal to teach us about the nature of light.

5.2 Scalar theory of diffraction

As we have proved in 3.7 and 3.8 the electromagnetic waves are very rich in their nature and the nature that can create waves. In this chapter we're going to study these wave behavior of the light, however, these are some introductory lectures, so solving these two equations are hard, and if we pretend to solve them analytically, well, that is completely impossible.

To do physics it is very often to impose assumptions on the model to make it easier to understand and develop, or even, to calculate analytical solutions. In this chapter we are going to counter how many assumptions do we need to arrive to the scalar diffraction theory. Having this on count, let's start with the first one.

Assumption 1 *In the propagation of light the magnetic interaction is completely null.*

This must be done because, in nature, the magnetic interactions are richer than the electric ones, however, it may sound funnier but it makes the solutions of 3.8 a lot harder to be calculated. Therefore, the electromagnetic formulation can be understood only by means of the electric wave equation 3.7

$$\left(\nabla^2 - \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \right) \vec{E} = \frac{1}{\epsilon_0} \nabla \rho + \mu_0 \frac{\partial}{\partial t} \vec{J}.$$

However, this wave equation is really hard to solve exactly, most of the solutions are going to be found in numerical ones. So to make the life a lot easier, we're going to use the next assumption

Assumption 2 *The electric wave propagates in the vacuum, there are no charge densities nor current densities.*

Under these assumptions, the wave equation adopts the well known form

$$\nabla^2 \vec{E} = \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \vec{E} \tag{5.1}$$

In the chapter 3.2 we start the study of the vector behavior of the electromagnetic radiation and we arrived to the concept of polarisation, which describes the vector behavior of light, and we found that, in general, this vector behavior is confined on an ellipse, the well known as the polarisation ellipse (see figure 3.3). Thus, in general, the electric vector is given by

$$\vec{E}(\vec{r}, t) = E(\vec{r}, t) \hat{e}(\vec{r}, t), \quad (5.2)$$

where $E(\vec{r}, t)$ denotes the amplitude of the field and $\hat{e}(\vec{r}, t)$ represents the polarisation vector, which besides varying in space may evolve in time in a stochastic manner.

The associated wave equation then takes the form

$$\nabla^2 [E(\vec{r}, t) \hat{e}(\vec{r}, t)] - \mu\varepsilon \frac{\partial^2}{\partial t^2} [E(\vec{r}, t) \hat{e}(\vec{r}, t)] = 0, \quad (5.3)$$

whose direct solution presents notable complexities. To tackle this problem one resorts to simplifying strategies, such as decomposing the field into scalar components. These scalar solutions may afterwards be combined to reconstruct the full vector behaviour.

The temporal variation of amplitude, phase and polarisation is analysed rigorously through coherence theory (a topic explored in depth in Chapter 6). Nevertheless, under certain approximations the temporal dependence of the amplitude is attributed solely to the intrinsic oscillations of the field, while the polarisation is regarded as static. The electric field then simplifies to

$$\vec{E}(\vec{r}, t) = E(\vec{r}, t) \hat{e}(\vec{r}), \quad (5.4)$$

allowing its expansion in an orthogonal polarisation basis,

$$\vec{E}(\vec{r}, t) = E_1(\vec{r}, t) \hat{e}_1 + E_2(\vec{r}, t) \hat{e}_2. \quad (5.5)$$

Here $\hat{e}_1$ and $\hat{e}_2$ correspond to opposite polarisation states on the Poincaré sphere. Each scalar component E_i individually satisfies the wave equation

$$\nabla^2 E_i - \mu\varepsilon \frac{\partial^2 E_i}{\partial t^2} = 0 \quad (i = 1, 2), \quad (5.6)$$

thereby decoupling the vector problem into manageable scalar equations.

In situations where one component is negligible (for instance $E_2 \rightarrow 0$), the field reduces to

$$\vec{E}(\vec{r}, t) = E(\vec{r}, t) \hat{e}, \quad (5.7)$$

defining a purely scalar case. This formalism underlines how the general vector description can emerge from the controlled superposition of scalar solutions.

5.3 The scalar wave equation (Helmholtz equation)

Let $\Psi(\mathbf{r}, t)$ be an arbitrary component of the electromagnetic field. This quantity satisfies the classical wave equation

$$\nabla^2 \Psi(\mathbf{r}, t) - \mu\epsilon \frac{\partial^2 \Psi(\mathbf{r}, t)}{\partial t^2} = 0, \quad (5.8)$$

where Ψ represents a polychromatic field. To decompose the field temporally we apply the Fourier transform

$$\Psi(\mathbf{r}, t) = \int_{\mathbb{R}^+} U(\mathbf{r}, \nu) \exp(-2\pi i \nu t) d\nu. \quad (5.9)$$

Substituting (5.9) into (5.8), the equation is re-expressed as

$$\int_{\mathbb{R}^+} \left[\nabla^2 U(\mathbf{r}, \nu) + \frac{\omega^2}{v^2} U(\mathbf{r}, \nu) \right] \exp(-2\pi i \nu t) d\nu = 0, \quad (5.10)$$

which leads directly to the Helmholtz equation for each spectral component:

$$(\nabla^2 + k^2)U(\mathbf{r}, \nu) = 0. \quad (5.11)$$

Here $k = 2\pi/\lambda$ denotes the wavenumber, and $\lambda = c/(n\nu)$ is the wavelength in a medium of refractive index n . This formulation describes mathematically how each monochromatic component of the field (characterised by its complex amplitude U) obeys an equation independent of the temporal frequency.

5.4 The Kirchhoff scalar boundary-value problem

The scalar theory of diffraction, based on solutions of the wave equation under boundary conditions, describes the propagation of fields in paraxial approximations or with smooth phase modulations. Despite its scalar formulation, the formalism - developed by Kirchhoff by means of Green's functions - is adaptable to vector fields.

The original inconsistency of Kirchhoff, pointed out by Poincaré, lies in the over-determination of the boundary conditions (prescribing both the field and its normal derivative on a plane), which violates the uniqueness theorems of differential equations. Sommerfeld resolved this by redefining the Green's function so as to eliminate the redundancy, guaranteeing mathematical consistency.

We therefore start from the physical entity under study - the electromagnetic field - and as a first step it is appropriate to recall its wave equation

$$\nabla^2 \vec{E}(\vec{r}, t) - \frac{1}{v^2} \frac{\partial^2 \vec{E}(\vec{r}, t)}{\partial t^2} = 0, \quad (5.12)$$

which is of a vector nature. We may restrict our phenomenon to the study of the propagation of its components, grouping the vector character into the polarisation vector $\hat{e}(\vec{r}, t)$. In order to avoid possible spacetime correlations between the evolution of the components and the polarisation of the field, the dependence of the polarisation vector on these variables is dropped, so that

$$\left(\nabla^2 E(\vec{r}, t) - \frac{1}{v^2} \frac{\partial^2 E(\vec{r}, t)}{\partial t^2} \right) \hat{e} \longrightarrow (\nabla^2 + k^2) E(\vec{r}, t) = 0, \quad (5.13)$$

thus obtaining the Helmholtz equation for the scalar electric field.

5.4.1 Green's theorem

Green's theorem, a direct corollary of Gauss's theorem, establishes fundamental relations between volume and surface integrals for scalar fields. Let $f(\vec{r})$ and $g(\vec{r})$ be scalar functions with continuous second derivatives in a domain V . We start from the differential identity

$$\nabla \cdot [f(\vec{r}) \nabla g(\vec{r})] = f(\vec{r}) \nabla^2 g(\vec{r}) + \nabla f(\vec{r}) \cdot \nabla g(\vec{r}). \quad (5.14)$$

Subtracting the analogous expression with f and g interchanged, we obtain

$$\nabla \cdot [f \nabla g - g \nabla f] = f \nabla^2 g - g \nabla^2 f. \quad (5.15)$$

Integrating this relation over a volume V and applying the Gauss divergence theorem,

$$\int_V \nabla \cdot [f(\vec{r}) \nabla g(\vec{r}) - g(\vec{r}) \nabla f(\vec{r})] dV = \int_V (f \nabla^2 g - g \nabla^2 f) dV, \quad (5.16)$$

which allows the volume integral on the right-hand side to be converted into a surface integral over ∂V :

$$\oint_{\Sigma(V)} [f(\vec{r}) \nabla g(\vec{r}) - g(\vec{r}) \nabla f(\vec{r})] \cdot d\vec{\sigma} = \int_V (f \nabla^2 g - g \nabla^2 f) dV, \quad (5.17)$$

where:

- $d\vec{\sigma} = \hat{n} d\sigma$ is the differential surface element oriented along the outward normal $\hat{n}$ to the surface $\Sigma(V)$.
- $\frac{\partial}{\partial n} = \hat{n} \cdot \nabla$ denotes the directional derivative along $\hat{n}$.

- The operator ∇ acts exclusively on the spatial coordinates $\vec{r}$.

This identity allows the solution of inhomogeneous differential equations (e.g. the Helmholtz equation) to be expressed in terms of integrals over the sources and the boundary conditions. The presence of $\hat{n}$ emphasises that the surface contribution depends critically on the geometry of the volume V .

5.4.2 The HelmholtzKirchhoff integral theorem

The HelmholtzKirchhoff integral theorem adapts Green's formalism to the propagation of electromagnetic fields. Starting from Green's identity and adding the term $k^2 fg$ to both sides, one obtains

$$\oint_{\Sigma(V)} [f(\vec{r})\nabla g(\vec{r}) - g(\vec{r})\nabla f(\vec{r})] \cdot d\vec{\sigma} = \int_V f(\vec{r}) (\nabla^2 + k^2) g(\vec{r}) dV - \int_V g(\vec{r}) (\nabla^2 + k^2) f(\vec{r}) dV, \quad (5.18)$$

where f and g are scalar functions. Choosing $f = U(\vec{r}, \nu)$, a solution of the Helmholtz equation, and g as an arbitrary smooth function, the expression simplifies to

$$\oint_{\Sigma(V)} [U(\vec{r}, \nu)\nabla g(\vec{r}) - g(\vec{r})\nabla U(\vec{r}, \nu)] \cdot d\vec{\sigma} = \int_V U(\vec{r}, \nu) (\nabla^2 + k^2) g(\vec{r}) dV. \quad (5.19)$$

The key lies in selecting g as the Green's function $G(\vec{r}, \vec{r}')$, defined by

$$(\nabla^2 + k^2) G(\vec{r}, \vec{r}') = 4\pi\delta(\vec{r} - \vec{r}'). \quad (5.20)$$

Remarkably, we may add an arbitrary number of Green's functions while keeping the solution invariant. What we can do is add an infinity of image charges, but *outside* the integration volume bounded by the surface $\Sigma(V)$, as shown in Figure 5.1.

In this spirit, equation (5.20) becomes

$$(\nabla^2 + k^2)G(\vec{r}, \vec{r}') = 4\pi\delta(\vec{r} - \vec{r}') + 4\pi \sum_{i=1}^{\infty} \delta(\vec{r} - \vec{r}'_i). \quad (5.21)$$

The explicit solution for a homogeneous medium of equation (5.20) is

$$G(\vec{r}, \vec{r}') = \frac{e^{ik\|\vec{r}-\vec{r}'\|}}{\|\vec{r}-\vec{r}'\|}, \quad (5.22)$$

corresponding to an outgoing spherical wave centred at $\vec{r}'$. Substituting G into Green's identity, one derives




Figure 5.1: On the boundary $\Sigma(V)$ the contribution of the field components $U(\vec{r})$ is shown; inside the region the total field is built from the contribution of all the points on the boundary. One may add infinitely many point Green charges of arbitrary propagation nature; the condition for the invariance of equation (5.20) is that these image charges lie outside $\Sigma(V)$.

$$U(\vec{r}', \nu) = \frac{1}{4\pi} \oint_{\Sigma(V)} [U(\vec{r}, \nu) \nabla G(\vec{r}, \vec{r}') - G(\vec{r}, \vec{r}') \nabla U(\vec{r}, \nu)] \cdot d\vec{\sigma}, \quad (5.23)$$

known as the Helmholtz-Kirchhoff integral formula. This equation allows $U(\vec{r}', \nu)$ to be computed at any point of the volume V by means of an integral over its boundary $\Sigma(V)$.




Figure 5.2: Representation of the Green's functions for point sources. Under these boundary conditions the Green's functions behave as point charges, one the mirror image of the other.

Before continuing, it is worth noting that the point-charge nature of the Green's functions $G(\vec{r}, \vec{r}')$ is an assumption taken for granted; there is, however, no restriction on their physical nature. As is well known in electromagnetic theory, radiation sources may have any volume and shape, so our problem (Figure 5.2) can be made more general, as shown in Figure 5.3, and the differential equation for Green's functions with volume would read

$$(\nabla^2 + k^2) G(\vec{r}, \vec{r}') = 4\pi\rho(\vec{r} - \vec{r}'_+) - 4\pi\rho(\vec{r} - \vec{r}'_-). \quad (5.24)$$

Let us continue with the development for point Green's functions. To simplify the evaluation of the integral, we consider a volume V bounded by an infinite plane Σ_1 ($z = 0$) and a hemisphere Σ_2




Figure 5.3: Representation of the Green's functions for non-point sources. Under these boundary conditions the Green's functions still behave as mirror images, but now with given volume and geometry; the only constraint on the emission of these sources is that, on the boundary $z = 0$, the total field must vanish, as required by the assumed Dirichlet conditions.

of radius R (see Figure 5.2). The contribution of Σ_2 vanishes as $R \rightarrow \infty$ (owing to the Sommerfeld radiation condition), reducing the expression to

$$U(\vec{r}', \nu) = \frac{1}{4\pi} \int_{\Sigma_1} \left[U(\vec{r}, \nu) \frac{\partial G}{\partial n} - G(\vec{r}, \vec{r}') \frac{\partial U}{\partial n} \right] d\sigma, \quad (5.25)$$

where $\frac{\partial}{\partial n} = \hat{n} \cdot \nabla$ denotes the normal derivative. However, this formulation requires specifying simultaneously U and $\partial U / \partial n$ on Σ_1 , which over-determines the problem. This difficulty motivates the introduction of physically realistic boundary conditions, such as those of Kirchhoff, in order to resolve the ambiguity.

This problem is aggravated by a theorem of complex analysis, attributed to Riemann, which states that if a function and its normal derivative vanish on an open region of a plane, then the function must be identically zero throughout the domain. This restriction motivates the strategic need to eliminate one of the terms in the Helmholtz-Kirchhoff surface integral.

The key is to exploit the degrees of freedom inherent in the choice of the auxiliary Green's function G . Kirchhoff initially proposed the intuitive boundary conditions

- $U = 0$ outside the aperture on Σ (blocked field),
- $\partial U / \partial n = 0$ on opaque regions (vanishing normal derivative).

Although these conditions reproduced the experimental results of diffraction, they introduced mathematical inconsistencies by violating the Cauchy-Riemann uniqueness theorem. Poincaré showed that such assumptions generate contradictions in the mixed derivatives of the field. The resolution came with Sommerfeld, who redefined G by means of a double-layer (mirror) Green's function, ensuring that

- $G = 0$ on Σ (Dirichlet condition),
- $\partial G / \partial n$ remains non-singular,

thus eliminating the redundancy in the boundary conditions and preserving the self-consistency of the formalism.

5.4.3 The Rayleigh-Sommerfeld formulation

Returning to equation (5.23) and choosing the Dirichlet condition for the function G , we have

$$U(\vec{r}', \nu) = \frac{1}{4\pi} \int_{\Sigma_1} [U(\vec{r}, \nu) \nabla G \cdot \hat{n}] d\sigma. \quad (5.26)$$

Given the integration surface under consideration, we may express (5.22) as

$$G(\vec{r}, \vec{r}') = \frac{e^{ik\|\vec{r}-\vec{r}'\|}}{\|\vec{r}-\vec{r}'\|} - \frac{e^{ik\|\vec{r}-\vec{r}''\|}}{\|\vec{r}-\vec{r}''\|}, \quad (5.27)$$

where

$$\|\vec{r}-\vec{r}''\| = \|\vec{r}-\vec{r}'\|, \quad x'' = x', \quad y'' = y', \quad z'' = -z', \quad (5.28)$$

since $\vec{r}''$ is the mirror image of $\vec{r}'$ across the plane Σ_1 . It follows that

$$\nabla G \cdot \hat{z} = 2 \left[ik - \frac{1}{\|\vec{r}-\vec{r}''\|} \right] \frac{e^{ik\|\vec{r}-\vec{r}''\|}}{\|\vec{r}-\vec{r}''\|} \cos(\vec{r}-\vec{r}', \hat{z}), \quad (5.29)$$

and with this, (5.26) becomes

$$U(\vec{r}', \nu) = \frac{1}{2\pi} \int_{\Sigma} U(\vec{r}, \nu) \left[ik - \frac{1}{\|\vec{r}-\vec{r}''\|} \right] \frac{e^{ik\|\vec{r}-\vec{r}''\|}}{\|\vec{r}-\vec{r}''\|} \cos(\vec{r}-\vec{r}', \hat{z}) d\sigma. \quad (5.30)$$

We may now make the approximation appropriate to the RayleighSommerfeld regime, in which $\|\vec{r}-\vec{r}'\| \gg \lambda$, obtaining

$$U(\vec{r}', \nu) = \frac{ik}{2\pi} \int_{\Sigma} U(\vec{r}, \nu) \frac{e^{ik\|\vec{r}-\vec{r}''\|}}{\|\vec{r}-\vec{r}''\|} \cos(\vec{r}-\vec{r}', \hat{z}) d\sigma, \quad (5.31)$$

a mathematical expression that embodies the HuygensFresnel principle: as can be appreciated, there is one spherical wave per point of the integration surface.

5.5 The Fresnel approximation

Starting from the field

$$U(\vec{r}') = \frac{ik}{2\pi} \int_{\Sigma} U(\vec{r}) \frac{e^{ik\|\vec{r}-\vec{r}''\|}}{\|\vec{r}-\vec{r}''\|} \cos(\vec{r}-\vec{r}', \hat{z}) d\sigma, \quad (5.32)$$

it is possible to carry out the so-called mid-field or Fresnel approximation. To this end we take

$$\begin{aligned} \|\vec{r}-\vec{r}'\| &= [(x-x')^2 + (y-y')^2 + (z-z')^2]^{1/2} \\ &= z \left(1 + \frac{(x-x')^2 + (y-y')^2}{z^2} \right)^{1/2} \\ &\approx z + \frac{(x-x')^2 + (y-y')^2}{2z}. \end{aligned} \quad (5.33)$$

It is clear that, for the factor $\frac{1}{\|\vec{r} - \vec{r}'\|}$, an expansion up to the first term of (5.33) suffices; this is not enough for the factor $e^{ik\|\vec{r} - \vec{r}'\|}$, however, since the latter has oscillatory behaviour over its domain, so both terms are required. We additionally take $\cos(\vec{r} - \vec{r}', \hat{z}) = \frac{z}{\|\vec{r} - \vec{r}'\|} \approx 1$.

In this spirit, we obtain an expression for the field of the form

$$U(\vec{r}') = \frac{ie^{ikz}}{\lambda z} \int U(\vec{r}) \exp\left(\frac{ik}{2z}((x - x')^2 + (y - y')^2)\right) d\sigma, \quad (5.34)$$

which describes the propagation or diffraction of the electromagnetic field in the Fresnel regime. It is worth stressing that equation (5.34) describes a parabolic wave, and although, at least conceptually, such propagation is valid for a mid-field, in practice it has proved more than sufficient for sufficiently short distances.

5.5.1 Example: Fraunhofer diffraction between two confocal spheres

Starting from the Fresnel integral (5.34), show that between two confocal spheres one obtains Fraunhofer diffraction.

Before anything else it is convenient to expand the square in the exponent of (5.34). Writing $r^2 = x^2 + y^2$ for the transverse radius on the emitting plane and $r'^2 = x'^2 + y'^2$ for the one on the receiving plane,

$$(x - x')^2 + (y - y')^2 = r^2 + r'^2 - 2(xx' + yy'), \quad (5.35)$$

so that the Fresnel integral separates into three factors,

$$U(\vec{r}') = \frac{ie^{ikz}}{\lambda z} \int U(\vec{r}) e^{\frac{ik}{2z}r^2} e^{\frac{ik}{2z}r'^2} e^{-\frac{ik}{z}(xx' + yy')} d\sigma. \quad (5.36)$$

The last factor is already the kernel of a Fourier transformation. The obstacle is the pair of quadratic factors, and the whole point of the exercise is that they are not an obstacle at all if one agrees to read the field on a curved surface instead of on a plane.

Consider a spherical cap tangent to a transverse plane at the optical axis, with its centre of curvature on the axis at the signed distance c from the vertex, counted positive in the direction of propagation. The point of the cap at the transverse position $\vec{r}$ is displaced axially from the tangent plane by the sagitta

$$\delta(r) = c - \text{sgn}(c)\sqrt{c^2 - r^2} = c - \text{sgn}(c)|c|\sqrt{1 - \frac{r^2}{c^2}} \approx c - \text{sgn}(c)\left(1 - \frac{r^2}{2c^2}\right),$$

where we used $\text{sgn}(c)|c| = c$ and expanded the root to first order, which is the same paraxial expansion already performed in (5.33). The two terms in red cancel and there remains

$$\delta(r) = \frac{r^2}{2c}. \quad (5.37)$$

Since a field propagating in the positive direction acquires the factor $e^{ik\delta}$ over an axial displacement δ , the field read on the cap and the field read on the tangent plane are related by

$$u_{\text{cap}}(\vec{r}) = U(\vec{r}) \exp\left(\frac{ik r^2}{2c}\right). \quad (5.38)$$

Two spheres are said to be **confocal** when each one has its centre of curvature at the vertex of the other. Let therefore $\mathcal{A}$ be the cap tangent to the emitting plane $\mathcal{P}$ at the vertex O and centred at O' , and let $\mathcal{F}$ be the cap tangent to the receiving plane $\mathcal{Q}$ at the vertex O' and centred at O , as shown in Figure 5.4. Both radii are then equal to the propagation distance, and with the sign convention of (5.38) the two signed distances are

$$c_{\mathcal{A}} = +z, \quad c_{\mathcal{F}} = -z, \quad (5.39)$$

the second one being negative because the centre of $\mathcal{F}$ lies upstream of its own vertex.

Applying (5.38) on each side with the radii (5.39),

$$U(\vec{r}) = u_{\mathcal{A}}(\vec{r}) e^{-\frac{ik}{2z}r^2}, \quad u_{\mathcal{F}}(\vec{r}') = U(\vec{r}') e^{-\frac{ik}{2z}r'^2}, \quad (5.40)$$

and substituting both relations into (5.36),

$$u_{\mathcal{F}}(\vec{r}') = \frac{ie^{ikz}}{\lambda z} e^{-\frac{ik}{2z}r'^2} e^{\frac{ik}{2z}r'^2} \int u_{\mathcal{A}}(\vec{r}) e^{-\frac{ik}{2z}r^2} e^{\frac{ik}{2z}r^2} e^{-\frac{ik}{z}(xx'+yy')} d\sigma,$$

where the four exponentials in red cancel in pairs, since in each pair the two exponents are opposite. We are left with

$$\boxed{u_{\mathcal{F}}(\vec{r}') = \frac{ie^{ikz}}{\lambda z} \int u_{\mathcal{A}}(\vec{r}) \exp\left[-\frac{ik}{z}(xx' + yy')\right] d\sigma,} \quad (5.41)$$

which is exactly the Fraunhofer expression (5.45) *without the curvature factor*. This proves the assertion.

Three comments are in order, because the result is stronger than it looks.

In the derivation of (5.45) the factor $e^{ikr^2/2z}$ was *neglected* by appealing to the far field. Here it has not been neglected, it has been *absorbed* by the choice of reference surface, and (5.41) is




Figure 5.4: Geometry of the two confocal spheres. The cap $\mathcal{A}$ is tangent to the emitting plane $\mathcal{P}$ at the vertex O and has its centre of curvature at O' , while the cap $\mathcal{F}$ is tangent to the receiving plane $\mathcal{Q}$ at O' and has its centre of curvature at O , so that both radii equal the propagation distance z . Read on these two surfaces, the Fresnel propagation reduces to an exact Fourier transformation, which in the language of the fractional Fourier transformation of Section 5.7 is the order $\alpha = \pi/2$.

therefore exact within the Fresnel approximation, for any propagation distance z . The lesson is that Fraunhofer diffraction is not really a regime of distances but a statement about geometry: it is what Fresnel propagation looks like when the field is read on two confocal spheres. The curvature factor that survives outside the integral in (5.45) is nothing but the price of insisting on reading the field on a plane, and this is precisely the interpretation anticipated in Figure 5.5.

Introducing the spatial frequency associated with the observation point,

$$\vec{\nu} = \frac{\vec{r}'}{\lambda z}, \quad \frac{k}{z} (xx' + yy') = \frac{2\pi}{\lambda z} \vec{r} \cdot \vec{r}' = 2\pi \vec{r} \cdot \vec{\nu}, \quad (5.42)$$

equation (5.41) becomes

$$u_{\mathcal{F}}(\lambda z \vec{\nu}) = \frac{ie^{ikz}}{\lambda z} \widehat{u_{\mathcal{A}}}(\vec{\nu}), \quad \widehat{u_{\mathcal{A}}}(\vec{\nu}) = \int u_{\mathcal{A}}(\vec{r}) e^{-2\pi i \vec{r} \cdot \vec{\nu}} d\sigma, \quad (5.43)$$

so that the observation coordinate on the sphere $\mathcal{F}$ is, up to the scale factor λz , the spatial frequency of the field on the sphere $\mathcal{A}$. This is the precise sense in which a diffracting system performs a Fourier analysis of the object, the statement that Abbe had reached qualitatively in 1873 (Abbe, 1873).

The only approximations used are those already contained in (5.34) together with the first order expansion (5.37) of the sagitta, so the result holds as long as the illuminated region satisfies $r_{\max} \ll z$, which is the condition for the caps not to depart appreciably from their tangent planes. When this ceases to hold one must abandon the paraxial description, exactly as noted at the end of the following section.

Finally, it is worth recording that this example is the first instance of a much more general fact. Nothing forces the two reference spheres to be confocal, and if they are given a common radius R different from z the two quadratic factors no longer cancel but combine into a chirp, and the propagation becomes a fractional Fourier transformation of an order fixed by the ratio z/R . The confocal case treated here is the particular order $\alpha = \pi/2$. This is the subject of Section 5.7.

5.6 The Fraunhofer approximation

Finally, from expression (5.34) a second approximation may be carried out, corresponding to the far field or Fraunhofer regime, in which the observation distances are considerably larger than the area over which one wishes to study the propagated field (whether through a slit or another object). We focus on the factor

$$\exp\left(\frac{ik}{2z}((x-x')^2 + (y-y')^2)\right) = \exp\left(\frac{ik}{2z}(x^2 + y^2)\right) \exp\left(\frac{ik}{2z}(x'^2 + y'^2)\right) \exp\left(-\frac{ik}{z}(xx' + yy')\right), \quad (5.44)$$

where we may make the far-field assumption $\exp\left(\frac{ik}{2z}(x^2 + y^2)\right) \approx 1$, so that we obtain

$$U(\vec{r}') = \frac{e^{ikz}}{\lambda z} \exp\left(\frac{ik}{2z}(x'^2 + y'^2)\right) \int U(\vec{r}) \exp\left(-\frac{ik}{z}(xx' + yy')\right) d\sigma. \quad (5.45)$$

There we can still appreciate a quadratic phase term outside the integral, usually called the curvature factor, which shows precisely the dephasing induced by the difference of optical path between a plane called the observation screen and a spherical surface of radius z (the propagation distance of the field along the optical axis) known as the cardinal sphere. The introduction of this spherical surface is rather useful when one wishes to describe Fraunhofer diffraction on a screen as a Fresnel propagation through two confocal cardinal spheres, as shown in Figure 5.5.

This makes more sense once we analyse the effect of each propagation step. It is clear that, without imposing far-field approximations from the outset, propagating the field at $\mathcal{A}$ towards $\mathcal{F}$ with the Fresnel approximation, we would obtain

$$U_{\mathcal{F}}(\vec{r}') = \frac{ie^{ikz}}{\lambda z} \exp\left(\frac{ik}{2z}(x^2 + y^2)\right) \int u_{\mathcal{A}} \exp\left(\frac{ik}{2z}(x'^2 + y'^2)\right) \exp\left(-\frac{ik}{z}(xx' + yy')\right) d\sigma', \quad (5.46)$$




Figure 5.5: Scheme of the confocal cardinal spheres, with radii equal to the propagation distance z of the field.

where we may associate $u_{\mathcal{A}}$, i.e. the field on the sphere $\mathcal{A}$, with the field on $\mathcal{P}$ multiplied by its corresponding quadratic factor $u_{\mathcal{P}} = u_{\mathcal{A}}(\vec{r}') \exp\left(\frac{ik}{2z}(x'^2 + y'^2)\right)$, while the term $\exp\left(\frac{ik}{2z}(x^2 + y^2)\right)$ may be understood as the factor resulting from passing the field from the sphere $\mathcal{F}$ to the screen $\mathcal{Q}$, so that the field on $\mathcal{F}$ would correspond to

$$u_{\mathcal{Q}}(\vec{r}) = \frac{ie^{ikz}}{\lambda z} \int u_{\mathcal{P}} \exp\left(-\frac{ik}{z}(xx' + yy')\right) d\sigma', \quad (5.47)$$

thus recovering a Fraunhofer expression for diffraction on $\mathcal{Q}$. It should be noted that one must check whether the study of the propagated field is valid within a paraxial regime, where the field is confined to the neighbourhood of the optical axis and hence the curvature of the cardinal sphere does not differ much from the plane (the screen). In cases where the curvature factor ceases to be negligible, the paraxial regime loses validity and the problem must be addressed from a meta-axial standpoint.

Studying the propagation of the field through Fresnel diffraction is of great importance because, as will be discussed later, it bears a close relationship with what is known as the fractional Fourier transform, a tool that arises naturally when one chains paraxial propagation steps.

5.7 The fractional Fourier transformation

The last section closed with the remark that Fresnel propagation bears a close relationship with the fractional Fourier transformation. That remark is the subject of the present section, and it deserves a careful treatment for two reasons. The first is that the fractional Fourier transformation is not a formal curiosity but the natural language in which paraxial propagation should be written, in the same sense in which the ordinary Fourier transformation is the natural language of the far field. The second is that the object itself, a continuous family of operators interpolating between the identity and the Fourier transformation, has a long and instructive history, one in which the same idea was rediscovered independently in pure analysis, in quantum mechanics, in signal processing and in optics before anybody realised that all four communities were talking about the same one parameter group.

5.7.1 A historical context of Fourier analysis and of its fractional powers

The story begins, as so many stories in mathematical physics do, with a vibrating string. In 1747 Jean le Rond d'Alembert wrote down the wave equation of a stretched string and solved it by superposing two travelling waves (d'Alembert, 1747), a solution that Leonhard Euler immediately generalised to arbitrary initial shapes (Euler, 1748). Six years later Daniel Bernoulli proposed something that looked, to his contemporaries, physically obvious and mathematically outrageous at the same time, namely that every possible motion of the string could be written as a sum of the sinusoidal modes of vibration (Bernoulli, 1753). Euler and d'Alembert both rejected the claim. Their

objection was not aesthetic but logical, since a sum of sines is a smooth and periodic object while the initial shape of a plucked string has a corner in it, and no eighteenth century mathematician was willing to accept that an infinite sum of analytic functions could reproduce a curve with a kink. The dispute lasted half a century and was never settled by the people who started it.

It was settled, in a sense, by somebody who was not trying to settle it. In 1807 Joseph Fourier submitted to the Institut de France a memoir on the propagation of heat in solids in which he asserted, and used, precisely the representation that Euler had rejected. The referees, Lagrange among them, refused the memoir. Lagrange in particular never accepted the generality of the trigonometric expansion, and Fourier had to wait until 1822 to publish the mature version of his ideas as the *Théorie Analytique de la Chaleur* (Fourier, 1822). What Fourier had understood, and what his referees had not, is that the question of whether the series converges to the function is a separate question from whether the series is useful, and that the coefficients can be computed by an integral regardless of one's philosophical position about convergence. The passage from the series on a finite interval to the integral on the whole line, which is what we now call the Fourier transformation, is already present in that book as a limiting case.

Fourier's boldness left a debt that the nineteenth century spent decades paying. Peter Gustav Lejeune Dirichlet gave in 1829 the first genuine convergence theorem, showing that the series of a piecewise monotone function with finitely many discontinuities converges to the mean of the lateral limits (Dirichlet, 1829). Bernhard Riemann, in his habilitation of 1854 published posthumously in 1867, asked the converse question of which functions admit a trigonometric representation at all, and in doing so was forced to invent a theory of integration (Riemann, 1867). Georg Cantor, studying the uniqueness of trigonometric representations, was led to classify the exceptional sets on which two different series can agree, and out of that classification grew set theory itself (Cantor, 1872). Henri Lebesgue built a new integral partly in order to make the theory of trigonometric series behave (Lebesgue, 1903). It is not an exaggeration to say that a good part of modern analysis exists because Fourier wrote down a formula that nobody could justify.

The theory reached its definitive form in the twentieth century. Michel Plancherel proved in 1910 the identity that bears his name, establishing that the Fourier transformation is an isometry of the space of square integrable functions onto itself (Plancherel, 1910). This is the single most important structural fact about the transformation for our purposes, because an isometry of a Hilbert space onto itself is a unitary operator, and unitary operators are exactly the objects that can be raised to continuous powers. Norbert Wiener (1933) and Edward Titchmarsh (1937) consolidated the analytic theory in the two treatises that defined the subject for a generation, and in 1965 James Cooley and John Tukey rediscovered an algorithm that reduced the numerical cost of the discrete transformation from a quadratic to a logarithmic law and thereby moved Fourier analysis from the blackboard into every laboratory instrument (Cooley and Tukey, 1965).

Optics entered the story from a different door. Ernst Abbe had understood already in 1873, while trying to explain why his microscopes could not resolve arbitrarily fine detail, that the objective of a microscope performs a decomposition of the object into spatial frequencies and that the aperture of the instrument truncates that decomposition (Abbe, 1873). The idea remained essentially qualitative until Pierre-Michel Duffieux published in 1946 the first systematic treatise on the Fourier integral in

optics (Duffieux, 1946), which is the direct ancestor of the modern treatment found in Goodman (2005). By the middle of the twentieth century it was a commonplace that the far field of an aperture is the Fourier transform of the aperture, which is the statement we derived in the section on the Fraunhofer approximation.

The idea of a *fractional* power of the Fourier transformation has a separate and much less linear genealogy. Its seed is a purely analytic observation. Charles Hermite had introduced in 1864 the polynomials that carry his name (Hermite, 1864), and Ferdinand Gustav Mehler had found in 1866 the bilinear generating function of those polynomials (Mehler, 1866), a formula that will do all the work for us later in this section. In 1929 Norbert Wiener published a short note in which he observed that the Hermite functions are eigenfunctions of the Fourier transformation with eigenvalues that are the four fourth roots of unity (Wiener, 1929). The observation is elementary once stated, but it is the whole idea, because an operator whose spectrum is known can be raised to any power one likes simply by raising the eigenvalues to that power.

Edward Uhler Condon took that step explicitly in 1937 in a paper whose title says exactly what it does, namely to immerse the Fourier transformation in a continuous group of functional transformations (Condon, 1937). Condon was interested in the group theoretical structure and in the connection with the quantum harmonic oscillator, and he already noticed that the generator of the continuous family is the oscillator Hamiltonian. Two years later Hermann Kober, working from the point of view of integral transformations, constructed roots of the Fourier and Hankel transformations and gave the corresponding kernels (Kober, 1939). Valentine Bargmann arrived at the same family in 1961 from yet another direction, through his Hilbert space of entire functions, where the fractional powers act as simple rotations of the complex variable (Bargmann, 1961). Nicolaas de Bruijn recovered them once more in 1973 in his theory of generalised functions built on the Wigner distribution (de Bruijn, 1973). Each of these authors had the object in his hands and none of them gave it a name that stuck.

The name and the systematic treatment are due to Victor Namias, who in 1980 published the paper that the optics community now takes as the origin of the subject (Namias, 1980). Namias' contribution was not merely terminological. He defined the transformation of fractional order in two equivalent ways, as a differential operator obtained by exponentiating the harmonic oscillator Hamiltonian and as an integral operator with an explicit kernel, showed that the two definitions agree, and then used the resulting tool to solve concrete problems in quantum mechanics that are awkward in the ordinary Fourier language. It is his derivation that we shall reproduce in full in the next subsection.

Namias' treatment was, by his own standards, a working physicist's treatment. The integral he wrote does not converge absolutely for every function one would like to transform, the square root appearing in the normalising constant is written without specifying which branch is meant, and the additivity of the orders is verified on the eigenfunction expansion rather than proved on a well defined domain. These are not pedantic objections, because a careless choice of branch makes the composition law fail by a sign. Adam McBride and Fiona Kerr repaired all of this in 1987 (McBride and Kerr, 1987), giving a definition that converges absolutely on the Schwartz space, fixing the branch in a way that makes the composition law hold for all real orders, and proving that the

resulting family is a genuine continuous one parameter group. Kerr extended the construction shortly afterwards to spaces of generalised functions (Kerr, 1988a) and to the space of square integrable functions with applications to differential equations (Kerr, 1988b).

While the analysts were tidying up, the optics community rediscovered the object experimentally. In 1993 David Mendlovic and Haldun Ozaktas published a pair of papers showing that a graded index medium of the appropriate length performs a fractional Fourier transformation on the field that traverses it (Mendlovic and Ozaktas, 1993; Ozaktas and Mendlovic, 1993), and Adolf Lohmann showed in the same year that the fractional order is nothing other than a rotation angle of the Wigner distribution of the field, which gave the whole construction a transparent phase space meaning and an immediate bulk optics implementation with two lenses (Lohmann, 1993). One year later Pierre Pellat-Finet proved the result that concerns us most directly, namely that Fresnel diffraction between two spherical caps is exactly a fractional Fourier transformation, with an order fixed by the geometry (Pellat-Finet, 1994), a result developed at length with Georges Bonnet (Pellat-Finet and Bonnet, 1994) and later incorporated into a full treatise on metaxial and fractional Fourier optics (Pellat-Finet, 2009). In parallel, Luís Almeida introduced the transformation into signal processing as an interpolation between the time and the frequency representations (Almeida, 1994), and the whole body of optical applications was collected by Ozaktas, Zalevsky and Kutay (2001).

It is worth pausing on the fact that the literature does not use a single notion of fractional order, and a reader who moves between papers will meet at least five conventions. The first is the one introduced by Namias, in which the order is an angle α measured in radians, the ordinary Fourier transformation corresponding to $\alpha = \pi/2$ and the identity to $\alpha = 0$. The second, standard in signal processing after Ozaktas, Mendlovic and Lohmann, replaces the angle by a dimensionless order a defined through $\alpha = a\pi/2$, so that the ordinary transformation is the transformation of order one and the parity operator is the transformation of order two. The third is the optical order used by Pellat-Finet, which is again an angle but is tied to a pair of reference spheres rather than to an abstract eigenvalue assignment, and which therefore depends on the geometry chosen to describe the propagation. The fourth is the complex order, introduced by Chung-Chieh Shih (1995b) and realised optically by Luís Bernardo and Olívio Soares (1996), in which the parameter is allowed to leave the real axis and the transformation acquires a Gaussian weighting in addition to its chirp. The fifth is not a convention but a genuine ambiguity, and it deserves emphasis. Since the ordinary Fourier transformation satisfies $\mathcal{F}^4 = \mathcal{I}$, the problem of defining $\mathcal{F}^a$ is the problem of choosing a fourth root of the identity in a continuous way, and that choice is not unique. Namias assigns to the eigenfunction of index n the eigenvalue e^{-ina} , which is the choice that makes the generator the harmonic oscillator Hamiltonian, but Shih showed that a different and perfectly consistent assignment, based on weighting the four powers of $\mathcal{F}$, produces an inequivalent family (Shih, 1995a). Gianfranco Cariolaro and collaborators later classified all admissible fractionalisations in a single framework (Cariolaro et al., 1998), and Çağatay Candan, Alper Kutay and Ozaktas solved the corresponding discrete problem (Candan et al., 2000). When a paper speaks of *the* fractional Fourier transformation without further qualification, it almost always means the Namias family, and that is the one we shall construct.

Finally, it should be said that the fractional Fourier transformation is not an isolated object but

the simplest nontrivial member of a much larger family. Marcos Moshinsky and Christiane Quesne introduced in 1971 the linear canonical transformations, the unitary representations of the group of linear transformations of phase space that preserve the symplectic form (Moshinsky and Quesne, 1971), and Kurt Bernardo Wolf developed their theory as integral transformations (Wolf, 1979). Stuart Collins had written in 1970 the diffraction integral of an arbitrary paraxial system in terms of the matrix of that system (Collins, 1970), which is the same family seen from the engineering side. In that language the fractional Fourier transformations are exactly the elliptic elements of the group, that is, the rotations of phase space, while free propagation over a plane and the action of a thin lens are parabolic elements and magnifications are hyperbolic elements. The reason the fractional Fourier transformation occupies such a privileged place in optics is that, as we shall prove, once the field is referred to the right pair of spherical surfaces, free propagation becomes a pure rotation.

5.7.2 The fractional Fourier transformation as a differential and as an integral operator

Throughout this subsection we work in one transverse variable, since the two dimensional case follows by taking the tensor product of two identical one dimensional transformations, and we adopt the symmetric convention

$$\mathcal{F}[f](x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(y) e^{-ixy} dy, \quad (5.48)$$

which is the convention that makes $\mathcal{F}$ unitary on $L^2(\mathbb{R})$ by the theorem of Plancherel (1910), and which agrees with the sign of the kernel that appeared in the Fraunhofer integral (5.45). Applying (5.48) twice gives the parity operator and four times the identity,

$$\mathcal{F}^2 = \mathcal{P}, \quad \mathcal{P}f(x) = f(-x), \quad \mathcal{F}^4 = \mathcal{I}, \quad (5.49)$$

so that the integer powers of $\mathcal{F}$ form a cyclic group of order four. The entire content of this subsection is the construction of a continuous group in which that cyclic group of four elements sits as a subgroup.

Before starting, a point of vocabulary must be settled, because the two words involved are routinely confused and the distinction is essential to everything that follows.

Definition 25 (*Transformation and transform*) A **transformation** is an operator, that is, a map $\mathcal{F}_\alpha : \mathcal{H} \rightarrow \mathcal{H}$ from a function space into itself. A **transform** is the image of one particular element under that map, that is, the function $\mathcal{F}_\alpha f$ obtained by applying the transformation to a given f .

The distinction is not pedantic. Everything that makes the fractional Fourier construction worth having is a statement about *transformations* and not about transforms: the family $\{\mathcal{F}_\alpha\}_{\alpha \in \mathbb{R}}$ is a one parameter group, its composition law is $\mathcal{F}_\alpha \mathcal{F}_\beta = \mathcal{F}_{\alpha+\beta}$, its generator is the harmonic oscillator Hamiltonian, and its elements are unitary. None of these sentences can even be formulated at the

level of a single transform, because a transform is merely a function and carries no group structure whatsoever. In the optical application of the last subsection the point becomes physical rather than linguistic: it is the propagation of the field over a distance that is a transformation, while the field observed on the receiving surface is a transform, and the reason propagation composes so simply is a property of the operators, not of any particular field.

5.7.2.1 The eigenfunctions of the Fourier transformation

The construction rests on a single fact, observed by Wiener in 1929 (Wiener, 1929), namely that the Fourier transformation possesses a complete orthonormal set of eigenfunctions built out of the Hermite polynomials (Hermite, 1864). Let us derive it, since every later step depends on the precise value of the eigenvalues. Recall the generating function of the Hermite polynomials,

$$e^{2xt-t^2} = \sum_{n=0}^{\infty} \frac{H_n(x)}{n!} t^n, \quad (5.50)$$

and define the generating function of the Hermite functions by multiplying (5.50) by the Gaussian $e^{-x^2/2}$,

$$G(x, t) = e^{-x^2/2} e^{2xt-t^2} = e^{-x^2/2} \sum_{n=0}^{\infty} \frac{H_n(x)}{n!} t^n. \quad (5.51)$$

We now transform (5.51) with respect to x . Writing the full exponent of the integrand of (5.48),

$$\begin{aligned} -\frac{x^2}{2} + 2xt - t^2 - i\xi x &= -\frac{x^2}{2} + x(2t - i\xi) - t^2 = \\ &= -\frac{1}{2} \left[x^2 - 2x(2t - i\xi) + \frac{(2t - i\xi)^2}{2} \right] + \frac{(2t - i\xi)^2}{2} - t^2, \end{aligned}$$

where the two terms in blue have been added and subtracted, which is legitimate because their sum is zero, and which turns the first bracket into a perfect square,

$$= -\frac{1}{2} [x - (2t - i\xi)]^2 + \frac{(2t - i\xi)^2}{2} - t^2.$$

The remaining integral is the Gaussian $\int_{-\infty}^{\infty} e^{-\frac{1}{2}(x-a)^2} dx = \sqrt{2\pi}$, which is valid for complex a by a translation of the integration contour, so that the factor $1/\sqrt{2\pi}$ in (5.48) is cancelled exactly and

$$\mathcal{F}[G(\cdot, t)](\xi) = \exp \left[\frac{(2t - i\xi)^2}{2} - t^2 \right]. \quad (5.52)$$

It remains to recognise the right hand side. Expanding the square,

$$\begin{aligned} \frac{4t^2 - 4it\xi - \xi^2}{2} - t^2 &= 2t^2 - 2it\xi - \frac{\xi^2}{2} - t^2 = -\frac{\xi^2}{2} - 2i\xi t + t^2 = \\ &= -\frac{\xi^2}{2} + 2\xi(-it) - (-it)^2, \end{aligned}$$

since $-(-it)^2 = +t^2$, and comparing with the definition (5.51) we conclude that

$$\mathcal{F}[G(\cdot, t)](\xi) = G(\xi, -it). \quad (5.53)$$

The generating function is mapped into itself with the parameter rotated by a quarter turn in the complex plane. Expanding both sides of (5.53) in powers of t by means of (5.51),

$$\sum_{n=0}^{\infty} \frac{t^n}{n!} \mathcal{F}[H_n e^{-x^2/2}](\xi) = \sum_{n=0}^{\infty} \frac{(-it)^n}{n!} H_n(\xi) e^{-\xi^2/2},$$

and identifying the coefficients of t^n , which is legitimate because both series converge for every t in a neighbourhood of the origin, we obtain the result we were after. Introducing the normalised Hermite functions

$$\psi_n(x) = (2^n n! \sqrt{\pi})^{-1/2} H_n(x) e^{-x^2/2}, \quad \int_{-\infty}^{\infty} \psi_n(x) \psi_m(x) dx = \delta_{nm}, \quad (5.54)$$

which form a complete orthonormal basis of $L^2(\mathbb{R})$, the eigenvalue relation reads

$$\mathcal{F}[\psi_n] = (-i)^n \psi_n = e^{-in\pi/2} \psi_n. \quad (5.55)$$

Equation (5.55) contains the whole idea in a single line. The eigenvalues of the Fourier transformation are $e^{-in\pi/2}$, and the number $\pi/2$ appears there not because of any deep property of the exponential but simply because $\mathcal{F}$ is the fourth root of the identity. Nothing forbids us from replacing $\pi/2$ by an arbitrary angle.

5.7.2.2 Namias' definition and the differential operator

Following Namias (1980), we define the fractional Fourier transformation of order α by prescribing its action on the basis (5.54),

$$\mathcal{F}_\alpha[\psi_n] = e^{-in\alpha} \psi_n, \quad (5.56)$$

and extending it to an arbitrary $f \in L^2(\mathbb{R})$ by linearity and continuity. Writing the expansion of f in the basis,

$$f(x) = \sum_{n=0}^{\infty} c_n \psi_n(x), \quad c_n = \int_{-\infty}^{\infty} f(y) \psi_n(y) dy, \quad (5.57)$$

the transform of order α is by definition

$$F_\alpha(x) \equiv \mathcal{F}_\alpha[f](x) = \sum_{n=0}^{\infty} c_n e^{-in\alpha} \psi_n(x). \quad (5.58)$$

The definition is manifestly consistent, since $\alpha = 0$ gives $F_0 = f$ by completeness, $\alpha = \pi/2$ reproduces (5.55), $\alpha = \pi$ gives $\sum c_n (-1)^n \psi_n$, which is the parity operator because ψ_n has the parity of n , and $\alpha = 2\pi$ returns the identity. The family is therefore 2π periodic in α and unitary, since it multiplies the coefficients of an orthonormal expansion by unimodular numbers.

Namias' first observation is that (5.58) can be freed from the basis and written as a differential operator. Differentiating (5.58) with respect to the order,

$$\frac{\partial F_\alpha}{\partial \alpha}(x) = -i \sum_{n=0}^{\infty} n c_n e^{-in\alpha} \psi_n(x), \quad (5.59)$$

we are left with the index n standing inside the sum, which is inconvenient because n is not an operator acting on x . The remedy is the differential equation of the Hermite functions. Substituting (5.54) into the Hermite equation $H_n'' - 2xH_n' + 2nH_n = 0$ one obtains, after the elementary computation of the two derivatives of the Gaussian factor,

$$\psi_n''(x) + (2n + 1 - x^2) \psi_n(x) = 0, \quad (5.60)$$

which is precisely the stationary Schrödinger equation of the quantum harmonic oscillator in natural units. Solving (5.60) for the index,

$$n \psi_n(x) = \frac{1}{2} [x^2 \psi_n(x) - \psi_n''(x) - \psi_n(x)], \quad (5.61)$$

and inserting (5.61) into (5.59),

$$\begin{aligned} \frac{\partial F_\alpha}{\partial \alpha}(x) &= -\frac{i}{2} \sum_{n=0}^{\infty} c_n e^{-in\alpha} [x^2 \psi_n(x) - \psi_n''(x) - \psi_n(x)] = \\ &= -\frac{i}{2} \left[x^2 \sum_n c_n e^{-in\alpha} \psi_n(x) - \frac{\partial^2}{\partial x^2} \sum_n c_n e^{-in\alpha} \psi_n(x) - \sum_n c_n e^{-in\alpha} \psi_n(x) \right], \end{aligned}$$

where the index n has disappeared entirely and each of the three sums is again F_α by (5.58). We therefore arrive at the differential formulation of the fractional Fourier transformation,

$$i \frac{\partial F_\alpha}{\partial \alpha} = \frac{1}{2} \left[-\frac{\partial^2}{\partial x^2} + x^2 - 1 \right] F_\alpha, \quad F_\alpha|_{\alpha=0} = f. \quad (5.62)$$

Equation (5.62) is Namias' definition of the fractional Fourier transformation as a differential operator. It is worth reading it twice. It states that the transform of order α of a given function is obtained by letting that function evolve, according to the Schrödinger equation of the harmonic oscillator, for a *time* equal to α , the constant -1 merely removing the zero point energy so that the ground state does not accumulate a phase. In operator language, introducing the oscillator Hamiltonian

$$\hat{H} = \frac{1}{2} \left(-\frac{\partial^2}{\partial x^2} + x^2 \right), \quad \hat{H}\psi_n = \left(n + \frac{1}{2} \right) \psi_n, \quad (5.63)$$

the solution of (5.62) is written at once as the exponential of the generator,

$$\mathcal{F}_\alpha = \exp \left[-i\alpha \left(\hat{H} - \frac{1}{2} \right) \right] = \exp \left[\frac{i\alpha}{2} \left(\frac{\partial^2}{\partial x^2} - x^2 + 1 \right) \right], \quad (5.64)$$

which reproduces (5.56) immediately, since $\exp[-i\alpha(\hat{H} - \frac{1}{2})]\psi_n = e^{-i\alpha n}\psi_n$. This identification is the reason the fractional Fourier transformation appears wherever a harmonic oscillator appears, and it explains in retrospect why Condon had already met the same family in 1937 while studying continuous groups of functional transformations (Condon, 1937).

Two structural consequences follow immediately from (5.64). First, since the exponentials of a single generator commute and add their arguments,

$$\mathcal{F}_\alpha \mathcal{F}_\beta = \mathcal{F}_{\alpha+\beta}, \quad \mathcal{F}_\alpha^{-1} = \mathcal{F}_{-\alpha}, \quad (5.65)$$

so the transformations form a commutative one parameter group. Second, since $\hat{H}$ is self adjoint, $\mathcal{F}_\alpha$ is unitary, and Plancherel's theorem extends to every order,

$$\int_{-\infty}^{\infty} |\mathcal{F}_\alpha[f](x)|^2 dx = \int_{-\infty}^{\infty} |f(x)|^2 dx, \quad (5.66)$$

which in the optical reading of the next subsection is nothing other than the conservation of the energy carried by the field.

5.7.2.3 The integral form of the transformation

The differential form (5.62) is compact and conceptually transparent, but for computation one wants a kernel. Since the transformation is linear and bounded, it must be representable as

$$\mathcal{F}_\alpha[f](x) = \int_{-\infty}^{\infty} K_\alpha(x, y) f(y) dy, \quad (5.67)$$

and inserting the expansion (5.57) into (5.58) and exchanging the sum with the integral, the kernel is read off as the bilinear spectral sum

$$K_\alpha(x, y) = \sum_{n=0}^{\infty} e^{-in\alpha} \psi_n(x) \psi_n(y). \quad (5.68)$$

Substituting the explicit Hermite functions (5.54),

$$K_\alpha(x, y) = \frac{e^{-(x^2+y^2)/2}}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{H_n(x) H_n(y)}{2^n n!} e^{-in\alpha}, \quad (5.69)$$

we are left with a bilinear generating function of Hermite polynomials, which is exactly the object Mehler computed in 1866 (Mehler, 1866),

$$\sum_{n=0}^{\infty} \frac{H_n(x) H_n(y)}{2^n n!} t^n = \frac{1}{\sqrt{1-t^2}} \exp \left[\frac{2xyt - (x^2 + y^2)t^2}{1-t^2} \right], \quad |t| < 1. \quad (5.70)$$

We must set $t = e^{-i\alpha}$, which lies on the boundary of the disc of convergence, so (5.70) is to be understood as the limit $t \rightarrow e^{-i\alpha}$ taken radially from inside the disc, a limit that exists in the sense of distributions and that is legitimate for every α that is not an integer multiple of π . Let us now carry out the algebra step by step. The denominator factorises as

$$1 - e^{-2i\alpha} = e^{-i\alpha} (e^{i\alpha} - e^{-i\alpha}) = e^{-i\alpha} 2i \sin \alpha, \quad (5.71)$$

so that the crossed term of the exponent of (5.70) becomes

$$\frac{2xy e^{-i\alpha}}{e^{-i\alpha} 2i \sin \alpha} = \frac{xy}{i \sin \alpha} = -\frac{ixy}{\sin \alpha} = -ixy \csc \alpha, \quad (5.72)$$

where the two exponentials in red cancel. For the quadratic term we must not forget the Gaussian prefactor of (5.69), so we collect both contributions,

$$\begin{aligned} & -\frac{x^2 + y^2}{2} - \frac{(x^2 + y^2) e^{-2i\alpha}}{e^{-i\alpha} 2i \sin \alpha} = -\frac{x^2 + y^2}{2} - \frac{(x^2 + y^2) e^{-i\alpha}}{2i \sin \alpha} = \\ & -\frac{x^2 + y^2}{2} \left[1 + \frac{e^{-i\alpha}}{i \sin \alpha} \right] = -\frac{x^2 + y^2}{2} \cdot \frac{i \sin \alpha + e^{-i\alpha}}{i \sin \alpha}, \end{aligned}$$

and the numerator of this last fraction collapses beautifully,

$$i \sin \alpha + e^{-i\alpha} = i \sin \alpha + \cos \alpha - i \sin \alpha = \cos \alpha, \quad (5.73)$$

so that, using $-1/i = i$,

$$-\frac{x^2 + y^2}{2} \cdot \frac{\cos \alpha}{i \sin \alpha} = \frac{i}{2} (x^2 + y^2) \cot \alpha. \quad (5.74)$$

There remains the normalising constant. Using (5.71) once more,

$$\frac{1}{\sqrt{\pi}} \cdot \frac{1}{\sqrt{1 - e^{-2i\alpha}}} = \frac{1}{\sqrt{\pi}} \cdot \frac{e^{i\alpha/2}}{\sqrt{2i \sin \alpha}} = \frac{e^{i(\frac{\alpha}{2} - \frac{\pi}{4})}}{\sqrt{2\pi \sin \alpha}}, \quad (5.75)$$

since $1/\sqrt{i} = e^{-i\pi/4}$. This constant is more often written in the compact form that Namias used, and the two agree because

$$1 - i \cot \alpha = \frac{\sin \alpha - i \cos \alpha}{\sin \alpha} = \frac{-i(\cos \alpha + i \sin \alpha)}{\sin \alpha} = \frac{e^{-i\pi/2} e^{i\alpha}}{\sin \alpha}, \quad (5.76)$$

whose square root divided by $\sqrt{2\pi}$ is precisely (5.75). Collecting (5.72), (5.74) and (5.75) we obtain the integral form of the transformation,

$$\boxed{\mathcal{F}_\alpha[f](x) = \sqrt{\frac{1 - i \cot \alpha}{2\pi}} \int_{-\infty}^{\infty} \exp \left\{ \frac{i}{2} (x^2 + y^2) \cot \alpha - ixy \csc \alpha \right\} f(y) dy.} \quad (5.77)$$

Let us verify that (5.77) does what it should. For $\alpha = \pi/2$ we have $\cot \alpha = 0$ and $\csc \alpha = 1$, so the constant reduces to $1/\sqrt{2\pi}$ and the kernel to e^{-ixy} , recovering the ordinary transformation (5.48) exactly. For $\alpha \rightarrow 0$ the coefficient $\cot \alpha$ diverges and the kernel oscillates infinitely fast everywhere except on the diagonal, where the exponent $\frac{i}{2}(x - y)^2 \cot \alpha$ vanishes, and a standard stationary phase argument shows that $K_\alpha \rightarrow \delta(x - y)$, which is the identity. For $\alpha \rightarrow \pi$ the same argument applied to $\frac{i}{2}(x + y)^2 \cot \alpha$ gives $K_\pi(x, y) = \delta(x + y)$, the parity operator. The three degenerate orders $\alpha = 0, \pi, 2\pi$, at which the closed formula (5.77) is meaningless because $\csc \alpha$ blows up, are therefore perfectly well defined in the spectral form (5.68), a discrepancy that we shall have to come back to.

5.7.2.4 The inverse transformation and the index law

The inverse is obtained without any new computation, because (5.68) makes the composition of two transformations transparent. Consider the kernel of the product $\mathcal{F}_\alpha \mathcal{F}_\beta$,

$$\begin{aligned} \int_{-\infty}^{\infty} K_\alpha(x, s) K_\beta(s, y) ds &= \int_{-\infty}^{\infty} \left[\sum_n e^{-in\alpha} \psi_n(x) \psi_n(s) \right] \left[\sum_m e^{-im\beta} \psi_m(s) \psi_m(y) \right] ds = \\ &= \sum_n \sum_m e^{-in\alpha} e^{-im\beta} \psi_n(x) \psi_m(y) \underbrace{\int_{-\infty}^{\infty} \psi_n(s) \psi_m(s) ds}_{\delta_{nm}} = \sum_n e^{-in(\alpha+\beta)} \psi_n(x) \psi_n(y), \end{aligned}$$

where the orthonormality (5.54) has collapsed the double sum into a single one, and the last expression is by definition $K_{\alpha+\beta}(x, y)$. We have therefore proved the **index law**

$$\mathcal{F}_\alpha \mathcal{F}_\beta = \mathcal{F}_{\alpha+\beta}, \quad (5.78)$$

which is the property that justifies calling α an order, since orders add under composition exactly as exponents do. Setting $\beta = -\alpha$ and using completeness,

$$K_0(x, y) = \sum_{n=0}^{\infty} \psi_n(x) \psi_n(y) = \delta(x - y), \quad (5.79)$$

we conclude that $\mathcal{F}_\alpha \mathcal{F}_{-\alpha} = \mathcal{I}$, that is,

$$\mathcal{F}_\alpha^{-1} = \mathcal{F}_{-\alpha}, \quad (5.80)$$

so that the inverse transformation is the transformation of the opposite order. Explicitly, replacing α by $-\alpha$ in (5.77) and using $\cot(-\alpha) = -\cot \alpha$ and $\csc(-\alpha) = -\csc \alpha$,

$$f(y) = \sqrt{\frac{1 + i \cot \alpha}{2\pi}} \int_{-\infty}^{\infty} \exp \left\{ -\frac{i}{2} (x^2 + y^2) \cot \alpha + i xy \csc \alpha \right\} F_\alpha(x) dx, \quad (5.81)$$

and comparing (5.81) with (5.77) we see that $K_{-\alpha}(y, x) = \overline{K_\alpha(x, y)}$, which is precisely the statement that $\mathcal{F}_\alpha$ is unitary, in agreement with (5.66).

5.7.2.5 The regularisation of McBride and Kerr

The derivation we have just carried out is Namias' derivation, and as a piece of physics it is complete. As a piece of analysis it has three gaps, all of which were identified and repaired by Adam McBride and Fiona Kerr in 1987 (McBride and Kerr, 1987). Since the modern optical literature uses their version without always saying so, it is worth stating what they changed.

The first gap is the **branch of the square root**. The constant in (5.77) is written as $\sqrt{1 - i \cot \alpha}$, and a square root of a complex number is defined only up to a sign. The spectral sum (5.68) has no such ambiguity, so the sign must be fixed by matching the two, and the correct determination is the one already contained in (5.75), which written without ambiguity reads

$$C_\alpha = \frac{\exp \left[i \left(\frac{\alpha}{2} - \frac{\pi}{4} \operatorname{sgn}(\sin \alpha) \right) \right]}{\sqrt{2\pi |\sin \alpha|}}. \quad (5.82)$$

The discrete factor $\operatorname{sgn}(\sin \alpha)$ is not cosmetic. If one chooses the principal branch naively and composes two transformations whose orders add up to more than π , the index law (5.78) fails by a factor -1 . This is the elementary manifestation of a well known phenomenon, namely that the

group of transformations is not represented faithfully by the kernels but by a double cover of them, the extra sign being what is called the Maslov index in the theory of canonical transformations (Moshinsky and Quesne, 1971; Wolf, 1979).

The second gap is the **domain and the convergence of the integral**. For a general $f \in L^2(\mathbb{R})$ the integral in (5.77) does not converge absolutely, because the modulus of the kernel is constant in y and the integrand therefore decays no faster than f itself. McBride and Kerr define the transformation first on the Schwartz space $\mathcal{S}$ of rapidly decreasing functions, where absolute convergence is guaranteed, prove that $\mathcal{F}_\alpha$ maps $\mathcal{S}$ onto $\mathcal{S}$ homeomorphically, and only then extend it by duality to the space $\mathcal{S}'$ of tempered distributions. Their technical device for enlarging the domain is to trade one power of decay in the integration variable for one derivative in the observation variable, replacing (5.77) by an expression of the form

$$\mathcal{F}_\alpha[f](x) = C_\alpha e^{\frac{i}{2}x^2 \cot \alpha} \frac{d}{dx} \int_{-\infty}^{\infty} \frac{e^{-ixy \csc \alpha}}{-iy \csc \alpha} e^{\frac{i}{2}y^2 \cot \alpha} f(y) dy, \quad (5.83)$$

in which the extra factor $1/y$ makes the integral absolutely convergent for functions that are not themselves integrable, while the derivative taken outside the integral restores the original kernel. Once the resulting operator is shown to be continuous on $\mathcal{S}$, the transposition to $\mathcal{S}'$ is automatic, and the degenerate orders $\alpha = n\pi$ acquire their correct meaning as the distributional kernels $\delta(x \mp y)$ instead of appearing as divergent integrals.

The third gap is that Namias verifies the index law (5.78) on the eigenfunction expansion, which is a formal manipulation of a doubly infinite sum. McBride and Kerr prove it as a theorem for all real α and β on a stated domain, and thereby establish that

$$\{\mathcal{F}_\alpha\}_{\alpha \in \mathbb{R}}, \quad \mathcal{F}_0 = \mathcal{I}, \quad \mathcal{F}_{\alpha+2\pi} = \mathcal{F}_\alpha, \quad (5.84)$$

is a genuine continuous one parameter group of homeomorphisms of $\mathcal{S}$, unitary on $L^2(\mathbb{R})$. Kerr afterwards developed the distributional theory in detail (Kerr, 1988a) and the L^2 theory together with applications to differential equations (Kerr, 1988b). With these three repairs, and only with them, is one entitled to manipulate $\mathcal{F}_\alpha$ as an algebraic object, which is what we shall do freely in the next subsection.

5.7.2.6 Why the construction is useful

The two faces of the transformation, the differential and the integral one, are useful in different circumstances, and it is worth separating them.

The **differential form** (5.62) is useful whenever the harmonic oscillator is hiding in the problem, which happens far more often than one would expect. Namias' own motivation was quantum mechanical (Namias, 1980): since $\mathcal{F}_\alpha$ is the evolution operator of the oscillator, the fractional transform of an initial wave packet *is* the wave packet at a later time, so that problems of time evolution become problems of choosing an order, and the well known revival of a wave packet after a

full period is simply the 2π periodicity (5.84). More generally, since $\mathcal{F}_\alpha$ interchanges the operators x and $-i\partial_x$ by a rotation of angle α , a differential equation with polynomial coefficients can frequently be rotated into a simpler one, which is the technique Kerr systematised (Kerr, 1988b). The same generator governs the Landau problem, the paraxial wave equation in a quadratic index medium, and the linearised dynamics of any system near a stable equilibrium, so the differential form transfers immediately to all of them.

The **integral form** (5.77) is useful whenever one needs to compute rather than to reason. In signal processing it provides a continuum of representations interpolating between the time domain and the frequency domain, and since a linear chirp is a plane wave in some fractional domain, a chirped signal buried in noise becomes a sharp peak at the right order, which is the basis of chirp detection in radar and sonar and of filtering in fractional domains (Almeida, 1994; Sejdić et al., 2011). Lohmann's identification of the order with a rotation angle of the Wigner distribution (Lohmann, 1993), later placed on a rigorous footing by Mustard (1996), gives the whole construction a geometrical reading in phase space, and the discrete version of Candan, Kutay and Ozaktas (2000) makes it computable with the same logarithmic cost as the ordinary discrete transformation (Cooley and Tukey, 1965). In optics the integral form is what allows a fractional transformation to be built in the laboratory, either with a graded index medium of the appropriate length (Mendlovic and Ozaktas, 1993; Ozaktas and Mendlovic, 1993) or with a pair of lenses (Lohmann, 1993), and it underlies applications to beam propagation, phase retrieval, optical tomography, filtering and image encryption (Ozaktas et al., 2001).

For our purposes, however, the decisive application is none of these. It is the fact, proved in the next subsection, that the Fresnel integral (5.34) is itself a fractional Fourier transformation. The consequence is that the whole of paraxial propagation, which in the plane to plane description of the previous sections looked like a family of integrals labelled by a distance and composing in a complicated way, becomes a one parameter group labelled by an angle and composing by simple addition.

5.7.3 The role of Fourier analysis in the theory of diffraction

We are now in a position to prove the statement announced at the end of the section on the Fraunhofer approximation. The result is due to Pellat-Finet (Pellat-Finet, 1994; Pellat-Finet and Bonnet, 1994) and it may be stated as follows: *Fresnel diffraction between two spherical caps of equal radius is exactly a fractional Fourier transformation, whose order is fixed by the ratio of the propagation distance to the radius of the caps.* Since the transformation acts on two transverse variables we shall use the two dimensional kernel, which is separable and therefore the product of two copies of (5.77),

$$\mathcal{F}_\alpha[g](\vec{v}) = \frac{1 - i \cot \alpha}{2\pi} \int_{\mathbb{R}^2} \exp \left\{ \frac{i}{2} (u^2 + v^2) \cot \alpha - i \vec{u} \cdot \vec{v} \csc \alpha \right\} g(\vec{u}) d^2u, \quad (5.85)$$

where $u^2 = \|\vec{u}\|^2$ and $v^2 = \|\vec{v}\|^2$, the constant being the square of the one dimensional constant because the two dimensional transformation is the tensor product of two identical one dimensional ones.

5.7.3.1 Why a plane cannot be the right reference surface

It is instructive to begin by showing that the naive attempt fails, because the failure tells us exactly what to repair. Expanding the square in the Fresnel integral (5.34),

$$U(\vec{r}') = \frac{i e^{ikz}}{\lambda z} \int_{\mathbb{R}^2} U(\vec{r}) \exp \left[\frac{ik}{2z} (r^2 + r'^2) \right] \exp \left[-\frac{ik}{z} \vec{r} \cdot \vec{r}' \right] d^2r, \quad (5.86)$$

where we used $\|\vec{r} - \vec{r}'\|^2 = r^2 + r'^2 - 2\vec{r} \cdot \vec{r}'$, we see that (5.86) has exactly the algebraic shape of (5.85): two quadratic terms with a common coefficient and one crossed term. Introduce therefore the reduced, dimensionless variables

$$\vec{u} = \frac{\vec{r}}{\omega}, \quad \vec{v} = \frac{\vec{r}'}{\omega}, \quad (5.87)$$

with ω a scale length to be determined, and identify the coefficients term by term,

$$\frac{k}{2z} \omega^2 = \frac{\cot \alpha}{2} \quad (\text{quadratic terms}), \quad (5.88)$$

$$\frac{k}{z} \omega^2 = \csc \alpha \quad (\text{crossed term}). \quad (5.89)$$

Dividing (5.88) by (5.89) the unknown scale ω disappears and we are left with

$$\frac{1}{2} = \frac{\cot \alpha}{2 \csc \alpha} = \frac{\cos \alpha}{2} \implies \cos \alpha = 1,$$

that is $\alpha = 0$, which is the identity and for which $\csc \alpha$ is infinite, so that (5.89) cannot be satisfied by any finite ω . The conclusion is unambiguous.

Theorem 7 *Fresnel diffraction between two planes is never a fractional Fourier transformation, whatever the order. In the language of the group of linear canonical transformations it is a parabolic element, while the fractional Fourier transformations are the elliptic ones.*

The obstruction is entirely due to the relative weight of the quadratic and the crossed terms, which in (5.86) is fixed at the value one half whereas the kernel (5.85) demands $\cos \alpha/2$. Since the crossed term is the only one that couples the two planes, and is therefore the one we cannot touch, the only freedom we have is to modify the quadratic terms. That is exactly what referring the field to a curved surface does.

5.7.3.2 Reference spheres and the fractional order

Let the emitting plane be at the origin and the receiving plane at the distance z along the axis. Consider a spherical cap tangent to a transverse plane at the axis and whose centre of curvature lies on the axis at the signed distance c from the vertex, counted positive in the direction of propagation. In the paraxial regime the point of the cap at the transverse position $\vec{r}$ is displaced axially from the tangent plane by the sagitta

$$\delta(\vec{r}) = c - \text{sgn}(c) \sqrt{c^2 - r^2} = \frac{r^2}{2c} + \mathcal{O}\left(\frac{r^4}{c^3}\right), \quad (5.90)$$

which is the same expansion we performed in (5.33). Since a field travelling in the positive direction acquires the factor $e^{ik\delta}$ over an axial displacement δ , the field referred to the cap and the field on the tangent plane are related by

$$u_{\text{cap}}(\vec{r}) = U(\vec{r}) \exp\left(\frac{ik r^2}{2c}\right). \quad (5.91)$$

Let now $\mathcal{A}$ be a cap tangent to the emitting plane with centre of curvature at c_1 , and $\mathcal{B}$ a cap tangent to the receiving plane with centre of curvature at the signed distance c_2 from its own vertex. Using (5.91) on both sides,

$$U(\vec{r}) = u_{\mathcal{A}}(\vec{r}) \exp\left(-\frac{ik r^2}{2c_1}\right), \quad u_{\mathcal{B}}(\vec{r}') = U(\vec{r}') \exp\left(\frac{ik r'^2}{2c_2}\right), \quad (5.92)$$

and substituting both relations into (5.86),

$$\begin{aligned} u_{\mathcal{B}}(\vec{r}') &= \frac{i e^{ikz}}{\lambda z} e^{\frac{ikr'^2}{2c_2}} \int u_{\mathcal{A}}(\vec{r}) e^{-\frac{ikr^2}{2c_1}} \exp\left[\frac{ik}{2z} (r^2 + r'^2)\right] e^{-\frac{ik}{z} \vec{r} \cdot \vec{r}'} d^2r = \\ &= \frac{i e^{ikz}}{\lambda z} \exp\left[\frac{ikr'^2}{2} \left(\frac{1}{z} + \frac{1}{c_2}\right)\right] \int u_{\mathcal{A}}(\vec{r}) \exp\left[\frac{ikr^2}{2} \left(\frac{1}{z} - \frac{1}{c_1}\right)\right] e^{-\frac{ik}{z} \vec{r} \cdot \vec{r}'} d^2r, \end{aligned}$$

where we have merely collected the coefficients of r^2 and of r'^2 . The kernel (5.85) requires the two quadratic coefficients to be equal, which imposes

$$\frac{1}{z} - \frac{1}{c_1} = \frac{1}{z} + \frac{1}{c_2} \implies \frac{\textcolor{red}{1}}{\textcolor{red}{z}} - \frac{1}{c_1} - \frac{\textcolor{red}{1}}{\textcolor{red}{z}} - \frac{1}{c_2} = 0 \implies c_2 = -c_1, \quad (5.93)$$

the two terms in red cancelling identically. We therefore set

$$c_1 = R, \quad c_2 = -R, \quad (5.94)$$

which describes a perfectly definite geometry: *two spherical caps of the same radius R , the emitting one having its centre of curvature downstream at the abscissa R , and the receiving one, whose vertex*

sits at the abscissa z , having its centre of curvature upstream at the abscissa $z - R$. These are the cardinal spheres of radius R associated with the propagation, and their geometry is the one drawn in Figure 5.6.




Figure 5.6: The two cardinal spheres of radius R associated with a propagation over the distance z . The emitting cap $\mathcal{A}$ is tangent to its plane at V_A with its centre of curvature downstream at C_A , of abscissa R , and the receiving cap $\mathcal{B}$ is tangent to its plane at V_B with its centre upstream at C_B , of abscissa $z - R$, so that the two caps bow towards each other. The quantity δ is the sagitta (5.90). The confocal case of Figure 5.4 is recovered when $R = z$, since then C_A falls on V_B and C_B on V_A .

With (5.94) the two quadratic coefficients collapse into the single quantity

$$Q = \frac{k}{2} \left(\frac{1}{z} - \frac{1}{R} \right), \quad (5.95)$$

and the propagation reads

$$u_B(\vec{r}') = \frac{i e^{ikz}}{\lambda z} \int_{\mathbb{R}^2} u_A(\vec{r}) \exp [iQ (r^2 + r'^2)] \exp \left[-\frac{ik}{z} \vec{r} \cdot \vec{r}' \right] d^2r, \quad (5.96)$$

which now has exactly the structure of (5.85) with independent quadratic and crossed coefficients. Repeating the identification of the previous subsection with the reduced variables (5.87),

$$Q \omega^2 = \frac{\cot \alpha}{2} \implies \cot \alpha = k \omega^2 \left(\frac{1}{z} - \frac{1}{R} \right), \quad (5.97)$$

$$\frac{k}{z} \omega^2 = \csc \alpha, \quad (5.98)$$

and dividing (5.97) by (5.98) the scale ω cancels once again,

$$\cos \alpha = \frac{\cot \alpha}{\csc \alpha} = \frac{\textcolor{red}{k} \omega^2 \left(\frac{1}{z} - \frac{1}{R} \right)}{\textcolor{red}{k} \omega^2 \frac{1}{z}} = z \left(\frac{1}{z} - \frac{1}{R} \right),$$

so that we obtain the central result

$$\boxed{\cos \alpha = 1 - \frac{z}{R}}. \quad (5.99)$$

This time there is no contradiction, because the radius R of the reference caps is at our disposal. The scale length follows from (5.98),

$$\omega^2 = \frac{z}{k \sin \alpha} = \frac{\lambda z}{2\pi \sin \alpha}, \quad (5.100)$$

and, eliminating α by means of (5.99) and of $\sin^2 \alpha = 1 - \cos^2 \alpha$,

$$\sin^2 \alpha = \textcolor{red}{1} - \left(\textcolor{red}{1} - \frac{2z}{R} + \frac{z^2}{R^2} \right) = \frac{z(2R - z)}{R^2} \implies \sin \alpha = \frac{\sqrt{z(2R - z)}}{R},$$

the two units in red cancelling, so that the scale length is expressed entirely in terms of the geometry,

$$\omega^2 = \frac{\lambda R}{2\pi} \sqrt{\frac{z}{2R - z}}. \quad (5.101)$$

5.7.3.3 The normalising constant and the final result

It remains to compare the constants. Changing variables in (5.96) according to (5.87), so that $d^2 r = \omega^2 d^2 u$,

$$u_B(\omega \vec{v}) = \frac{i e^{ikz} \omega^2}{\lambda z} \int_{\mathbb{R}^2} u_A(\omega \vec{u}) \exp \left[\frac{i}{2} (u^2 + v^2) \cot \alpha - i \vec{u} \cdot \vec{v} \csc \alpha \right] d^2 u, \quad (5.102)$$

and using (5.100) the prefactor simplifies to

$$\frac{i\omega^2}{\lambda z} = \frac{i}{\lambda z} \cdot \frac{\lambda z}{2\pi \sin \alpha} = \frac{i}{2\pi \sin \alpha}. \quad (5.103)$$

On the other hand the constant of (5.85), rewritten by means of the identity (5.76), is

$$\frac{1 - i \cot \alpha}{2\pi} = \frac{1}{2\pi} \cdot \frac{e^{-i\pi/2} e^{i\alpha}}{\sin \alpha} = \frac{-i e^{i\alpha}}{2\pi \sin \alpha}. \quad (5.104)$$

Dividing (5.103) by (5.104),

$$\frac{\frac{i}{2\pi \sin \alpha}}{\frac{-i e^{i\alpha}}{2\pi \sin \alpha}} = \frac{i}{-i} e^{-i\alpha} = -e^{-i\alpha},$$

and therefore, collecting everything, we have proved the theorem.

Theorem 8 (*Fresnel diffraction as a fractional Fourier transformation*) *Let the field be referred to two cardinal spherical caps of radius R separated by the distance z along the axis, in the sense of (5.94), and let the transverse coordinates be reduced by the scale ω of (5.101). Then the Fresnel propagation (5.34) is the fractional Fourier transformation*

$$u_{\mathcal{B}}(\omega \vec{v}) = e^{i(kz + \pi - \alpha)} (\mathcal{F}_{\alpha} u_{\mathcal{A}})(\vec{v}), \quad (5.105)$$

whose order is given by $\cos \alpha = 1 - z/R$.

A remark on the residual phase is in order. The additive π in (5.105) is inherited from the sign of the constant written in front of the Fresnel integral (5.34). With the normalisation $1/(i\lambda z)$, which is the one for which the Fresnel kernel tends to $\delta(\vec{r} - \vec{r}')$ as $z \rightarrow 0$, the very same computation gives the residual factor $e^{-i\alpha}$ with no additive constant, and the order α is then exactly the accumulated Gouy phase, a point to which we shall return.

Let us test (5.105) on the three cases we can check independently.

1. $z = 0$. Equation (5.99) gives $\cos \alpha = 1$, hence $\alpha = 0$ and $\mathcal{F}_0 = \mathcal{I}$: the field on the receiving cap is the field on the emitting cap, as it must be.
2. $z = R$. Now $\cos \alpha = 0$, hence $\alpha = \pi/2$ and the transformation is the ordinary Fourier transformation. This is not a coincidence: when $z = R$ the emitting cap has its centre at the abscissa $R = z$, which is the vertex of the receiving cap, and the receiving cap has its centre at $z - R = 0$, which is the vertex of the emitting cap. The two caps are therefore *the confocal cardinal spheres* of Figure 5.5, and we recover exactly the Fraunhofer result (5.45) of the previous section. The far field description is the particular case $\alpha = \pi/2$ of the present one.

3. $z = 2R$. Here $\cos \alpha = -1$, hence $\alpha = \pi$ and $\mathcal{F}_\pi = \mathcal{P}$: the field on the receiving cap is the field on the emitting cap turned upside down. Two consecutive Fourier transformations invert the image, and the geometry says the same thing.

The gain in economy is considerable. In the plane to plane description, chaining two propagations of distances z_1 and z_2 requires composing two Fresnel integrals, and the composite kernel must be recomputed each time. In the present description the composition is the index law (5.78),

$$\mathcal{F}_{\alpha_2} \mathcal{F}_{\alpha_1} = \mathcal{F}_{\alpha_1 + \alpha_2}, \quad (5.106)$$

so that propagation is described by the addition of angles. The distances z do not add in any useful way, but the orders do, and this is the precise sense in which α , and not z , is the natural parameter of free paraxial propagation.

5.7.3.4 Physical meaning of a real order

When α is real, (5.99) requires $|1 - z/R| \leq 1$, that is

$$0 \leq z \leq 2R, \quad (5.107)$$

and the order sweeps the interval $[0, \pi]$ monotonically as the observation cap recedes. The reading of α is then completely transparent, and there are three equivalent ways of stating it.

As a degree of diffraction. The order measures how far the field has travelled from the near field towards the far field, on a scale in which $\alpha = 0$ is the aperture itself and $\alpha = \pi/2$ is its Fraunhofer pattern. A field observed at $\alpha = \pi/4$ is halfway between its own distribution and its Fourier transform, in a sense that is not metaphorical but exact, since applying the same transformation twice produces the Fourier transform. The continuum of intermediate regimes that the Fresnel approximation describes only implicitly is here labelled by a single number.

As a rotation of phase space. By the theorem of Lohmann (1993), later established rigorously by Mustard (1996), the transformation $\mathcal{F}_\alpha$ acts on the Wigner distribution of the field simply by rotating the plane of the transverse position and the transverse spatial frequency by the angle α . Free propagation between cardinal spheres is therefore a rigid rotation of phase space, which is why the orders add: rotations compose by adding angles. This also explains geometrically why plane to plane propagation is not a fractional transformation, since in the plane description propagation is a shear of phase space rather than a rotation.

As a Gouy phase. The residual factor $e^{-i\alpha}$ of the footnote to (5.105) is not an accident. If instead of arbitrary caps one chooses as reference surfaces the wavefronts of a Gaussian beam of Rayleigh range z_R , the order that comes out of the same computation is

$$\alpha = \arctan\left(\frac{z}{z_R}\right), \quad (5.108)$$

which is precisely the accumulated Gouy phase of the beam (Ozaktas and Mendlovic, 1995). The identification is satisfying, since (5.108) tends to $\pi/2$ as $z \rightarrow \infty$, recovering once more the statement that the far field is the Fourier transform of the near field, and it shows that the mysterious extra phase of a focused beam is nothing but the order of the fractional transformation the beam has undergone.

One point must be emphasised, because it is a frequent source of confusion. *The order is not a property of the propagation alone.* It is a property of the pair formed by the propagation and the two reference caps, since (5.99) involves both z and R . Given a propagation distance z , any order whatsoever can be obtained by choosing

$$R = \frac{z}{1 - \cos \alpha}, \quad (5.109)$$

and conversely the same pair of caps assigns different orders to different distances. What is intrinsic is not α but the group structure: whatever reference caps are chosen, propagation is represented by an element of the same one parameter group.

5.7.3.5 Physical meaning of a complex order

The condition (5.107) is a real restriction, and it is natural to ask what happens when it is violated. If the radius R of the caps is held fixed while the propagation distance exceeds $2R$, or if the propagation is run backwards so that $z < 0$, then $\cos \alpha$ leaves the interval $[-1, 1]$ and the order becomes complex. Writing $\alpha = \alpha_r + i\alpha_i$ there are two cases.

For $\cos \alpha > 1$, which corresponds to $z < 0$ with $R > 0$, the order is purely imaginary, $\alpha = i\beta$ with

$$\cosh \beta = 1 - \frac{z}{R}. \quad (5.110)$$

For $\cos \alpha < -1$, which corresponds to $z > 2R$, the order is $\alpha = \pi + i\beta$ with $\cosh \beta = z/R - 1$, the additive π carrying the inversion of the image that we already met at $z = 2R$.

The consequence for the kernel is immediate and physically eloquent. Taking the first case and using $\cot(i\beta) = -i \coth \beta$ and $\csc(i\beta) = -i \operatorname{csch} \beta$, the exponent of (5.85) becomes

$$\begin{aligned} & \frac{i}{2} (u^2 + v^2) (-i \coth \beta) - i \vec{u} \cdot \vec{v} (-i \operatorname{csch} \beta) = \\ & \frac{1}{2} (u^2 + v^2) \coth \beta - \vec{u} \cdot \vec{v} \operatorname{csch} \beta, \end{aligned}$$

which is *entirely real*. The kernel has ceased to be a pure chirp and has become a real Gaussian, and since

$$\coth \beta - \operatorname{csch} \beta = \frac{\cosh \beta - 1}{\sinh \beta} > 0 \quad (\beta > 0),$$

the associated quadratic form is positive definite, so the kernel grows rather than oscillates. This is the analytic signature of an ill posed problem, and it is exactly what one should expect, since $z < 0$ is back propagation and back propagation amplifies the high spatial frequency content of the field without bound. The complex order does not merely describe the difficulty, it quantifies it: the imaginary part β is the exponent of the Gaussian amplification.

More generally, a complex order admits three complementary physical readings.

1. **Group theoretically**, a real order corresponds to an elliptic element of the group of linear canonical transformations, that is, to a rotation of phase space, while a complex order corresponds to an element that also contains a squeeze, that is, a magnification along one direction of phase space and a compression along the other. A complex order therefore announces that the transformation is a rotation composed with a scaling, and the imaginary part measures the scaling.
2. **For Gaussian beams**, the wavefront curvature is not a real number but the complex parameter q defined by $q^{-1} = R^{-1} - i\lambda/(\pi w^2)$, where w is the beam radius. Substituting q for R in (5.99) gives a complex order automatically, whose real part is the phase space rotation and whose imaginary part carries the Gaussian envelope of the beam. Complex order fractional transformations are, for this reason, the natural language of Gaussian beam propagation, and they were realised optically by Shih (1995b) and by Bernardo and Soares (1996).
3. **For apodised systems**, a Gaussian aperture e^{-r^2/a^2} placed in the beam contributes exactly a real Gaussian factor of the kind that a complex order produces. The imaginary part of the order then measures the attenuation of the large $|\vec{u}|$ content of the field, that is, a Gaussian low pass filtering in the fractional domain, which in the image is a loss of resolution. A complex order is, in this reading, the bookkeeping device for soft apertures and for losses.

Finally, it should be stressed that a complex order never signals a physical impossibility. Since (5.109) shows that the order depends on the reference caps, any forward propagation with $z > 0$ can be described with a real order simply by choosing $R \geq z/2$. What a complex order signals is that the reference surfaces chosen are not the ones with respect to which the propagation is a pure rotation. The limiting case of this statement is the plane, for which $R \rightarrow \infty$ gives $\cos \alpha \rightarrow 1$ and, by (5.100), $\omega \rightarrow \infty$: the transformation degenerates, the rotation angle collapses to zero while the scale diverges, and we recover the theorem of the first subsection that plane to plane Fresnel propagation lies outside the fractional Fourier family. The whole content of Pellat-Finet's result is that the family is recovered as soon as one agrees to look at the field on a sphere.

Chapter 6

Theory of Coherence

*Coherence is the thread that weaves order out of
the chaos of random light.*

“Time advances in the direction in which correlations increase.”
– attributed to Ludwig Boltzmann

6.1 A historical context of coherence theory

At its heart, coherence asks a single, almost childlike question: how much does one thing resemble another, and does that resemblance survive as time goes by? It sounds too simple to matter, and yet the search for a rigorous answer to that question grew, over three centuries, into one of the great unifying ideas of physics: a language that explains why some light sources cast crisp interference fringes while others cast none at all, why a laser beam behaves so differently from a candle flame, how radio astronomers photograph the shadow of a black hole without ever building a telescope the size of the Earth, and why the delicate superpositions inside a quantum computer are so maddeningly hard to keep alive. The story of coherence is really two stories that grew up separately and only later discovered they were talking about the same thing: one begins at a gambling table in seventeenth century France, and the other begins in a darkened optics laboratory. It is worth following both threads from the start.

The first thread is the story of correlation, and it begins, fittingly, with a bet. In 1654 the nobleman and gambler Antoine Gombaud, better known as the Chevalier de Méré, wrote to Blaise Pascal asking how the stakes of an interrupted game of dice should fairly be divided between two players. The resulting correspondence between Pascal and Pierre de Fermat, working out the mathematics of expectation and combinatorics needed to answer such questions, is generally credited as the birth of probability theory. Their ideas matured slowly. In 1713, sixty years later, Jacob Bernoulli's posthumous book *Ars Conjectandi* Bernoulli (1713) proved the law of large numbers, the remarkable fact that if you repeat a random experiment often enough, the observed frequency of an outcome settles down, ever more reliably, around its true theoretical probability. A century later still, in 1812, Pierre Simon Laplace gathered these scattered results, together with his own theory of how new evidence should update our beliefs (what we now call Bayesian inference), into the monumental *Théorie Analytique des Probabilités* Laplace (1812). What was still missing from this whole tradition was a number, a single figure that could say not just how uncertain one quantity is, but how strongly two different quantities move together.

That missing number arrived through an unexpected route: the very practical business of dealing with noisy measurements. In 1809, working out how to fit the imperfect, scatter riddled observations of planetary orbits, Carl Friedrich Gauss introduced the method of least squares in his *Theoria Motus* Gauss (1809), a technique built entirely on the idea that errors cluster statistically around the truth. Decades later, in the 1880s, the polymath Francis Galton, fascinated by how strongly a child's height tends to track its parents', coined the very word "correlation" to describe exactly this kind of joint tendency Galton (1888). It fell to his student Karl Pearson, in 1896, to turn Galton's intuition into a precise formula: the Pearson correlation coefficient Pearson (1896), a single number, always between minus one and plus one, that tells you at a glance whether two quantities march together, drift apart, or ignore each other completely. For the first time, resemblance between two things could be measured rather than merely felt.

The second thread, meanwhile, was unfolding almost entirely without reference to the first, in the study of light itself. In 1801, Thomas Young let sunlight fall through two narrow slits and watched it paint a row of bright and dark bands on a screen behind them Young (1804), the unmistakable

signature of two waves sometimes reinforcing and sometimes cancelling one another. For those fringes to appear at all, the light leaving the two slits had to keep a steady, dependable phase relationship with itself, a property nobody yet had a name for, though everybody could see its footprint on the screen. Between 1815 and 1820, Augustin Jean Fresnel built the full mathematical theory of light as a transverse wave, and with it interference and polarisation stopped being curiosities and became consequences of a single physical picture. And yet a puzzle quietly remained. Young's beam, freshly split from a single narrow slit, produced fringes beautifully. But ordinary daylight, or the light of a candle, or of the Sun itself, never produced anything of the sort when simply allowed to interfere with a delayed copy of itself beyond a certain separation. Something about real, everyday light was different from the idealised, perfectly regular wave that Fresnel's mathematics described, and for a hundred years nobody could quite say what.

The answer, when it finally came in the twentieth century, is that ordinary light sources are not single, orderly waves at all. A hot filament, a flame, or the surface of the Sun is an enormous crowd of independent atoms, each one emitting its own short burst of light and then falling silent, entirely out of step with its neighbours, so that the overall wave forms and reforms its phase billions of times each second in a way that is, for all practical purposes, random. Describing such light properly meant marrying the two separate traditions that had grown up in isolation: the statistical machinery of Pascal, Gauss, Galton and Pearson had to be grafted onto the wave optics of Young and Fresnel. That graft is exactly what the word "coherence" now names: not whether a wave has a phase, every wave does, but whether that phase remains predictable enough, for long enough and over a wide enough region, to leave a visible trace when the wave is made to interfere with a copy of itself.

The decisive mathematical step was taken in the 1930s. In 1934 the Dutch physicist Pieter Hendrik van Cittert worked out how the statistical correlations in the light from an incoherent, extended source, a distant star, for instance, change as that light travels away from the source van Cittert (1934), and in 1938 Frits Zernike sharpened and generalised the result into the theorem that today carries both their names Zernike (1938). The van Cittert Zernike theorem makes a beautifully counterintuitive claim: light that starts out utterly incoherent, with every point on the source flickering independently of every other, becomes measurably, predictably more coherent purely by virtue of travelling a long distance. This is not a minor footnote. It is the physical principle that allows astronomers to measure the angular size of stars far too small for any telescope to resolve directly, by comparing the light collected at two separated points and watching how the fringes fade as the separation grows, exactly the trick Albert Michelson and Francis Pease used in 1920 to measure the diameter of the red giant Betelgeuse Michelson and Pease (1921). Decades later, the very same theorem, scaled up from a handful of metres to a network of radio dishes spanning the whole planet, is what let the Event Horizon Telescope collaboration reconstruct, in 2019, the first direct image of a black hole.

Classical coherence theory found its definitive, mature form in the work of Max Born and Emil Wolf, whose treatise *Principles of Optics* Born and Wolf (1999) gathered every strand into a single object, the mutual coherence function, which simply measures how correlated the light wave is between two points in space and two moments in time. Squeezed inside that one function are both of the qualities an experimenter actually cares about: the temporal coherence, essentially how pure

a colour the light is and for how long it keeps a steady rhythm with itself, and the spatial coherence, how faithfully the wave stays in step across an extended region rather than just at a single point. Leonard Mandel and Wolf later expanded this picture into the full treatise *Optical Coherence and Quantum Optics* Mandel and Wolf (1995), sharpening the practical notions of coherence time and coherence length that every laboratory now takes for granted. None of this stayed confined to the blackboard. The Michelson interferometer of the 1890s already exploited temporal coherence to measure wavelengths with a precision no ruler could match, and the invention of the laser in 1960, whose light is coherent to a degree no natural source ever achieves, turned coherence from a subject of theoretical fascination into the working principle behind fibre optic communication, precision metrology, holography and modern medicine.

Nor does coherence belong to optics alone. Wherever many independent parts of a system can be made to oscillate in step, something with exactly the same mathematical anatomy appears. In condensed matter physics, superconductivity is, at bottom, a coherence phenomenon: below a critical temperature, an enormous population of paired electrons locks into a single, shared quantum phase across an entire macroscopic sample, which is why a supercurrent flows without any resistance at all. Nuclear magnetic resonance and the magnetic resonance imaging built upon it work by tipping billions of nuclear spins into a common, coherent precession and then watching, with exquisite sensitivity, how quickly that shared rhythm decays. Every atomic clock, and with it the entire Global Positioning System, depends on an oscillator whose phase stays coherent for as long as humanly achievable. Coherence, in short, is not a peculiarity of light; it is what allows countless independent actors, electrons, spins, atoms or photons, to behave, briefly and fragilely, as one.

While all of this classical machinery was being consolidated, an entirely separate and stranger story was unfolding in the quantum world. Its opening scene took place in 1956, when Robert Hanbury Brown and Richard Twiss, trying to measure the angular sizes of stars with radio and then optical detectors, found that photons arriving from a thermal source at two separated detectors were not independent: they tended to arrive in correlated little bunches Hanbury Brown and Twiss (1956). The result seemed to many physicists at the time to be flatly impossible. Photons, everyone assumed, should behave like classical, independent particles once detected, and no such correlation should survive the act of measurement; a real and lively controversy followed before the effect was accepted. Untangling it properly required a fully quantum theory of coherence, and that theory was supplied in 1963, almost simultaneously and independently, by Roy Glauber Glauber (1963) and George Sudarshan Sudarshan (1963). Glauber introduced the coherent states, the quantum states of light that behave as closely as quantum mechanics permits to Fresnel's classical waves, and defined a whole hierarchy of quantum correlation functions capable of explaining Hanbury Brown and Twiss's bunching rigorously, and of predicting its mirror image, photon antibunching, a behaviour so stubbornly non classical that no wave theory of light could ever reproduce it, later confirmed in the laboratory. Sudarshan, working from a different direction, found the precise mathematical boundary, now called the Glauber Sudarshan representation, that separates light whose quantum nature can safely be ignored from light for which it cannot. For this quantum theory of optical coherence, Glauber received the Nobel Prize in 2005, and the field he founded is what we now call quantum optics.

Quantum coherence turned out to matter for reasons that reach far beyond photon counting. It is quantum coherence, the delicate, phase locked superposition of many possible states at once, that makes a quantum bit fundamentally different from an ordinary one, and it is the loss of that coherence through unavoidable contact with the environment, a process physicists call decoherence, that explains why the everyday world around us looks so stubbornly classical even though it is built from quantum parts. Keeping coherence alive for long enough to be useful is, to this day, the single greatest engineering obstacle standing between today's quantum computers and tomorrow's. The same idea reappears in the coldest matter physicists know how to make: in 1995 Eric Cornell, Carl Wieman and Wolfgang Ketterle cooled clouds of atoms to nanokelvin temperatures and watched millions of them collapse into a single, shared quantum state, a Bose Einstein condensate, in effect one enormous coherent matter wave, an achievement recognised with the Nobel Prize in 2001. It even reaches into biology: careful experiments on the light harvesting complexes that plants and certain bacteria use to capture sunlight have found that quantum coherence in the transfer of that energy survives, astonishingly, for measurable fractions of a second at room temperature, hinting that evolution may have stumbled upon and exploited quantum coherence long before physicists gave it a name.

Binding the statistical and the wave pictures of coherence together, classical and quantum alike, is a single piece of mathematics: the Wiener Khinchin theorem, developed independently by Norbert Wiener in 1930 Wiener (1930) and Aleksandr Khinchin in 1934 Khinchin (1934), which states that the correlation of a fluctuating signal with itself and the way that signal's energy is spread across frequency are simply two faces of the same object, translated into one another by a Fourier transform. It is, as later chapters will show, the quiet mathematical hinge on which the whole subject turns.

In the twenty first century coherence has acquired one further identity: it is now understood, and formally quantified, as a genuine physical resource, something that can be created, spent and depleted much like energy or entanglement, with its own rigorous accounting rules inside the broader theory of quantum information Baumgratz et al. (2014). Seen end to end, the history of coherence runs from a seventeenth century argument about dice, through the statistics of inherited height and the interference of sunlight, to the imaging of a black hole and the fragile inner workings of a quantum processor. In its classical guise it explains why stars twinkle the way they do, why a laser can cut steel while a torch cannot, and why an interferometer the size of a planet can photograph something no single telescope ever could. In its quantum guise it explains why the world of atoms feels so unlike the world of tables and chairs, and it now sits at the very centre of the effort to build technologies that think and communicate in ways no classical machine ever could. Coherence, whether counted in correlated dice throws, interfering light waves or entangled qubits, is ultimately always the same thing: order, briefly and remarkably, surviving in a world that constantly conspires to erase it.

6.2 How is statistical similarity quantified?

Suppose we have a completely random phenomenon, such as the electromagnetic radiation emitted by an incandescent bulb that is not in thermodynamic equilibrium, the number of cars in some

capital city throughout the day, the motion of an ion in a plasma, or something as simple as throwing a ball into the air under non-ideal conditions.

When we wish to analyse statistically a single object or random variable, the familiar tools appear: the probability density, the mean value, the standard deviation, the kurtosis, and so on. What happens when we want to study at least two random variables and how they relate? When we wish to study the statistics of two or more random variables we must study the statistical similarity between them; but statistical similarity is a rather broad concept, since we may have similarity in the mean values, or in the standard deviations, or in the probability densities. To address this problem we return to linear algebra.

If we have two vectors $\vec{A}, \vec{B} \in \mathbb{R}^n$, the dot product $\vec{A} \cdot \vec{B}$ is interpreted as the magnitude of the projection of $\vec{A}$ onto $\vec{B}$ and vice versa; it can also be read as *how much of $\vec{A}$ lies along $\vec{B}$* .

It is worth pausing on what that operation actually does mechanically, because the whole of coherence theory is built on this single gesture. Written out component by component, the dot product is

$$\vec{A} \cdot \vec{B} = \sum_{i=1}^n A_i B_i, \quad (6.1)$$

that is, *multiply the two lists entry by entry and add up the results*. Nothing more. And yet the sign of each individual term already carries exactly the information we are after: whenever A_i and B_i lean the same way, their product is positive and pushes the sum upwards; whenever they lean opposite ways, the product is negative and drags it down. If the two vectors tend to agree along most of their components, the positive terms dominate and the total comes out large; if they systematically disagree, the total comes out large but negative; and if they agree just as often as they disagree, the terms cancel one another in the sum and the total collapses to nearly zero. So “multiply the pairs and add” is already, all by itself, a similarity meter, and everything that follows in this chapter is that same gesture applied to progressively richer objects.

This behaviour can be extrapolated to continuous vector spaces, where one usually employs the notion of inner product. The step is smaller than it looks: a function is simply a vector with a continuum of components, since instead of the finite list $A_1, A_2, \dots, A_n$ we now carry the values $f_A(x)$, one for every x . The index i becomes the label x , the sum over a discrete index becomes an integral over a continuous one, and the dot product becomes the inner product. If f_A, f_B are two elements of the Hilbert space $L^2(\mathbb{R})$ of square integrable functions of one real variable, one can define an inner product as

$$\langle f_A, f_B \rangle = \int_{\mathbb{R}} f_A^*(x) f_B(x) W(x) dx, \quad (6.2)$$

where the complex conjugate on the first factor is what keeps $\langle f_A, f_A \rangle$ real and non negative even when the functions are complex, as the optical fields of this chapter will be, and where $W(x)$ is

some weight function.¹ The interpretation is still valid: this can be read as how much the function f_A over all the reals coincides with the function f_B over all the reals.

To reach the notion of correlation from linear algebra we use a bit of bracket notation, where the ket $|f_A\rangle$ represents a vector of an arbitrary Hilbert space L^2 and the bra $\langle f_A|$ represents an element of the dual of the same Hilbert space,² and the inner product of a state with itself, $\langle f_A, f_A \rangle = \langle f_A | f_A \rangle$, accumulates the probability of finding the system anywhere at all; it is the integrand $|f_A(q)|^2$, rather than the completed integral, that plays the role of the probability density of finding f_A in a state $q \pm \Delta q$, where q represents some generalised coordinate. Suppose that $f_A, f_B \in L^2$ have the form

$$|f_A\rangle = \hat{A} |\Psi_A\rangle, \quad |f_B\rangle = \hat{B} |\Psi_B\rangle, \quad (6.3)$$

where $\hat{A}, \hat{B}$ are the operators of the physical quantities of $|\Psi_A\rangle, |\Psi_B\rangle$, so that we may interpret $|f_A\rangle, |f_B\rangle$ as the wave functions steered by their expected value. Using the inner product (6.2) with unit weight function, $W = 1$, the inner product reads

$$\langle f_A | f_B \rangle = \int_{\mathbb{R}^n} \hat{A} \hat{B} \langle \Psi_A | \Psi_B \rangle d^n q = \int_{\mathbb{R}^n} \hat{A} \hat{B} P(A; B) d^n q, \quad (6.4)$$

where we see that the similarity between $|f_A\rangle$ and $|f_B\rangle$ can be viewed as the product of $\hat{A}$ with $\hat{B}$ times the probability density of having states A and B .³ That is, the similarity, or how *correlated* two variables are, can be seen as the integral of the product of the variables times their joint probability density.

With this analogy in mind we can understand correlation as a first measure of the similarity between one or more variables across space and time. If $x_1(\vec{r}, t_1), x_2(\vec{r}', t_2)$ are two random variables and $p_2(x_1, t_1; x_2, t_2)$ is the joint probability density, then the correlation between x_1, x_2 is defined as

$$\Gamma(\vec{r}, \vec{r}'; t_1, t_2) = E \{ x_1(\vec{r}, t_1) x_2^*(\vec{r}', t_2) \} = \iint x_1 x_2^* p_2(x_1, t_1; x_2, t_2) dx_1 dx_2, \quad (6.5)$$

that is, as the ensemble average of $x_1 \cdot x_2^*$, weighted by how likely each pair of values is to occur together. The conjugation is placed on the second factor here to agree with the convention used for the coherence function in the next section; for real valued variables it does nothing at all, and

¹ $W(x)$ receives this name because, as it says, it gives a weight to each value the integrand may take; there are no uniform contributions, but rather each possible value of the integrand within the domain contributes weighted by the value of W at that point.

²The Hilbert space is a particular case of the L^p spaces, the space of p -norm Lebesgue-integrable functionals; for Hilbert $p = 2$, and it has the property that the dual of L^2 is also L^2 , which lets quantum mechanics arise naturally in this space.

³This last step is deliberately heuristic: we are treating the operators as though they could simply be pulled out of the inner product and replaced by the values they measure, which is only legitimate when acting on their own eigenstates. The aim here is not a derivation but a bridge, to make plausible where the definition that follows comes from; the rigorous treatment belongs to Mandel and Wolf (1995).

for complex ones it merely conjugates Γ as a whole, leaving every magnitude we shall care about untouched.

Structurally this is the same object we have been building all along. The sum over components in (6.1) has simply become an average over the ensemble, and the role of the weight function W in (6.2) is now played by the joint probability density p_2 , which is the natural weight to use when the “components” being paired up are the possible outcomes of a random experiment rather than the entries of a list. Correlation, in other words, is the dot product of two random variables, taken in the only sense available when the quantities involved refuse to hold still.

The same sign argument that made the dot product a similarity meter now tells us how to read Γ , and it is worth spelling out the three cases explicitly. If the two variables rise and fall together, so that they are large at the same instants and small at the same instants, then the product $x_1 x_2^*$ is predominantly positive, every realisation of the experiment contributes with the same sign, and the average survives as a large positive number. If they move in strict opposition, one peaking whenever the other troughs, the product is predominantly negative and the average survives as a large negative number, which is just as informative: perfect anticorrelation is a form of perfect similarity, merely with a sign attached. But if the two variables know nothing about each other, then for every realisation in which the product happens to come out positive there is, somewhere in the ensemble, an equally likely realisation in which it comes out negative, the contributions annihilate one another, and the average collapses to zero. **The averaging is therefore not an incidental technicality but the entire mechanism:** it is precisely what destroys the terms that carry no shared information while allowing the terms that do to accumulate.

Two features of the definition (6.5) deserve a comment, because both differ from the Pearson coefficient met in the historical introduction. The first is that we have not subtracted the mean values before multiplying. Subtracting them would produce the covariance, which asks how the *fluctuations* about the average conspire; keeping them, as we do here, asks about the total quantity, and for the optical fields of this chapter the distinction happily evaporates, since a thermal field oscillates symmetrically about zero and its mean vanishes on its own. The second is that we have not divided by anything, so Γ still carries the physical units of the fields it was built from and its size depends on how intense the light happens to be, not only on how similar it is. A correlation of 10^{-6} between two feeble beams may represent far tighter agreement than a correlation of unity between two powerful ones. Repairing this, by dividing Γ by its own value at zero separation, which is precisely the intensity, so that the result is a pure number confined between zero and one that measures similarity and similarity alone, is exactly the step that turns the correlation function into the *complex degree of coherence*, and it is the point the next section works towards.

6.3 Random variables

In the previous section we spoke freely of random variables, of joint probability densities and of ensemble averages, and we did so on purpose, because the aim there was to make the definition of the correlation function plausible rather than to justify it. That debt must now be paid. Coherence

theory is, from beginning to end, a theory *about* averages, and an average that is not defined precisely defines nothing precisely. Worse, the objects we shall manipulate are not averages of numbers but averages over an infinite collection of entire field histories, and for such objects naive intuition is not merely imprecise, it is occasionally wrong.

The purpose of this section is therefore to build the notion of a random variable from the ground up, in the measure theoretic language in which it was cast by Kolmogorov (Kolmogorov, 1933), and then to apply it to the vector electromagnetic field, constructing explicitly the probability space on which the whole of the following theory lives. Every ingredient will be given its physical reading as it appears, because each one of them corresponds to something an optician does in a laboratory.

6.3.1 The sample space and the physical meaning of an event

Definition 26 (*Sample space*) A **sample space** is a nonempty set Ω . Its elements $\omega \in \Omega$ are called **elementary outcomes** or **sample points**.

The temptation is to read ω as a number, and it is essential to resist it. In optics an elementary outcome is not a field value and not a measurement; it is one complete **realisation** of the source, that is, one entire spacetime history that the field could have had. A thermal lamp contains of the order of 10^{20} atoms radiating independently, each with an emission instant and an initial phase that we neither control nor know. One particular assignment of all those microscopic details, together with everything that follows from it by Maxwell's equations, is one ω . The set of all such assignments is Ω .

This is precisely the picture that Figure 6.1 will show in the next section, where several curves $^{(i)}E(\vec{r}, t)$ are drawn one above the other: each curve is one ω , and the label i is only a bookkeeping device, since in general Ω is uncountable. What the laboratory delivers on any given afternoon is a single ω , chosen by nature and not revealed to us. The whole of statistical optics consists in extracting statements about Ω from access to one of its points.

Definition 27 (*Event*) An **event** is a subset $A \subseteq \Omega$.

The physical content of this definition is that an event is a **statement about the realisation that can be declared true or false**. The statement “the field at $\vec{r}_0$ at the instant t_0 exceeded the value E_0 ” is not a number; it is the set of all realisations for which that happens to be the case. To say that the event occurred is to say that the ω nature selected belongs to that subset. Events and statements are the same thing seen from two sides, and this identification is what allows probability to be built out of set theory.

6.3.2 The sigma algebra, or which questions may be asked

Definition 28 (*Sigma algebra*) A **sigma algebra** on Ω is a collection $\mathcal{F}$ of subsets of Ω such that

1. $\Omega \in \mathcal{F}$,
2. if $A \in \mathcal{F}$ then $A^c = \Omega \setminus A \in \mathcal{F}$,
3. if $A_1, A_2, A_3, \dots \in \mathcal{F}$ then $\bigcup_{n=1}^{\infty} A_n \in \mathcal{F}$.

The pair $(\Omega, \mathcal{F})$ is called a **measurable space**.

A sigma algebra looks like a piece of bookkeeping and is in fact a physical statement. It is the **catalogue of admissible questions**, the list of statements to which the theory is prepared to attach a probability, and the three axioms are exactly the three properties that any honest catalogue of questions must have.

The first says that the question “did anything at all happen?” is admissible. The second says that if we can ask whether an event occurred, we can ask whether it did not, which is the formal expression of the fact that an apparatus capable of detecting something is by that very fact capable of detecting its absence. The third says that if each of a sequence of questions is admissible, so is the question of whether at least one of them holds.

The word *countable* in the third axiom is not a technical convenience but a genuine physical restriction, and it deserves to be pointed out. Any real experimental protocol is a *sequence* of operations: one records a reading, then another, then another. No laboratory has ever performed an uncountable number of operations, and the theory is built so as to promise nothing about what would happen if one could.

One might ask why we do not simply take $\mathcal{F}$ to be the collection of *all* subsets of Ω and be done with it. The answer is that we are not allowed to. Vitali showed in 1905 (Vitali, 1905) that on the real line there exist sets so pathological that no translation invariant notion of size can be assigned to them without contradiction. If we insisted on measuring everything we would be forced into inconsistency, and the sigma algebra is precisely the device that keeps such monsters outside the theory while retaining every set that a physicist would ever have occasion to ask about.

The example we shall need constantly is the following.

Definition 29 (*Borel sigma algebra*) The **Borel sigma algebra** $\mathcal{B}(\mathbb{R}^n)$ is the smallest sigma algebra on $\mathbb{R}^n$ containing all the open sets.

Physically, $\mathcal{B}(\mathbb{R})$ is generated by the questions “does the reading lie between a and b ?”, together with everything that can be built from countably many such questions by negation and union. This is exactly the class of questions that a real detector, which has a finite resolution and reports an interval rather than a point, is able to answer. The Borel sigma algebra is not an abstraction imposed on physics; it is the mathematical transcription of what a graduated instrument does.

6.3.3 Measure, the Lebesgue measure and the probability measure

Definition 30 (*Measure*) A **measure** on a measurable space $(\Omega, \mathcal{F})$ is a function $\mu : \mathcal{F} \rightarrow [0, \infty]$ such that $\mu(\emptyset) = 0$ and such that, for every sequence $A_1, A_2, \dots$ of pairwise disjoint elements of $\mathcal{F}$,

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n). \quad (6.6)$$

The triple $(\Omega, \mathcal{F}, \mu)$ is a **measure space**.

A measure is a consistent way of assigning a *size*. Condition (6.6) is countable additivity, and it states nothing more exotic than that the size of a whole assembled out of separate pieces is the sum of the sizes of the pieces. That is what the word size means operationally, and a function which violated it would not deserve the name.

Among all measures one is singled out by the geometry of space itself.

Definition 31 (*Lebesgue measure*) The **Lebesgue measure** λ is the unique translation invariant measure on $\mathcal{B}(\mathbb{R}^n)$ satisfying $\lambda\left(\prod_{j=1}^n [a_j, b_j]\right) = \prod_{j=1}^n (b_j - a_j)$.

The Lebesgue measure (Lebesgue, 1902) is the mathematical form of the words *length*, *area*, *volume* and *duration*. Every integral we have written so far in these notes has been an integral against λ : when we integrated over an aperture in the theory of diffraction we were measuring *how much aperture*, and when we integrate over the response time of a detector we are measuring *how long we looked*. The Lebesgue measure knows about the extent of the domain and it knows nothing whatever about probability.

It is worth insisting on this separation, because in coherence theory both measures appear at once and they do entirely different jobs. **The Lebesgue measure records how long the detector integrated and how large the aperture was; the probability measure records how likely each realisation of the source was.** The time average of a single beam is an integral against λ on the time axis; the ensemble average over realisations is an integral against P on Ω . These are different operations on different spaces, and the fact that they so often agree is not a triviality but a theorem with hypotheses, to which we return at the end of this section.

The Lebesgue measure also supplies the missing justification for the densities of the previous section. When in (6.5) we wrote $p_2(x_1, t_1; x_2, t_2) dx_1 dx_2$ we were tacitly assuming that the probability measure on the space of values is absolutely continuous with respect to λ , so that by the theorem of Radon and Nikodym (Nikodym, 1930) it possesses a density. That is a physical hypothesis and not a formality: it asserts that no field value occurs with strictly positive probability, that is, that the source has no infinitely sharp features. A perfectly monochromatic and perfectly stabilised laser would violate it, since its distribution would be concentrated on a circle in the complex plane, a set of zero Lebesgue measure, and no density with respect to the plane would exist.

Definition 32 (*Probability space*) A **probability measure** is a measure P with $P(\Omega) = 1$. The triple $(\Omega, \mathcal{F}, P)$ is called a **probability space**.

The normalisation $P(\Omega) = 1$ is the statement that the source does *something*: some realisation occurs, and the alternatives listed in Ω are exhaustive. Countable additivity, read now in P , is the statement that mutually exclusive alternatives have probabilities that add.

Here lies the crucial physical remark of this subsection. **The probability measure is where the physics of the source resides.** A thermal lamp, a laser and a single photon emitter may perfectly well share the same Ω and the same $\mathcal{F}$, since all three could in principle produce any of the same field histories; what distinguishes them is that they weight those histories differently. To specify a source is to specify P , and every statement that coherence theory will make about light is, in the last analysis, a statement about P .

6.3.4 Random variables and the meaning of measurability

Definition 33 (*Random variable*) Let $(\Omega, \mathcal{F}, P)$ be a probability space and $(S, \mathcal{S})$ a measurable space. A map $X : \Omega \rightarrow S$ is a **random variable** if it is **measurable**, that is, if

$$X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\} \in \mathcal{F} \quad \text{for every } B \in \mathcal{S}. \quad (6.7)$$

Two things must be said about this definition, and both are usually said too quickly.

The first is that **a random variable is not a variable that is random**. It is a perfectly deterministic function. Once the realisation ω has been fixed, $X(\omega)$ is a definite element of S , with nothing capricious about it. All the randomness lives in the selection of ω , which is to say in our ignorance of which realisation nature has produced. The field at a point of spacetime is not fluctuating because it is undecided; it is definite in each realisation and we simply do not know which realisation we are in.

The second is that the measurability condition (6.7) is the mathematical statement that **the instrument is a legitimate instrument**. It demands that the question “did the reading fall inside B ?” be one of the admissible questions of $\mathcal{F}$. A non measurable map would be an apparatus that reports an answer to a question the theory refuses to assign a probability to, which is a polite way of saying that it reports numbers that mean nothing. Measurability is the licence to exist.

Since the outcome ω is inaccessible, what an experiment actually reconstructs is not P but its image.

Definition 34 (*Law of a random variable*) The **law or distribution** of X is the probability measure P_X on $(S, \mathcal{S})$ defined by the push forward

$$P_X(B) = P(X^{-1}(B)), \quad B \in \mathcal{S}. \quad (6.8)$$

If P_X is absolutely continuous with respect to the Lebesgue measure, its Radon-Nikodym derivative $p = dP_X/d\lambda$ is the **probability density** of X .

The law is what a histogram estimates. Two sources built on completely different sample spaces may possess identical laws for a given observable, and are then indistinguishable by that observable no matter how long one measures. The density p appearing here is exactly the p_2 that was used without apology in (6.5).

Definition 35 (*Expectation*) The **expectation** of an integrable random variable X is the Lebesgue integral of X over the probability space,

$$E\{X\} = \int_{\Omega} X(\omega) dP(\omega) = \int_S x dP_X(x) = \int_S x p(x) d\lambda(x), \quad (6.9)$$

the second equality being the change of variables induced by (6.8).

Equation (6.9) retroactively legitimises the definition (6.5) of the correlation function. What was written there as an ensemble average and what was written as an integral against a joint density are not two definitions that happen to agree; they are one definition, given on Ω , and its image on the space of values under the push forward. The first form is conceptually primary and the second is the one we can compute with.

There is now a payoff which is worth stating explicitly, because it converts the central analogy of the previous section from a metaphor into a theorem. Consider the set of random variables of finite second moment,

$$L^2(\Omega, \mathcal{F}, P) = \{X : \Omega \rightarrow \mathbb{C} \mid E\{|X|^2\} < \infty\}, \quad (6.10)$$

equipped with the inner product

$$\langle X, Y \rangle = E\{XY^*\}. \quad (6.11)$$

This is a Hilbert space, and comparing (6.11) with the definition (6.5) of the correlation function we see that **the correlation function is an inner product**, not by analogy but literally, in the space of random variables of finite energy. The dot product with which the previous section began was not an illustrative parallel; it was the same object, waiting for the space to be named. The weight function W of (6.2) has become the measure P , and the sum over components of (6.1) has become the integral over Ω .

From this identification one inequality follows immediately, and it is the one that makes the normalised degree of coherence meaningful. The Cauchy-Schwarz inequality in $L^2(\Omega, \mathcal{F}, P)$ reads

$$|E\{XY^*\}|^2 \leq E\{|X|^2\} E\{|Y|^2\}, \quad (6.12)$$

so that the correlation of two random variables can never exceed the geometric mean of their intensities. When we divide by that geometric mean, as anticipated at the end of the previous section, the result is a pure number lying between zero and one. It is, quite literally, the cosine of the angle between two vectors in a Hilbert space, and this is why a degree of coherence larger than unity is not merely unobserved but impossible.

6.3.5 Stochastic processes

A single random variable describes the field at one point of spacetime. To describe it everywhere at once we need a family.

Definition 36 (*Stochastic process*) Let T be an index set. A **stochastic process** is a family $\{X_\tau\}_{\tau \in T}$ of random variables defined on a common probability space $(\Omega, \mathcal{F}, P)$, equivalently a map $X : T \times \Omega \rightarrow S$ such that $X(\tau, \cdot)$ is measurable for every $\tau \in T$.

A stochastic process admits two readings, and the whole intuition of the subject lies in holding both at once (Doob, 1953). If we fix the index τ and let ω vary, we obtain a random variable: a *vertical* slice through Figure 6.1, comparing all realisations at one instant. If instead we fix ω and let τ vary, we obtain a single function of the index, called a **sample path** or **realisation**: a *horizontal* slice, that is, one of the individual curves of the figure. The experimentalist observes horizontal slices and the theory speaks about vertical ones, and reconciling the two is the central practical problem of statistical optics.

6.3.6 The probability space of the electromagnetic field

We can now carry out the construction that this whole section was preparing, and build explicitly the probability space on which coherence theory operates. Let $D \subseteq \mathbb{R}^3 \times \mathbb{R}$ be the region of spacetime in which we intend to describe the field, and recall from the chapter on electromagnetic optics that the field is a complex, three component object, the analytic signal $\vec{\mathcal{E}} = (\mathcal{E}_x, \mathcal{E}_y, \mathcal{E}_z)$.

The sample space. We take

$$\Omega = \{\omega : D \rightarrow \mathbb{C}^3 \mid \omega \text{ satisfies Maxwell's equations in } D\}, \quad (6.13)$$

that is, the set of admissible field histories. The restriction is essential and it is physical: nature does not select arbitrary functions of spacetime but only those compatible with the dynamics. **Randomness enters through the sources and never through the propagation**, which remains exactly as deterministic as it was in the previous chapters. A probability measure on an optical field is a measure on the solution space of Maxwell's equations, and this is why, as we shall see in a later subsection, coherence itself obeys a wave equation.

With this choice the field becomes the simplest map imaginable, namely the evaluation of a function at a point,

$$\mathcal{E}_j(\vec{r}, t) : \Omega \rightarrow \mathbb{C}, \quad \omega \mapsto \omega_j(\vec{r}, t), \quad j \in \{x, y, z\}, \quad (6.14)$$

so that for each spacetime point and each Cartesian component the field is a complex random variable, and the vector $\vec{\mathcal{E}}(\vec{r}, t) : \Omega \rightarrow \mathbb{C}^3$ is a $\mathbb{C}^3$ valued random variable. The whole field is the stochastic process indexed by $T = D \times \{x, y, z\}$.

The sigma algebra. We equip Ω with the **cylinder sigma algebra** $\mathcal{F}$, generated by the sets

$$C = \{\omega \in \Omega \mid (\omega(\vec{r}_1, t_1), \dots, \omega(\vec{r}_n, t_n)) \in B\}, \quad B \in \mathcal{B}(\mathbb{C}^{3n}), \quad (6.15)$$

for every finite collection of spacetime points and every Borel set B .

The physical reading of (6.15) is the most important remark of this section. **A cylinder set is exactly a statement that can be checked by placing a finite number of detectors at a finite number of spacetime points and looking at their readings.** No experiment has ever measured a field everywhere; every experiment samples it at finitely many places and finitely many instants. The sigma algebra generated by the cylinder sets is therefore the collection of all statements decidable by countable sequences of finite measurements, which is to say, the collection of all statements an optical experiment can ever settle. The cylinder construction is not a mathematical convenience, it is the transcription of experimental practice.

The probability measure. It remains to place a measure on this enormous space, and it would be hopeless to attempt it directly. Kolmogorov's extension theorem (Kolmogorov, 1933) shows that it is unnecessary.

Theorem 9 (*Kolmogorov extension theorem*) *Given a family of finite dimensional distributions $P_{(\vec{r}_1, t_1), \dots, (\vec{r}_n, t_n)}$ on $\mathbb{C}^{3n}$, one for every finite collection of spacetime points, which is consistent under permutation of the points and under marginalisation over any of them, there exists a unique probability measure P on the cylinder sigma algebra $\mathcal{F}$ whose finite dimensional marginals are the given ones.*

The physical content is emancipating. **To specify a source it is enough to specify the joint statistics of any finite set of detector readings**, and this is precisely the data that an experiment can access. We never have to say what P does to the monstrous space of all field histories; we say what the joint distribution of two detectors is, and of three, and so on, taking care only that these prescriptions do not contradict one another, and Kolmogorov guarantees that they knit together into one consistent object. The probability space of an optical field is built out of exactly the information a laboratory can supply.

The finite energy hypothesis. We assume finally that

$$E \left\{ |\mathcal{E}_j(\vec{r}, t)|^2 \right\} < \infty \quad \text{for every } j \text{ and every } (\vec{r}, t) \in D, \quad (6.16)$$

which is the statement that the mean energy density is finite, that is, that the source does not radiate infinite power. Physically it is unobjectionable, and mathematically it is exactly the hypothesis that places every field component inside the Hilbert space (6.10). From this point on, every component of the field at every point of spacetime is a vector in one and the same Hilbert space, and the geometry of that space is the geometry of coherence.

The coherence tensor. We may now define, with every symbol having been given a meaning, the object that the rest of this chapter studies. For two spacetime points and two Cartesian components,

$$\Gamma_{jk}(\vec{r}, t; \vec{r}', t') = E \{ \mathcal{E}_j(\vec{r}, t) \mathcal{E}_k^*(\vec{r}', t') \} = \langle \mathcal{E}_j(\vec{r}, t), \mathcal{E}_k(\vec{r}', t') \rangle, \quad (6.17)$$

a 3×3 matrix attached to every pair of spacetime points, the inner product being the one of (6.11). This is the vector generalisation of the correlation function (6.5), and three of its properties follow at once from the construction.

1. **Hermitian symmetry.** Exchanging the two points and the two indices and conjugating,

$$\Gamma_{kj}^*(\vec{r}', t'; \vec{r}, t) = E \{ \mathcal{E}_k^*(\vec{r}', t') \mathcal{E}_j(\vec{r}, t) \} = \Gamma_{jk}(\vec{r}, t; \vec{r}', t'), \quad (6.18)$$

since conjugation may be taken inside the expectation, which is an integral of a complex function.

2. **Non negative definiteness.** For any finite set of spacetime points labelled by m and any complex coefficients $a_j^{(m)}$,

$$\sum_{m,n} \sum_{j,k} a_j^{(m)} a_k^{(n)*} \Gamma_{jk}(m; n) = E \left\{ \left| \sum_m \sum_j a_j^{(m)} \mathcal{E}_j(m) \right|^2 \right\} \geq 0, \quad (6.19)$$

because the expectation of a non negative quantity is non negative. Physically this says that no superposition of field components, at whatever points, can carry negative intensity.

3. **Bounded correlation.** By (6.12),

$$|\Gamma_{jk}(\vec{r}, t; \vec{r}', t')|^2 \leq \Gamma_{jj}(\vec{r}, t; \vec{r}, t) \Gamma_{kk}(\vec{r}', t'; \vec{r}', t'), \quad (6.20)$$

which is what allows the normalised degree of coherence to be confined to the interval $[0, 1]$.

It is worth noticing where polarisation fits into this. Evaluating (6.17) at a single spacetime point, $\vec{r} = \vec{r}'$ and $t = t'$, the diagonal entries Γ_{jj} give the mean intensity carried by each Cartesian component while the off diagonal entries measure the correlation between components, and the resulting matrix is exactly the polarisation matrix (3.67) introduced in the chapter on polarisation. **Polarisation is coherence between the components of the field at one point; coherence proper is correlation between the field at different points.** They are the same tensor read in two different regimes, which is the reason a unified treatment of coherence and polarisation is possible at all (Wolf, 2007).

6.3.7 Stationarity, ergodicity and the right to measure

One question remains, and it is the question on which the experimental relevance of everything above depends. The expectation (6.9) is an average over Ω , and Ω is not accessible: nature hands us one realisation. What we can average over is time, along that single realisation. Under what conditions do the two agree?

The first condition concerns the invariance of the source.

Definition 37 (*Stationarity*) Let $\theta_\tau : \Omega \rightarrow \Omega$ be the time shift defined by $(\theta_\tau \omega)(\vec{r}, t) = \omega(\vec{r}, t + \tau)$. The field is **stationary** if P is invariant under every such shift,

$$P(\theta_\tau^{-1}A) = P(A) \quad \text{for all } A \in \mathcal{F} \text{ and all } \tau \in \mathbb{R}. \quad (6.21)$$

This is the measure theoretic form of the statement, given in a more elementary way in the next section, that the statistical properties of the light do not depend on when we choose to start the clock. A lamp switched on an hour ago and left alone is stationary; a pulsed source is not. Its immediate consequence is that the coherence tensor (6.17) can depend on the two instants only through their difference, $\Gamma_{jk}(\vec{r}, \vec{r}'; t, t') = \Gamma_{jk}(\vec{r}, \vec{r}'; \tau)$ with $\tau = t' - t$, which is what makes the Wiener-Khinchin theorem of the following section possible at all.

The second condition is stronger and concerns the indecomposability of the source.

Definition 38 (*Ergodicity*) A stationary field is **ergodic** if every shift invariant event is trivial, that is, if $\theta_\tau^{-1}A = A$ for all τ implies $P(A) = 0$ or $P(A) = 1$.

Physically, ergodicity says that the source cannot be split into two sub ensembles that never communicate. If it could, a single realisation would remain trapped in one of them forever and could never report on the other. When it holds, Birkhoff's ergodic theorem (Birkhoff, 1931) applies and gives the result we need,

$$\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T \mathcal{E}_j(\vec{r}, t) \mathcal{E}_k^*(\vec{r}', t + \tau) dt = E \{ \mathcal{E}_j(\vec{r}, t) \mathcal{E}_k^*(\vec{r}', t + \tau) \} \quad (6.22)$$

for almost every realisation ω .

Equation (6.22) is the licence, and the only licence, to replace the inaccessible ensemble average by the accessible time average of a single beam, and it is the reason the whole theory is experimentally meaningful rather than merely elegant. Notice, finally, what the two sides of (6.22) are: on the left an integral against the Lebesgue measure on the time axis, on the right an integral against the probability measure on Ω . The two measures we were at pains to keep apart at the beginning of this section meet here, in a single identity, and ergodicity is precisely the hypothesis under which they may be exchanged. With that hypothesis in hand we may return to physics and define the coherence function itself.

6.4 The coherence function

The concept of the coherence function is, at the mathematical level, identical to that of the correlation function given in equation (6.5); both represent correlations. However, coherence is a term widely used in the study of electromagnetic radiation; let us see a little of why this is so.

In Figure 6.1 we observe a component of the electric field, represented by $E(\vec{r}, t)$, seen across several observations or realisations. Behaving as a random variable, in each realisation it has a




Figure 6.1: Representation of $E(\vec{r}, t)$ as a random variable.

different behaviour in time at a point $\vec{r}$; in general, here we have indexed $^{(i)}E(\vec{r}, t)$ with integers, but in general there could be an uncountable set of possible states for the electric field. If we look at Figure 6.1 there are two temporal points, t, t' , at which we isolate parts of these random oscillations; we could ask how correlated these oscillations are across the realisations over the interval $t' - t$, or in other words, how *coherent* the oscillations are across the realisations.

We speak of coherence when we want to analyse the correlation of electromagnetic radiation between spacetime points. If we have two sources emitting an electric field (for the moment understood as scalar), their coherence function is given by

$$\Gamma(\vec{r}, \vec{r}'; t, t') = E \{ \mathcal{E}(\vec{r}, t) \mathcal{E}^*(\vec{r}', t') \}, \quad (6.23)$$

where we are computing the ensemble average of the components $\mathcal{E}$, which are complex numbers as already discussed in these notes. If we take $\vec{r} = \vec{r}'$ we obtain what is known as the *mutual coherence function*

$$\Gamma(\vec{r}; t, t') = E \{ \mathcal{E}(\vec{r}, t) \mathcal{E}^*(\vec{r}, t') \}. \quad (6.24)$$

We can go even further by including the concept of *statistical stationarity*, but one must be careful since there are several kinds. Formally, a stochastic process $\{X_t\}$ is considered stationary when its statistical properties are invariant under time translations. This condition manifests in different degrees:

1. **Strict (strong) stationarity:** the most demanding form of stationarity requires that all joint probability distributions be time-invariant:

$$F_X(x_{t_1+\tau}, \dots, x_{t_k+\tau}) = F_X(x_{t_1}, \dots, x_{t_k}) \quad \forall \tau, k, t_i, \quad (6.25)$$

where F_X is the joint distribution function. This implies that:

- all moments (mean, variance, skewness, etc.) are constant;
- the marginal distributions are identical at any instant;
- it is a sufficient but not necessary condition for ARMA models.

2. **Weak (second-order) stationarity:** more practical in applications, it only requires invariance of the first two moments:

$$\begin{cases} \mathbb{E}[X_t] = \mu & (\text{constant}) \quad \forall t \\ \text{Var}(X_t) = \sigma^2 & (\text{constant}) \quad \forall t \\ \text{Cov}(X_t, X_{t+h}) = \gamma(h) & (\text{depends only on } h, \text{ not on } t) \end{cases} \quad (6.26)$$

This is the definition used in most econometric models (ARIMA, GARCH) and in the WienerKhinchin spectral analysis (which we shall use later).

3. **Trend stationarity (deterministic stationarity)**: a non-stationary process that can be made stationary by removing deterministic components:

$$X_t = \alpha + \beta t + \epsilon_t, \quad (6.27)$$

where ϵ_t is stationary. The non-stationarity here comes exclusively from the temporal trend βt .

We shall use a hypothesis of strong stationarity (although weak stationarity often suffices). In this spirit, if $\tau = t - t'$ we have

$$\Gamma(\vec{r}', t, t') = \Gamma(\vec{r}', t - t') = \Gamma(\vec{r}', \tau). \quad (6.28)$$

As we can see, the mutual coherence at a point $\vec{r}'$ under a stationarity hypothesis gives rise to an explicit dependence on the time difference τ in the ensemble average.

It is worth asking: how necessary is it to analyse the ensemble average; why not take the temporal average? Answering this is **very complicated**, since it touches on the problem of *ergodicity*. Ergodicity, in the sense of ergodic theory, requires that a dynamical system satisfy that the temporal averages coincide with the spatial averages for almost every initial condition. Its rigorous proof is highly non-trivial, to the point that the problems for which ergodicity has been proven with mathematical rigour are few.

From a phenomenological standpoint we shall assume, from now on, an *ergodic hypothesis* in which electromagnetic radiation has a stationary and ergodic character. Under these conditions the mutual coherence function coincides with the temporal average

$$\Gamma(\vec{r}'; t, t') = \int_{\mathbb{R}} \mathcal{E}(\vec{r}', t') \mathcal{E}^*(\vec{r}', t - t') dt'. \quad (6.29)$$

The mutual coherence function has several properties, some of which are:

- a) $\Gamma(\vec{r}', 0) \geq 0$. This follows from the definition, since $\Gamma(\vec{r}', 0) = \langle |\mathcal{E}(\vec{r}')|^2 \rangle$, i.e. it corresponds to the intensity of the electric field and can only be zero when trivially the field is zero at every instant.
- b) $\Gamma(\vec{r}', -\tau) = \Gamma^*(\vec{r}', \tau)$. This is known as the *Hermitian property* of the mutual coherence function, and follows from the fact that $\mathcal{E}(\vec{r}', t)$ is stationary, which allows arbitrary temporal translations from the origin. In this spirit,

$$\Gamma(\vec{r}', \tau) = \langle \mathcal{E}(\vec{r}') \mathcal{E}^*(\vec{r}', t + \tau) \rangle = \langle \mathcal{E}(\vec{r}', t - \tau) \mathcal{E}^*(\vec{r}') \rangle = \Gamma^*(\vec{r}', -\tau). \quad (6.30)$$

c) $|\Gamma(\vec{r}', \tau)| \leq \Gamma(\vec{r}', 0)$. This property follows from the *CauchySchwarz inequality*

$$|\Gamma(\vec{r}', \tau)|^2 = |\langle \mathcal{E}(\vec{r}') \mathcal{E}^*(\vec{r}', t + \tau) \rangle|^2 \leq \langle \mathcal{E}(\vec{r}', t) \mathcal{E}^*(\vec{r}', t) \rangle \langle \mathcal{E}(\vec{r}', t + \tau) \mathcal{E}^*(\vec{r}', t + \tau) \rangle = \Gamma^2(\vec{r}', 0), \quad (6.31)$$

which implies that $|\Gamma(\vec{r}', \tau)|$ can never exceed its initial value $\Gamma(\vec{r}', 0)$. (This property is, let us recall, exclusive to the mutual coherence function; when we consider the coherence between two sources we will see that it is not satisfied - on the contrary, it will increase.)

d) $\Gamma(\vec{r}', \tau)$ is non-negative. We can see this from the definition through the following, rather obvious, inequality

$$\left\langle \left| \int_{T_1}^{T_2} f(t) \mathcal{E}(t) dt \right|^2 \right\rangle \geq 0, \quad (6.32)$$

where $f(t)$ is an arbitrary function and T_1, T_2 are arbitrary times. In this way we obtain the inequality

$$\int_{T_1}^{T_2} \int_{T_1}^{T_2} f^*(\vec{r}', t) f(\vec{r}', t') \Gamma(\vec{r}', t' - t) dt dt' \geq 0. \quad (6.33)$$

e) If the electric field is random, stationary and differentiable, then we can find a velocity $\xi(\vec{r}', t) = \dot{\mathcal{E}}(\vec{r}', t)$, which is stationary, random and of zero mean value, and whose mutual coherence function is

$$\Gamma_\xi(\vec{r}', \tau) = -\ddot{\Gamma}_\mathcal{E}(\vec{r}', \tau). \quad (6.34)$$

f) $\Gamma(\vec{r}', \tau)$ is continuous. This holds even when there are jump discontinuities.

Finally, it is worth mentioning that there is a quantity known as the complex degree of coherence, given by

$$\gamma(\vec{r}', \tau) = \frac{\Gamma(\vec{r}', \tau)}{\Gamma(\vec{r}', 0)}, \quad (6.35)$$

and from the CauchySchwarz inequality we have $0 \leq |\gamma(\vec{r}', \tau)| \leq 1$.

6.4.1 Manifestations of coherence: the WienerKhinchin theorem

The mutual coherence function and the power spectral density are two sides of the same coin in the study of light and wave signals. Mutual coherence describes how the fluctuations of a luminous field are correlated in space and time, acting as a map that reveals to what extent the variations at

one point of space are synchronised with those at another, even with a certain temporal delay. The power spectral density, in turn, decomposes the energy of the signal into its frequency components, showing which tones or colours - in the case of light - dominate its behaviour. The intimate relation between the two concepts lies in the fact that, through mathematical transformations, it is possible to convert the detailed information on the spacetime correlations captured by mutual coherence into a precise description of how the energy is distributed in the frequency domain, and vice versa. This link allows researchers to jump between complementary perspectives: analysing interference patterns in an interferometer or identifying spectral signatures in a material, all from the same fundamental principles.

We are going to find one of the most important theorems in coherence theory, the *WienerKhinchin theorem*. Note that stationary random signals do not have a Fourier transform; hence we may define the following truncated random signals of the electric field

$${}^\omega \mathcal{E}_T(\vec{r}, t) = \begin{cases} {}^\omega \mathcal{E}(\vec{r}, t) & -\frac{T}{2} \leq t \leq \frac{T}{2} \\ 0 & \text{otherwise} \end{cases} \quad (6.36)$$

and we define the spectral power for a realisation ω as the squared magnitude of the Fourier transforms of the electric fields (6.36), that is,

$${}^\omega S_T(\vec{r}, \nu) = \left| \int_{\mathbb{R}} {}^\omega \mathcal{E}_T(\vec{r}, t) e^{i2\pi\nu t} dt \right|^2 = \left| \int_{-T/2}^{T/2} {}^\omega \mathcal{E}(\vec{r}, t) e^{i2\pi\nu t} dt \right|^2, \quad (6.37)$$

and the *power spectral density of the event* ω we compute as

$$\lim_{T \rightarrow \infty} \frac{{}^\omega S_T(\vec{r}, \nu)}{T} = {}^\omega S(\vec{r}, \nu). \quad (6.38)$$

Since we already possess a tool to study, in each event or realisation, the power spectral density, then, through its ensemble average, we can find the total density, that is,

$$S(\vec{r}, \nu) = E \{ {}^\omega S(\vec{r}, \nu) \}, \quad (6.39)$$

introducing equations (6.37) and (6.38) we then have

$$S(\vec{r}, \nu) = E \left\{ \lim_{T \rightarrow \infty} \left| \frac{1}{T} \int_{-T/2}^{T/2} {}^\omega \mathcal{E}(\vec{r}, t) e^{i2\pi\nu t} dt \right|^2 \right\}, \quad (6.40)$$

$$S(\vec{r}, \nu) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_{-T/2}^{T/2} \int_{-T/2}^{T/2} E \{ \mathcal{E}(\vec{r}, t) \mathcal{E}^*(\vec{r}', t') \} e^{i2\pi\nu(t-t')} dt dt', \quad (6.41)$$

$$S(\vec{r}, \nu) = \int_{\mathbb{R}} \Gamma(\vec{r}, \vec{r}'; \tau) e^{i2\pi\nu\tau} d\tau, \quad (6.42)$$

$$S(\vec{r}, \nu) = \mathcal{F} \{ \Gamma(\vec{r}, \vec{r}', \tau) \}. \quad (6.43)$$

Result (6.43) reads: the Fourier transform of the coherence function corresponds to the power spectral density, and is known as the WienerKhinchin theorem. Its use is fundamental, since it tells us that by methods such as spectrometry we can find the coherence function of any phenomenon; we can even go further, since this result really stems from something more general - the correlation function - meaning that every stochastic phenomenon has an associated spectrum, and from the measurements or estimates of that spectrum we can compute the correlations in the system.

If we make a simple observation, the optical range of radiation - the visible - is a band of very low frequencies relative to the whole spectrum; the visible runs roughly from 400 nm to 700 nm, and we may say that all of optics is *narrowband*. This obeys the inequality

$$\frac{\Delta\nu}{\nu} \ll 1, \quad (6.44)$$

and, in this narrowband regime, the coherence function takes the form

$$\Gamma(\vec{r}; \tau) = \tilde{\Gamma}(\vec{r}) e^{i2\pi\tilde{\nu}\tau}, \quad (6.45)$$

which we can represent graphically as shown in Figure 6.2, where one can see the behaviour of the coherence function in the narrowband regime.⁴ There one sees an envelope (the observable) enveloping all the fundamental frequencies.

6.4.2 The wave equation of coherence

Let us step a little out of the comfort zone and understand how correlations propagate through space and time. Let $\mathcal{E}^{(r)}(\vec{r}, t)$ be some **real** component of the electric field, viewed as a random signal. As is well known from classical electromagnetic theory, it obeys Maxwell's equations and therefore has an associated wave equation, namely

$$\nabla^2 \mathcal{E}^{(r)}(\vec{r}, t) = \frac{1}{C^2} \frac{\partial^2 \mathcal{E}^{(r)}(\vec{r}, t)}{\partial t^2}, \quad (6.46)$$

where C is the speed of light in vacuum. We also have, by Fourier analysis,

$$\mathcal{E}(\vec{r}, t) = \int_0^\infty \tilde{\mathcal{E}}(\vec{r}, \nu) e^{-i2\pi\nu t} d\nu. \quad (6.47)$$

⁴One usually finds in the literature a term similar but not equal to that of narrowband, known as the quasi-monochromaticity condition, written as $\frac{\Delta\lambda}{\lambda} \ll 1$; but this bounds the wavelength so that it is as monochromatic as possible, while the narrowband condition admits several frequencies, provided they are narrow relative to the whole frequency spectrum.




Figure 6.2: Graphical representation of the narrowband coherence function. In black, the envelope $\tilde{\Gamma}(\vec{r}, \tau)$ is shown, and in red the oscillations $e^{i2\pi\tilde{\nu}\tau}$. These oscillations are confined within something known as the coherence time τ_C , which we shall see later.

And if we apply the d'Alembertian operator $\square = \nabla^2 - \frac{1}{C^2} \frac{\partial^2}{\partial t^2}$ to equation (6.47), we find that $\tilde{\mathcal{E}}(\vec{r}, \nu)$ has the form of a Helmholtz-type equation, that is,

$$\nabla^2 \mathcal{E}(\vec{r}, t) - \frac{1}{C^2} \frac{\partial^2 \mathcal{E}(\vec{r}, t)}{\partial t^2} = \int_0^\infty \left[\nabla^2 \tilde{\mathcal{E}}(\vec{r}, \nu) + k^2 \tilde{\mathcal{E}}(\vec{r}, \nu) \right] e^{-i2\pi\nu t} d\nu, \quad (6.48)$$

which leads us to the fact that the electric field, as an analytic and complex signal, also obeys a wave-type equation

$$\nabla^2 \mathcal{E}(\vec{r}, t) = \frac{1}{C^2} \frac{\partial^2 \mathcal{E}(\vec{r}, t)}{\partial t^2}. \quad (6.49)$$

Taking the complex conjugate of the previous equation we have

$$\nabla^2 \mathcal{E}^*(\vec{r}, t) = \frac{1}{C^2} \frac{\partial^2 \mathcal{E}^*(\vec{r}, t)}{\partial t^2}. \quad (6.50)$$

If we multiply both sides of equation (6.50) by $\mathcal{E}(\vec{r}', t')$, and since we have taken the d'Alembertian with respect to the unprimed variables, using some sign rules we have

$$\nabla^2 [\mathcal{E}^*(\vec{r}, t) \mathcal{E}(\vec{r}', t')] = \frac{1}{C^2} \frac{\partial^2}{\partial t^2} [\mathcal{E}^*(\vec{r}, t) \mathcal{E}(\vec{r}', t')]. \quad (6.51)$$

Taking the ensemble average of equation (6.51) over all realisations of the field, this wave equation becomes

$$\nabla^2 \Gamma(\vec{r}, \vec{r}'; t, t') = \frac{1}{C^2} \frac{\partial^2}{\partial t^2} \Gamma(\vec{r}, \vec{r}'; t, t'), \quad (6.52)$$

and, similarly, using $\square' = \nabla'^2 - \frac{1}{C^2} \frac{\partial^2}{\partial t'^2}$ we have another wave equation given by

$$\nabla'^2 \Gamma(\vec{r}, \vec{r}'; t, t') = \frac{1}{C^2} \frac{\partial^2}{\partial t'^2} \Gamma(\vec{r}, \vec{r}'; t, t'). \quad (6.53)$$

Equations (6.52) and (6.53) tell us directly that the coherence functions between two spacetime points evolve in time following a wave-type equation, separately, with respect to each pair of spacetime coordinates - a result that is, without doubt, remarkable. We can take this result even further and study this propagation in the next section.




Figure 6.3: Propagation of the electric field from a plane $Z = 0$ to a plane Z , where each point of the wavefront carries its mean polarisation ellipse.

6.4.3 Propagation of coherence and the direction of time: the van CittertZernike theorem

As we know, the electric field obeys a wave-type equation and the orientation of the fields is given by the polarisation of the field; in turn, as we saw in the diffraction chapter (see Section 5.2), at the classical level light fundamentally propagates as spherical waves, as illustrated in Figure 6.3, and at each point of the wavefront the components of the electric field oscillate randomly, but within the mean polarisation ellipse of their representative state. Mathematically we can represent this component as

$$\mathcal{E}_z(\vec{r}_1, \nu) = \frac{i}{\lambda} \int_{\Sigma} \mathcal{E}_0(\vec{r}, \nu) \frac{e^{ik\|\vec{r}_1 - \vec{r}\|}}{\|\vec{r}_1 - \vec{r}\|} \cos(\hat{z}; \vec{r}_1 - \vec{r}) d\sigma. \quad (6.54)$$

Using equation (6.47) we have

$$\mathcal{E}_z(\vec{r}_1, t) = \frac{1}{i} \int_{\mathbb{R}^+} \frac{\nu}{V} \int_{\Sigma} \mathcal{E}_0(\vec{r}, \nu) \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r})}{\|\vec{r}_1 - \vec{r}\|} e^{i2\pi\nu\left(t + \frac{\|\vec{r}_1 - \vec{r}\|}{V}\right)} d\sigma d\nu, \quad (6.55)$$

where V is the propagation velocity of the envelope. Rewriting a little more, we have

$$\mathcal{E}_z(\vec{r}_1, t) = \frac{1}{i} \int_{\Sigma} \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r})}{\|\vec{r}_1 - \vec{r}\|} \int_{\mathbb{R}^+} \frac{\nu}{V} \mathcal{E}_0(\vec{r}, \nu) e^{i2\pi\nu\left(t + \frac{\|\vec{r}_1 - \vec{r}\|}{V}\right)} d\nu d\sigma, \quad (6.56)$$

and, taking the narrowband approximation, we have

$$\mathcal{E}_z(\vec{r}_1, t) = \frac{1}{i\lambda} \int_{\Sigma} \mathcal{E}_0 \left(\vec{r}, t + \frac{\|\vec{r}_1 - \vec{r}\|}{V} \right) \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r})}{\|\vec{r}_1 - \vec{r}\|} d\sigma. \quad (6.57)$$

We can visualise graphically how we intend to propagate in Figure 6.4, where we observe that we are starting from the coherence function between two points $\vec{r}, \vec{r}'$ at instants t, t' , respectively, and carrying it to a coherence function at the points $\vec{r}_1, \vec{r}_2$ at two instants t_1, t_2 , respectively.




Figure 6.4: Propagation of the coherence function between two points $\vec{r}, \vec{r}'$ on the initial plane to the coherence between two points $\vec{r}_1, \vec{r}_2$ on the observation plane.

Starting from equation (6.57) we can compute the coherence function in its propagation as

$$\Gamma_z(\vec{r}_1, \vec{r}_2; \tau) = E \{ \mathcal{E}_z(\vec{r}_1, t) \mathcal{E}_z^*(\vec{r}_2, t - \tau) \}, \quad (6.58)$$

$$\Gamma_z(\vec{r}_1, \vec{r}_2; \tau) = E \left\{ \frac{1}{i\tilde{\lambda}} \int_{\Sigma} \mathcal{E}_0 \left(\vec{r}, t + \frac{\|\vec{r}_1 - \vec{r}\|}{V} \right) \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r})}{\|\vec{r}_1 - \vec{r}\|} d\sigma \right. \quad (6.59)$$

$$\left. \cdot \frac{1}{-i\tilde{\lambda}} \int_{\Sigma'} \mathcal{E}_0^* \left(\vec{r}', t - \tau + \frac{\|\vec{r}_2 - \vec{r}'\|}{V} \right) \frac{\cos(\hat{z}; \vec{r}_2 - \vec{r}')}{\|\vec{r}_2 - \vec{r}'\|} d\sigma' \right\}, \quad (6.60)$$

$$\Gamma_z(\vec{r}_1, \vec{r}_2; \tau) = \frac{1}{\tilde{\lambda}^2} \int_{\Sigma} \int_{\Sigma'} E \left\{ \mathcal{E}_0 \left(\vec{r}, t + \frac{\|\vec{r}_1 - \vec{r}\|}{V} \right) \mathcal{E}_0^* \left(\vec{r}', t - \tau + \frac{\|\vec{r}_2 - \vec{r}'\|}{V} \right) \right\} \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r}) \cos(\hat{z}; \vec{r}_2 - \vec{r}')}{\|\vec{r}_2 - \vec{r}'\| \cdot \|\vec{r}_1 - \vec{r}\|} d\sigma d\sigma' \quad (6.61)$$

$$\Gamma_z(\vec{r}_1, \vec{r}_2; \tau) = \frac{1}{\tilde{\lambda}^2} \int_{\Sigma} \int_{\Sigma'} \Gamma_0 \left(\vec{r}, \vec{r}'; \tau + \frac{\|\vec{r}_1 - \vec{r}\| + \|\vec{r}_2 - \vec{r}'\|}{V} \right) \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r}) \cos(\hat{z}; \vec{r}_2 - \vec{r}')}{\|\vec{r}_2 - \vec{r}'\| \cdot \|\vec{r}_1 - \vec{r}\|} d\sigma d\sigma' \quad (6.62)$$

The propagation of the random oscillations across n realisations can be visualised in Figure 6.5.

We are going to impose the following condition on the coherence function

$$\Gamma(\vec{r}, \vec{r}'; \tau) = \Gamma(\vec{r}'; \tau) \delta(\vec{r} - \vec{r}'), \quad (6.63)$$

which, in a few words, means that, spatially speaking, the distribution representing the coherence function is point-localised and weighted by the mutual coherence at the point $\vec{r}'$. Imposing this on equation (6.62) we have

$$\Gamma_z(\vec{r}_1, \vec{r}_2; \tau) = \frac{1}{\tilde{\lambda}^2} \int_{\Sigma} \Gamma_0 \left(\vec{r}; \tau + \frac{\|\vec{r} - \vec{r}_1\| - \|\vec{r} - \vec{r}_2\|}{V} \right) \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r}) \cos(\hat{z}; \vec{r}_2 - \vec{r})}{\|\vec{r}_2 - \vec{r}\| \cdot \|\vec{r}_1 - \vec{r}\|} d\sigma. \quad (6.64)$$

In nature we will not always have narrowband sources; however, we can resort to another argument, namely that most instruments do not have an infinite or broad range but a short one - or, in other words, measurement instruments measure a narrow part of the spectrum. Therefore, including the narrowband condition we have

$$\Gamma_z(\vec{r}_1, \vec{r}_2) e^{i2\pi\tilde{\nu}\tau} = \frac{1}{\tilde{\lambda}^2} \int_{\Sigma} \Gamma_0(\vec{r}) e^{i2\pi\tilde{\nu} \left(\tau + \frac{\|\vec{r} - \vec{r}_1\| - \|\vec{r} - \vec{r}_2\|}{V} \right)} \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r}) \cos(\hat{z}; \vec{r}_2 - \vec{r})}{\|\vec{r}_2 - \vec{r}\| \cdot \|\vec{r}_1 - \vec{r}\|} d\sigma. \quad (6.65)$$

Given the coherence time defined by $\tau_C = \int_{\mathbb{R}^+} |\gamma(\vec{r}, \tau)|^2 d\tau$, and if we impose a condition of very small times, smaller than a coherence time $\tau \ll \tau_C$, we have

$$\Gamma_z(\vec{r}_1, \vec{r}_2) = \frac{1}{\tilde{\lambda}^2} \int_{\Sigma} I_0(\vec{r}) e^{i2\pi \left(\frac{\|\vec{r} - \vec{r}_1\| - \|\vec{r} - \vec{r}_2\|}{V\tilde{\nu}} \right)} \frac{\cos(\hat{z}; \vec{r}_1 - \vec{r}) \cos(\hat{z}; \vec{r}_2 - \vec{r})}{\|\vec{r}_2 - \vec{r}\| \cdot \|\vec{r}_1 - \vec{r}\|} d\sigma. \quad (6.66)$$




Figure 6.5: Evolution of the realisations upon propagating the electric field.

Taking the far-field approximation we can simplify the problem as follows

$$\Gamma_z(\vec{r}_1, \vec{r}_2) = \frac{1}{\tilde{\lambda}^2 z^2} \int_{\Sigma} I_0(\vec{r}) e^{i2\pi \left(\frac{\|\vec{r}-\vec{r}_1\| - \|\vec{r}-\vec{r}_2\|}{V/\tilde{\nu}} \right)} d\sigma. \quad (6.67)$$

We may now make a Taylor-series expansion, the same one performed in the diffraction chapter. In this spirit the following terms become

$$\|\vec{r} - \vec{r}_1\| = (z^2 + (x - x_1)^2 + (y - y_1)^2)^{1/2} = z \left(1 + \frac{(x - x_1)^2 + (y - y_1)^2}{z^2} \right)^{1/2} \approx z + \frac{(x - x_1)^2 + (y - y_1)^2}{2z}, \quad (6.68)$$

$$\|\vec{r} - \vec{r}_2\| \approx z + \frac{(x - x_2)^2 + (y - y_2)^2}{2z}. \quad (6.69)$$

Introducing this into integral (6.67) we obtain

$$\Gamma_z(\vec{r}_1, \vec{r}_2) = \frac{1}{\tilde{\lambda}^2 z^2} \int_{\mathbb{R}^2} I_0(x, y) e^{-\frac{i2\pi}{\tilde{\lambda}z}(xx_1 + yy_1)} e^{\frac{i2\pi}{\tilde{\lambda}z}(xx_2 + yy_2)} e^{-\frac{i\pi}{\tilde{\lambda}z}(x_1^2 + y_1^2)} dx dy. \quad (6.70)$$

Let us define the vectors $\vec{S}_1 = (x_1, y_1)$, $\vec{S}_2 = (x_2, y_2)$ and $\vec{S} = (x, y)$. Then

$$\Gamma_z(\vec{S}_1, \vec{S}_2) = \frac{e^{i\frac{\pi}{\tilde{\lambda}z}(\|\vec{S}_1\|^2 - \|\vec{S}_2\|^2)}}{\tilde{\lambda}^2 z^2} \int_{\mathbb{R}^2} I_0(\vec{S}) e^{-i\frac{2\pi}{\tilde{\lambda}z}\vec{S} \cdot (\vec{S}_1 - \vec{S}_2)} dS. \quad (6.71)$$

If we set $\vec{S}_2 = \vec{0}$ we arrive at the Fourier integral

$$\Gamma_z(\vec{S}_1) = \frac{e^{i\frac{\pi}{\tilde{\lambda}z}\|\vec{S}_1\|^2}}{\tilde{\lambda}^2 z^2} \int_{\mathbb{R}^2} \Gamma_0(\vec{S}) e^{-i\frac{2\pi}{\tilde{\lambda}z}\vec{S} \cdot \vec{S}_1} dS, \quad (6.72)$$

which, basically, means that the correlations, under all the approximations we have made, propagate by means of a Fourier integral; or, what is the same, the van CittertZernike theorem tells us that coherence diffracts - it propagates naturally as a diffraction integral (see Figure 6.6).

How can we tell in which direction time flows? This may sound like a question rather out of context for these notes, but if we could travel back to the golden age of Boltzmann, we would find a heated debate of the time on how statistical physics gives a direction to time, an *arrow*. Paraphrasing Boltzmann, it is said that *time advances in the direction in which correlations increase*. If we look at the van CittertZernike theorem, we can start from two completely uncorrelated sources but, as they propagate, they gradually become correlated; time grows and so does coherence, until a point is reached at which the radiation of those sources is coherent. This is, without doubt, one of the strongest and most beautiful results we can find in statistical physics.




Figure 6.6: The mutual coherence propagated to the far field is the Fourier transform of the intensity distribution $\Gamma_0(\vec{S})$ of the source.

Bibliography

- Abbe, E. (1873). Beiträge zur theorie des mikroskops und der mikroskopischen wahrnehmung. *Archiv für Mikroskopische Anatomie*, 9(1):413–468.
- Abbott, B. P. et al. (2016). Observation of gravitational waves from a binary black hole merger. *Physical Review Letters*, 116(6):061102.
- Almeida, L. B. (1994). The fractional fourier transform and time-frequency representations. *IEEE Transactions on Signal Processing*, 42(11):3084–3091.
- Azzam, R. M. A. and Bashara, N. M. (1977). *Ellipsometry and Polarized Light*. North-Holland, Amsterdam.
- Bargmann, V. (1961). On a hilbert space of analytic functions and an associated integral transform. part i. *Communications on Pure and Applied Mathematics*, 14(3):187–214.
- Baumgratz, T., Cramer, M., and Plenio, M. B. (2014). Quantifying coherence. *Physical Review Letters*, 113(14):140401.
- Bernardo, L. M. and Soares, O. D. D. (1996). Optical fractional fourier transforms with complex orders. *Applied Optics*, 35(17):3163–3166.
- Bernoulli, D. (1753). Réflexions et éclaircissemens sur les nouvelles vibrations des cordes. *Histoire de l'Académie Royale des Sciences et Belles-Lettres de Berlin*, 9:147–172.
- Bernoulli, J. (1713). *Ars Conjectandi*. Thurnisius Brothers, Basel.
- Białynicki-Birula, I. (1994). On the wave function of the photon. *Acta Physica Polonica A*, 86(1):97–116.
- Birkhoff, G. D. (1931). Proof of the ergodic theorem. *Proceedings of the National Academy of Sciences of the United States of America*, 17(12):656–660.
- Born, M. and Wolf, E. (1999). *Principles of Optics: Electromagnetic Theory of Propagation, Interference and Diffraction of Light*. Cambridge University Press, 7th edition.
- Brewster, D. (1815). On the laws which regulate the polarisation of light by reflexion from transparent bodies. *Philosophical Transactions of the Royal Society of London*, 105:125–159.
- Candan, Ç., Kutay, M. A., and Ozaktas, H. M. (2000). The discrete fractional fourier transform. *IEEE Transactions on Signal Processing*, 48(5):1329–1337.
- Cantor, G. (1872). Über die ausdehnung eines satzes aus der theorie der trigonometrischen reihen. *Mathematische Annalen*, 5(1):123–132.

- Cariolaro, G., Erseghe, T., Kraniuskas, P., and Laurenti, N. (1998). A unified framework for the fractional fourier transform. *IEEE Transactions on Signal Processing*, 46(12):3206–3219.
- Chandrasekhar, S. (1950). *Radiative Transfer*. Clarendon Press, Oxford.
- Collins, S. A. (1970). Lens-system diffraction integral written in terms of matrix optics. *Journal of the Optical Society of America*, 60(9):1168–1177.
- Condon, E. U. (1937). Immersion of the fourier transform in a continuous group of functional transformations. *Proceedings of the National Academy of Sciences of the United States of America*, 23(3):158–164.
- Cooley, J. W. and Tukey, J. W. (1965). An algorithm for the machine calculation of complex fourier series. *Mathematics of Computation*, 19(90):297–301.
- d’Alembert, J. l. R. (1747). Recherches sur la courbe que forme une corde tendue mise en vibration. *Histoire de l’Académie Royale des Sciences et Belles-Lettres de Berlin*, 3:214–219.
- de Bruijn, N. G. (1973). A theory of generalized functions, with applications to wigner distribution and weyl correspondence. *Nieuw Archief voor Wiskunde*, 21:205–280.
- Dirichlet, P. G. L. (1829). Sur la convergence des séries trigonométriques qui servent à représenter une fonction arbitraire entre des limites données. *Journal für die reine und angewandte Mathematik*, 4:157–169.
- Doob, J. L. (1953). *Stochastic Processes*. John Wiley and Sons, New York.
- Duffieux, P.-M. (1946). *L’Intégrale de Fourier et ses Applications à l’Optique*. Faculté des Sciences, Besançon.
- Euler, L. (1748). Sur la vibration des cordes. *Histoire de l’Académie Royale des Sciences et Belles-Lettres de Berlin*, 4:69–85.
- Fabry, C. and Perot, A. (1899). Théorie et applications d’une nouvelle méthode de spectroscopie interférentielle. *Annales de Chimie et de Physique*, 16:115–144.
- Feynman, R. P., Leighton, R. B., and Sands, M. (1965). *The Feynman Lectures on Physics, Volume III: Quantum Mechanics*. Addison-Wesley, Reading, MA.
- Fourier, J. (1822). *Théorie Analytique de la Chaleur*. Firmin Didot, Père et Fils, Paris.
- Fresnel, A.-J. (1819). Mémoire sur la diffraction de la lumière. In *Mémoires de l’Académie des Sciences*, volume 5, pages 339–475. Académie Royale des Sciences, Paris.
- Galton, F. (1888). Co-relations and their measurement, chiefly from anthropometric data. *Proceedings of the Royal Society of London*, 45:135–145.
- Gauss, C. F. (1809). *Theoria Motus Corporum Coelestium in Sectionibus Conicis Solem Ambientium*. Perthes and Besser, Hamburg.

- Glauber, R. J. (1963). The quantum theory of optical coherence. *Physical Review*, 130(6):2529–2539.
- Goodman, J. W. (2005). *Introduction to Fourier Optics*. Roberts and Company Publishers, 3rd edition.
- Grimaldi, F. M. (1665). *Physico-mathesis de lumine, coloribus, et iride, aliisque annexis libri duo*. Vittorio Bonati, Bologna.
- Hanbury Brown, R. and Twiss, R. Q. (1956). Correlation between photons in two coherent beams of light. *Nature*, 177:27–29.
- Hermite, C. (1864). Sur un nouveau développement en série de fonctions. *Comptes Rendus de l'Académie des Sciences de Paris*, 58:93–100.
- Hertz, H. (1893). *Electric Waves: Being Researches on the Propagation of Electric Action with Finite Velocity Through Space*. Macmillan and Company.
- Herzberger, M. (1966). Optics from euclid to huygens. *Applied Optics*, 5(9):1383–1393.
- Hopkins, H. H. (1950). *The Wave Theory of Aberrations*. Clarendon Press, Oxford.
- Hurwitz, H. and Jones, R. C. (1941). A new calculus for the treatment of optical systems. ii. proof of three general equivalence theorems. *Journal of the Optical Society of America*, 31(7):493–499.
- Huygens, C. (1690). *Traité de la Lumière*. Pierre van der Aa, Leiden.
- Jones, R. C. (1941). A new calculus for the treatment of optical systems. i. description and discussion of the calculus. *Journal of the Optical Society of America*, 31(7):488–493.
- Kerr, F. H. (1988a). A distributional approach to namias' fractional fourier transforms. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 108(1–2):133–143.
- Kerr, F. H. (1988b). Namias' fractional fourier transforms on l^2 and applications to differential equations. *Journal of Mathematical Analysis and Applications*, 136(2):404–418.
- Khinchin, A. (1934). Korrelationstheorie der stationären stochastischen prozesse. *Mathematische Annalen*, 109(1):604–615.
- Kingslake, R. and Johnson, R. B. (2010). *Lens Design Fundamentals*. Academic Press, Burlington, 2 edition.
- Kirchhoff, G. (1883). Zur theorie der lichtstrahlen. *Annalen der Physik*, 254(4):663–695.
- Kober, H. (1939). Wurzeln aus der hankel-, fourier- und aus anderen stetigen transformationen. *The Quarterly Journal of Mathematics*, 10(1):45–59.
- Kolmogorov, A. N. (1933). *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Julius Springer, Berlin.
- Laplace, P.-S. (1812). *Théorie Analytique des Probabilités*. Courcier, Paris.

- Lebesgue, H. (1902). Intégrale, longueur, aire. *Annali di Matematica Pura ed Applicata*, 7(1):231–359.
- Lebesgue, H. (1903). Sur les séries trigonométriques. *Annales Scientifiques de l'École Normale Supérieure*, 20:453–485.
- Lohmann, A. W. (1993). Image rotation, wigner rotation, and the fractional fourier transform. *Journal of the Optical Society of America A*, 10(10):2181–2186.
- Malus, É.-L. (1809). Sur une propriété de la lumière réfléchie. *Mémoires de Physique et de Chimie de la Société d'Arcueil*, 2:143–158.
- Mandel, L. and Wolf, E. (1995). *Optical Coherence and Quantum Optics*. Cambridge University Press.
- Maxwell, J. C. (1865). A dynamical theory of the electromagnetic field. *Philosophical Transactions of the Royal Society of London*, 155:459–512.
- McBride, A. C. and Kerr, F. H. (1987). On namias's fractional fourier transforms. *IMA Journal of Applied Mathematics*, 39(2):159–175.
- McMaster, W. H. (1954). Polarization and the stokes parameters. *American Journal of Physics*, 22(6):351–362.
- Mehler, F. G. (1866). Ueber die entwicklung einer function von beliebig vielen variablen nach laplaceschen functionen höherer ordnung. *Journal für die reine und angewandte Mathematik*, 66:161–176.
- Mendlovic, D. and Ozaktas, H. M. (1993). Fractional fourier transforms and their optical implementation: I. *Journal of the Optical Society of America A*, 10(9):1875–1881.
- Michelson, A. A. (1881). The relative motion of the earth and the luminiferous ether. *American Journal of Science*, 22(128):120–129.
- Michelson, A. A. and Morley, E. W. (1887). On the relative motion of the earth and the luminiferous ether. *American Journal of Science*, 34(203):333–345.
- Michelson, A. A. and Pease, F. G. (1921). Measurement of the diameter of alpha orionis with the interferometer. *The Astrophysical Journal*, 53:249–259.
- Mie, G. (1908). Beiträge zur optik trüber medien, speziell kolloidaler metallösungen. *Annalen der Physik*, 330(3):377–445.
- Moshinsky, M. and Quesne, C. (1971). Linear canonical transformations and their unitary representations. *Journal of Mathematical Physics*, 12(8):1772–1780.
- Mustard, D. (1996). The fractional fourier transform and the wigner distribution. *The Journal of the Australian Mathematical Society Series B*, 38(2):209–219.

- Namias, V. (1980). The fractional order fourier transform and its application to quantum mechanics. *IMA Journal of Applied Mathematics*, 25(3):241–265.
- Newton, I. (1704). *Opticks: or, A Treatise of the Reflexions, Refractions, Inflexions and Colours of Light*. Samuel Smith and Benjamin Walford, London.
- Nikodym, O. (1930). Sur une généralisation des intégrales de m. j. radon. *Fundamenta Mathematicae*, 15(1):131–179.
- Ozaktas, H. M. and Mendlovic, D. (1993). Fractional fourier transforms and their optical implementation: Ii. *Journal of the Optical Society of America A*, 10(12):2522–2531.
- Ozaktas, H. M. and Mendlovic, D. (1995). Fractional fourier optics. *Journal of the Optical Society of America A*, 12(4):743–751.
- Ozaktas, H. M., Zalevsky, Z., and Kutay, M. A. (2001). *The Fractional Fourier Transform with Applications in Optics and Signal Processing*. John Wiley and Sons, Chichester.
- Pearson, K. (1896). Mathematical contributions to the theory of evolution. iii. regression, heredity, and panmixia. *Philosophical Transactions of the Royal Society of London. Series A*, 187:253–318.
- Pellat-Finet, P. (1994). Fresnel diffraction and the fractional-order fourier transform. *Optics Letters*, 19(18):1388–1390.
- Pellat-Finet, P. (2009). *Optique de Fourier: Théorie Métaaxiale et Fractionnaire*. Springer, Paris.
- Pellat-Finet, P. and Bonnet, G. (1994). Fractional order fourier transform and fourier optics. *Optics Communications*, 111(1–2):141–154.
- Perrin, F. (1942). Polarization of light scattered by isotropic opalescent media. *The Journal of Chemical Physics*, 10(7):415–427.
- Plancherel, M. (1910). Contribution à l’étude de la représentation d’une fonction arbitraire par des intégrales définies. *Rendiconti del Circolo Matematico di Palermo*, 30(1):289–335.
- Poincaré, H. (1892). *Théorie Mathématique de la Lumière II*. Georges Carré, Paris.
- Riemann, B. (1867). Ueber die darstellbarkeit einer function durch eine trigonometrische reihe. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen*, 13:87–132.
- Santos, E. A., Castro, F., and Torres, R. (2018). Huygens-fresnel principle: Analyzing consistency at the photon level. *Physical Review A*, 97(4):043853.
- Schwarzschild, K. (1905). Untersuchungen zur geometrischen optik i–iii. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, 4(1–3):1–54.
- Sejdić, E., Djurović, I., and Stanković, L. (2011). Fractional fourier transform as a signal processing tool: An overview of recent developments. *Signal Processing*, 91(6):1351–1369.

- Shih, C.-C. (1995a). Fractionalization of fourier transform. *Optics Communications*, 118(5–6):495–498.
- Shih, C.-C. (1995b). Optical interpretation of a complex-order fourier transform. *Optics Letters*, 20(10):1178–1180.
- Sommerfeld, A. (1896). Mathematische theorie der diffraction. *Mathematische Annalen*, 47(2-3):317–374.
- Sommerfeld, A. (1954). *Optics: Lectures on Theoretical Physics, Volume IV*. Academic Press, New York.
- Stokes, G. G. (1852). On the composition and resolution of streams of polarized light from different sources. *Transactions of the Cambridge Philosophical Society*, 9:399–416.
- Sudarshan, E. C. G. (1963). Equivalence of semiclassical and quantum mechanical descriptions of statistical light beams. *Physical Review Letters*, 10(7):277–279.
- Titchmarsh, E. C. (1937). *Introduction to the Theory of Fourier Integrals*. Clarendon Press, Oxford.
- van Cittert, P. H. (1934). Die wahrscheinliche schwingungsverteilung in einer von einer lichtquelle direkt oder mittels einer linse beleuchteten ebene. *Physica*, 1(1-6):201–210.
- Vitali, G. (1905). *Sul Problema della Misura dei Gruppi di Punti di una Retta*. Tipografia Gamberini e Parmeggiani, Bologna.
- Welford, W. T. (1986). *Aberrations of Optical Systems*. Adam Hilger, Bristol.
- Wiener, N. (1929). Hermitian polynomials and fourier analysis. *Journal of Mathematics and Physics*, 8(1–4):70–73.
- Wiener, N. (1930). Generalized harmonic analysis. *Acta Mathematica*, 55:117–258.
- Wiener, N. (1933). *The Fourier Integral and Certain of its Applications*. Cambridge University Press, Cambridge.
- Wolf, E. (2007). *Introduction to the Theory of Coherence and Polarization of Light*. Cambridge University Press, Cambridge.
- Wolf, K. B. (1979). *Integral Transforms in Science and Engineering*. Plenum Press, New York.
- Young, T. (1802). The bakerian lecture: On the theory of light and colours. *Philosophical Transactions of the Royal Society of London*, 92:12–48.
- Young, T. (1804). The bakerian lecture: Experiments and calculations relative to physical optics. *Philosophical Transactions of the Royal Society of London*, 94:1–16.
- Young, T. (1807). *A Course of Lectures on Natural Philosophy and the Mechanical Arts*. Joseph Johnson, London.

- Yu, N., Genevet, P., Kats, M. A., Aieta, F., Tetienne, J.-P., Capasso, F., and Gaburro, Z. (2011). Light propagation with phase discontinuities: generalized laws of reflection and refraction. *Science*, 334(6054):333–337.
- Zernike, F. (1938). The concept of degree of coherence and its application to optical problems. *Physica*, 5(8):785–795.