

1. Administrative & Study Identification

Full Study Title ELRANATAMAB POST TRIAL ACCESS: AN OPEN-LABEL, SINGLE-ARM STUDY FOR PARTICIPANTS WITH MULTIPLE MYELOMA CONTINUING FROM PFIZER-SPONSORED ELRANATAMAB CLINICAL STUDIES
Short Title/Acronym Elranatamab Post Trial Access Study for Participants With Multiple Myeloma (MM) (MagnetisMM15) **Protocol Number** C1071015 / 2023-505200-33-00 **NCT Number** NCT06057402
Sponsor Name Pfizer (Company: Pfizer) **Investigational Product** Elranatamab (Trade Name: Elrexfio) **Phase of Development** Phase 4 (PHASE4) **Trial Status** Recruiting **Medical Monitor** Pfizer CT.gov Call Center / Principal Investigator

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To provide continued post-trial access to elranatamab for participants with Multiple Myeloma who have completed a qualifying Parent Study and are still deriving clinical benefit. **Secondary Objectives:** To monitor the long-term safety and tolerability of elranatamab, specifically focusing on the incidence of adverse events (AEs) and serious adverse events (SAEs) leading to discontinuation.

2.2 Background & Rationale This is a post-trial access (PTA) open-label, single-arm study designed to ensure continued treatment for multiple myeloma patients who successfully completed a Pfizer-sponsored parent study of elranatamab and continue to derive clinical benefit from elranatamab monotherapy. Because elranatamab targets BCMA on MM cells and CD3 on T cells, it is crucial to provide ongoing therapy without interruption until the drug becomes commercially accessible in the participant's country, disease progression occurs, or unacceptable toxicity is observed.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm **Blinding:** Open-label (Masking: None) **Randomization:** N/A

3.2 Planned Enrollment & Duration Trial Status: Recruiting (Status Description: Currently recruiting participants) **Target \$N\$:** 80 participants (Estimated) **Number of Sites:** Multicenter (Global) **Estimated Study Start:** 03-Oct-2023 (Actual) **Estimated Primary Completion:** 22-Feb-2032

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Participants must agree to follow the reproductive criteria as outlined in the protocol. 2. Participants have completed a qualifying Parent Study, were still receiving elranatamab when the Parent Study terminated or completed, and are deriving clinical benefit from elranatamab (as determined by the investigator).

4.2 Exclusion Criteria 1. Any medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior or laboratory abnormality that may increase the risk of study participation or, in the investigator's judgment, make the participant inappropriate for the study. 2. Participants not previously enrolled or who have discontinued study treatment in a Parent Study are ineligible for participation in this study.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Investigational Product: Elranatamab
Preferred UMLS Name / Drug Preferred Name: Elranatamab **Trade Name:** Elrexfio **EMA URL:** https://www.ema.europa.eu/en/documents/product-information/elrexfio-epar-product-information_en.pdf
Dosage/Frequency: Elranatamab administered at the previously established efficacious dose/schedule from the parent study.
Route of Administration: Subcutaneous injection. **Duration of Treatment:** Until disease progression, unacceptable toxicity, withdrawal of consent, study termination, or until elranatamab becomes commercially accessible in the participant's country.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant medications for co-morbidities as prescribed by the treating physician and permitted in the parent trial. **Prohibited:** Other investigational drugs or anti-myeloma therapies that would confound the safety assessments of elranatamab.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening/Transition	Day -14 to 0 Informed Consent, verification of parent study completion, confirmation of continued clinical benefit. Treatment Period Ongoing Administration of Elranatamab, routine safety monitoring, AE/SAE assessment. End of Study Follow-up Final safety assessment, tracking of AEs/SAEs for a minimum of 90 days after the last dose of study drug.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. **Incidence of nonserious adverse events (AEs) leading to permanent discontinuation.** **Measurement Method:** Clinical safety assessments measured up to a minimum of 90 days after the last dose of the study drug. 2. **Incidence of serious adverse events (SAEs).** **Measurement Method:** Monitored up to a minimum of 90 days after the last dose of the study drug.

7.2 Secondary & Exploratory Endpoints Not explicitly specified in the primary outcome structured data, but standard long-term safety and tolerability monitoring continues throughout the duration of the post-trial access.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard continuous monitoring at each site visit during the treatment phase and extending 90 days post-last dose. **Serious Adverse Events (SAEs):** Reported promptly according to regulatory timelines (typically 24 hours), closely monitoring for severe toxicities. **Data Monitoring Committee (DMC):** Internal safety monitoring mechanisms managed by Pfizer.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants who receive at least one dose of elranatamab in the PTA study). **Sample Size Justification:** The targeted enrollment ($N = 80$) is not based on statistical power calculations but rather on the estimated number of eligible participants transitioning from the qualifying Pfizer-sponsored Parent Studies. **Handling of Missing Data:** Summarized using descriptive statistics; last observation carried forward (LOCF) for relevant safety data as required.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring visits to ensure appropriate AE/SAE reporting and continued assessment of clinical benefit. **Audit Trail:** Details on eCRF locking and compliance checks with standard post-trial access protocol regulations.

11. Study Identifiers & Governance

Primary ID: C1071015 **Posting ID:** 1010680919594791619 **Registry Number:** NCT06057402 **Sponsor/Lead Organization:** Pfizer **Study**

Title: Elranatamab Post Trial Access Study for Participants With Multiple Myeloma (MM) **Official Regulatory Title:** ELRANATAMAB POST TRIAL ACCESS: AN OPEN-LABEL, SINGLE-ARM STUDY FOR PARTICIPANTS WITH MULTIPLE MYELOMA CONTINUING FROM PFIZER-SPONSORED ELRANATAMAB CLINICAL STUDIES **Source Level:** 5 **Criteria Type:** PHASE_REQUIREMENT (Criteria Text: Trial must be in PHASE4)

12. Clinical Framework (Multiple Myeloma Specific)

Primary Condition: Multiple Myeloma (MM) **Disease Area / Preferred Name:** Multiple Myeloma **Semantic Type:** Neoplastic Process
Definition: A bone marrow-based plasma cell neoplasm characterized by a serum monoclonal protein and skeletal destruction with osteolytic lesions, pathological fractures, bone pain, hypercalcemia, and anemia. Clinical variants include non-secretory myeloma, smoldering myeloma, indolent myeloma, and plasma cell leukemia. (WHO, 2001) **Condition Codes:** * **MeSH Code:** D009101 * **HPO Code:** HP:0006775 * **SNOMED ID:** 109989006 * **SNOMED Hierarchy:** - 269475001 | Malignant neoplasm of lymphoid, hematopoietic and/or related tissue | Malignant neoplasm of lymphoid, hematopoietic and/or related tissue (disorder) - 415111003 | Plasma cell neoplasm | Plasma cell neoplasm (disorder) - 1162576007 | Plasma cell myeloma | Plasma cell myeloma (morphologic abnormality) - 64572001 | Disease | Disease (disorder)

Diagnosis Threshold: Confirmed MM deriving continued clinical benefit from elranatamab as determined by the investigator following a parent study. **Medical Methodology:** Post-trial access to BCMA-CD3 bispecific antibody. **Optimization Target:** Sustained disease control and maintenance of clinical benefit.

13. Study Procedures & Methodology

Management Pathway: Continued elranatamab monotherapy for relapsed/refractory multiple myeloma transitioning from prior clinical trials. **Education Protocol (HCP):** AE management and protocol transition training. **Education Protocol (Patient):** Informed consent detailing the rationale and logistics of post-trial drug access. **Quality Audit Mechanism:** Safety reporting compliance and tracking of treatment discontinuation criteria.

14. Observation & Follow-Up Schedule

Total Duration: Expected treatment through February 2032 or commercial availability. **Onsite Visit Cadence:** Aligned with the treatment administration schedule. **Remote Monitoring Frequency:** As required by local site protocols. **Checklist Review Interval:** Ongoing AE/SAE safety review up to 90 days post-treatment.

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					