

Clinical Study Summary Report (CSSR) Document Status: Draft |
Version: 0.1 | **Date:** 07-Apr-2026 **Trial Status:** RECRUITING
(Currently recruiting participants) | **Posting ID:**
1073500494040370013

1. Administrative & Study Identification

Full Study Title: A Long-Term Study to Learn About The Study Medicine Called Ritlecitinib in Children With Severe Alopecia Areata. **Short Title/Acronym:** Ritlecitinib Pediatric AA Long-Term Extension (Study B7981028) **Protocol Number:** B7981028 **NCT Number:** NCT07029828 **Sponsor/Company Name:** Pfizer **Investigational Product:** Ritlecitinib (Trade Name: Litfulo) (oral JAK3/TEC family kinase inhibitor) **EMA Product Information:** [Litfulo EPAR](#) **Phase of Development:** Phase 3 **Medical Monitor:** Pfizer Clinical Trial Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the long-term safety and tolerability of ritlecitinib in pediatric participants (ages 6 to 14) with severe alopecia areata (AA) who have completed the parent studies B7981027 or B7981031. **Secondary Objectives:** To evaluate the long-term efficacy of ritlecitinib, durability of response, and the effect of ritlecitinib on patient-centered outcomes (including psychological/psychosocial status and neuropsychological/cognitive development).

2.2 Background & Rationale Alopecia areata is an autoimmune condition causing unpredictable hair loss, carrying significant emotional and social impacts, especially in pediatric populations. While ritlecitinib has been shown to be effective in treating severe AA, there is a medical necessity to evaluate its prolonged use in children to answer critical questions about long-term safety, sustained hair regrowth, and overall impact on quality of life. This 3-year extension study ensures continued access to the investigational medication for children who participated in previous trials while generating robust longitudinal safety and efficacy data.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Long-term Extension) **Control Type:** Placebo-controlled (for threshold discontinuation criteria) and Active Dose-comparison **Blinding:** Double-blind **Randomization:** Randomized (Participants rolling over from placebo or specific parent trials are randomized at a 1:1 ratio to receive higher or lower dose ritlecitinib)

3.2 Planned Enrollment & Duration Target **\$N\$**: 140 pediatric patients (evaluable for primary analysis) **Number of Sites**: Multicenter (Global locations including US sites in FL, IN, OR, TX) **Estimated Study Start**: Late 2024 / Early 2025 **Estimated Primary Completion**: Up to 3 years (36 months) per participant

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Children aged 6 to 14 years. 2. Participants with alopecia areata (AA), including alopecia totalis [AT] and alopecia universalis [AU], who completed studies B7981027 or B7981031. 3. For participants originating from Study B7981031: At least 50% scalp hair loss due to AA (i.e., a SALT score of $\geq 50\%$) at both the Screening and Baseline visits. 4. Received varicella vaccination (2 doses) or have been previously infected by the varicella-zoster virus based on blood test reports.

4.2 Exclusion Criteria 1. **From Study B7981027 (≤ 30 days between last dose and first visit)**: Presence of safety events requiring permanent discontinuation. Non-safety discontinuations must have issues resolved prior to enrollment. 2. **From Study B7981031 or B7981027 (>30 days gap)**: Any safety events requiring permanent discontinuation. 3. Any present malignancies or history of malignancies or lymphoproliferative disorders. 4. Evidence of untreated or inadequately treated active or latent Mycobacterium tuberculosis (TB) infection. 5. History (one or more episodes) of severe or serious cytomegalovirus (CMV), herpes zoster (shingles), disseminated herpes simplex, or infection with hepatitis B virus (HBV) or hepatitis C virus (HCV).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Ritlecitinib higher dose or lower dose capsule, taken once a day (QD). Participants will be evaluated for treatment continuation criteria at Months 3, 6, 9, and 12 depending on their original study assignment. Treatment will be discontinued for those who do not meet the continuation criteria. **Route of Administration**: Oral (capsules provided in HDPE bottles). **Duration of Treatment**: Up to 3 years (36 months).

5.2 Concomitant & Prohibited Therapy Permitted: Standard pediatric supportive care. **Prohibited**: Other systemic immunosuppressants, biologics, and live-attenuated vaccines during the study period.

6. Study Timeline & Visit Schedule

| Visit ID | Window | Key Activities/Procedures | | :--- | :---
| :--- | | **Screening/Baseline** | Day 0 | Rollover from B7981027/
B7981031, Randomization, First Dose | | **Interim Visits** | Months 1
- 35 | 17 onsite clinic visits: Safety Labs, SALT scoring, PROs
(CDLQI, PROMIS, BRIEF2), AE tracking | | **End of Study** | Month 36
| Final Efficacy Assessment, Exit Interview, IP Accountability |

(Note: The study team will also call participants once a month over the phone between clinic visits.)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Long-term safety and tolerability of ritlecitinib over 36 months, defined by: 1. Incidence of treatment-emergent adverse events (TEAEs). 2. Incidence of Serious Adverse Events (SAEs) and AEs leading to permanent discontinuation from the study. **(Additional Measurement):** Clinical laboratory abnormalities evaluated during regular remote/onsite safety monitoring.

7.2 Secondary & Exploratory Endpoints * Long-term efficacy measured by Severity of Alopecia Tool (SALT) score: 1. Proportion of participants achieving absolute SALT score $\leq 10\%$. 2. Proportion of participants achieving absolute SALT score $\leq 20\%$. 3. Change from baseline (CFB) in SALT score. * Eyebrow and eyelash regrowth assessments. * Patient Global Impression of Change (PGI-C). * Patient-Reported Outcomes including PROMIS (Parent Proxy - Anxiety Short Form 8a and Depressive Symptoms Short Form 6a), BRIEF2 (Executive Function), and modified Children's Dermatology Life Quality Index (CDLQI).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous collection via 17 onsite clinic visits and monthly phone calls. Specific monitoring for opportunistic infections (e.g., herpes zoster), systemic toxicities, and emotional wellbeing. **Serious Adverse Events (SAEs):** Standard 24-hour reporting protocols for all severe, life-threatening events, and AEs leading to permanent study discontinuation. **Data Monitoring Committee (DMC):** Independent safety monitoring to oversee pediatric risk profiles continuously throughout the 3-year extension.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Analysis Set (all participants who receive at least 1 dose of the study drug); Efficacy Analysis Set (Intent-to-Treat rollover population). **Sample Size Justification:**

Determined by the number of eligible patients successfully completing parent phase 3 trials (Target \$N = 140\$ evaluable for primary analysis). **Handling of Missing Data:** Standard handling for long-term extensions, utilizing Multiple Imputation (MI) or Last Observation Carried Forward (LOCF) for efficacy endpoints depending on regional regulatory requirements.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Strict safety and compliance monitoring comprising 17 onsite visits over 3 years and intermediary remote phone check-ins every month. **Audit Trail:** Standardized eCRF locking, tracking of protocol-defined treatment discontinuation criteria, and efficacy threshold monitoring. **Study Phase**

Criteria: PHASE_REQUIREMENT (Trial must be in PHASE3)

11. Study Identifiers & Governance

Primary ID: B7981028 **Registry Number:** NCT07029828 **Posting ID:**

1073500494040370013 **Sponsor/Lead Organization:** Pfizer **Study**

Title: A Long-Term Study to Learn About The Study Medicine Called Ritlecitinib in Children With Severe Alopecia Areata. **Official**

Regulatory Title: A LONG-TERM, DOUBLE-BLIND EXTENSION STUDY TO INVESTIGATE THE SAFETY AND EFFICACY OF RITLECITINIB IN

PARTICIPANTS WITH SEVERE ALOPECIA AREATA WHO PREVIOUSLY COMPLETED STUDIES B7981027 OR B7981031

12. Clinical Framework (Alopecia Areata Specific)

Disease Area / Primary Condition: Severe Alopecia Areata (including Alopecia Totalis and Alopecia Universalis) **Condition**

Definition: Severe Alopecia Areata is an autoimmune disorder characterized by extensive, non-scarring hair loss, where the body's immune system mistakenly attacks its own hair follicles. This severity is typically defined by hair loss affecting more than 50% of the scalp surface area, or involvement of other body hair, including eyebrows, eyelashes, or progression to complete scalp hair loss (alopecia totalis) or total body hair loss (alopecia universalis). **Diagnosis Threshold:** $\geq 50\%$ scalp

hair loss (SALT ≥ 50) at parent study baseline **Medical**

Methodology: Oral selective dual JAK3/TEC family kinase inhibitor therapy

Clinical Classifications & Ontologies: * **Preferred UMLS Name:**

Severe Alopecia Areata * **Preferred Name:** Alopecia Areata Symptom Impact Scale, severity of symptoms subscale * **MeSH Code:** D000523

* **SNOMED ID:** 402513000 * **SNOMED Hierarchy:** 717088002 | Alopecia

13. Study Procedures & Methodology

Management Pathway: Long-term extension maintenance and safety tracking **Education Protocol (HCP):** Clinical guidelines for managing pediatric JAK inhibitor safety profiles. **Education Protocol (Patient):** Regular site visits and monthly telephone check-ins. **Quality Audit Mechanism:** Tracking of SALT score efficacy thresholds; standardized PRO psychological testing (PROMIS, BRIEF2, CDLQI) to assess emotional health.

14. Observation & Follow-Up Schedule

Total Duration: 36 months (3 years) **Onsite Visit Cadence:** 17 total clinic visits spaced across the 3-year period **Remote Monitoring Frequency:** Telephone calls once a month **Checklist Review Interval:** Regular assessment of treatment efficacy thresholds at Months 3, 6, 9, and 12 to determine continuation of study therapy.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead	
Statistician	[Name]	_____	[DD-MMM-YYYY]				