

1. Administrative & Study Identification

Full Study Title: A 16-WEEK, MULTICENTER, INTERVENTIONAL, PHASE 3, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, PARALLEL GROUP STUDY TO INVESTIGATE EFFICACY AND SAFETY OF ABROCITINIB IN CHILDREN 6 TO LESS THAN 12 YEARS OF AGE WITH MODERATE-TO-SEVERE ATOPIC DERMATITIS **Short Title/Acronym:** A Study of the Medicine Called Abrocitinib in Children 6 to Less Than 12 Years of Age With Moderate-to-Severe Eczema **Protocol Number:** B7451023 **NCT Number:** NCT06807268 **Posting ID:** 1075556539970939144 **Trial Status:** RECRUITING (Currently recruiting participants) **Sponsor/Company Name:** Pfizer **Investigational Product:** Abrocitinib (Trade Name: CIBINQO™) - Oral JAK inhibitor **Drug ATC Code:** D11AH08 **EMA Product Information:** [EMA URL](#) **Phase of Development:** Phase 3 **Medical Monitor:** Pfizer Clinical Trial Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To investigate the efficacy and safety of abrocitinib compared to placebo in children 6 to <12 years of age with moderate-to-severe atopic dermatitis (AD). **Secondary Objectives:** To evaluate long-term effectiveness, reduction in itch intensity (WI-NRS), patient-reported outcomes via electronic diaries, and overall safety and tolerability of the oral JAK inhibitor.

2.2 Background & Rationale Atopic dermatitis (eczema) is a chronic inflammatory skin condition common in children. Eczema is a form of dermatitis that is characterized by scaly, pruritic, erythematous lesions located on flexural surfaces [<https://orcid.org/0000-0002-0736-9199>, PMID:32412211]. It can severely impact sleep and quality of life. This study is being conducted to address the unmet medical need for pediatric patients aged 6 to <12 who have not had adequate results from topical medications (like steroid creams or non-steroid ointments) and to determine if the oral JAK inhibitor abrocitinib is a safe and effective systemic treatment option.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** Randomized (Parallel Group, 2:1 ratio of abrocitinib to placebo) **Phase Criteria:** Trial must be in PHASE3 (Criteria Type: PHASE_REQUIREMENT)

3.2 Planned Enrollment & Duration Targeted Enrollment (\$N\$): 150 participants **Number of Sites:** Multicenter (Global, including sites in the US and Japan) **Estimated Study Start:** 2023 **Estimated Primary Completion:** ~Mid-2028 (Results expected approximately 2 years, 1 month from March 2026)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Children aged 6 to <12 years at the time of informed consent/assent weighing ≥ 15 kg. No contraception methods are required for male participants. 2. A documented diagnosis of chronic AD for at least 1 year prior to screening (confirmed according to Hanifin and Rajka criteria), with moderate-to-severe AD at baseline (BSA $\geq 10\%$, VIGA ≥ 3 , EASI ≥ 16 , and WI-NRS ≥ 4). 3. Documented history (within 6 months) of inadequate response to treatment with topical medical therapy for AD (e.g., TCS and TCI) for at least 4 weeks, and are candidates for systemic therapy.

4.2 Exclusion Criteria 1. Skin infections requiring treatment with systemic antimicrobials within 2 weeks prior to Day 1, superficial skin infections within 1 week of Day 1, or history of systemic infection requiring hospitalization or parenteral antimicrobial therapy within 1 month prior to Day 1. 2. Infection with HIV, hepatitis B, and/or hepatitis C, or evidence of active TB or inadequately treated latent TB. History of disseminated herpes zoster or simplex. 3. History of conditions associated with thrombocytopenia, coagulopathy, malignancies (leukemia, lymphoma, sarcoma), immunodeficiency disorders, skeletal dysplasia, or retinal detachment. 4. Psychiatric condition including active suicidal ideation in the past year or suicidal behavior in the past 5 years. SDQ total score ≥ 17 requires MHP evaluation. 5. Previous administration of an investigational drug within 30 days or 5 half-lives (whichever is longer) of Day 1.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Abrocitinib test medicine or matching placebo taken by mouth once daily. **Route of Administration:** Oral **Duration of Treatment:** 16 weeks of assigned treatment per protocol, with the total study lasting about 24 weeks.

5.2 Concomitant & Prohibited Therapy Permitted: Background topical therapies (e.g., topical corticosteroids [TCS]) applied to affected areas at least once daily. **Prohibited:** Prior treatment with a systemic JAK inhibitor for AD; live attenuated vaccines within 6 weeks prior to Day 1, during treatment, or within 6 weeks after the last dose; concomitant use of strong

inhibitors and inducers of CYP2C19 enzymes, strong inducers of CYP2C9 enzymes, and P-gp substrates.

6. Study Timeline & Visit Schedule

| Visit ID | Window (Days) | Key Activities/Procedures | | :---
| :--- | :--- | | **Screening** | Day -28 to -1 | Informed Consent/
Assent, Confirm AD severity (vIGA, EASI, BSA), Screen for
infections (TB, HIV, Hep B/C) | | **Baseline** | Day 1 |
Randomization (2:1), First Dose, Issue electronic diary and
scratching tracking wristwatch to participant/caregiver | |
Interim | Weeks 1-15 | Daily e-diary questionnaires, Scratching
tracking, Safety Labs, Efficacy Assessment, Background topical
therapy compliance | | **End of Treatment** | Week 16 | Primary
Efficacy Assessment (vIGA and EASI), Safety Review | | **End of
Study** | Week 24 | Final Follow-up Visit, Exit Interview, IP
Accountability |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. **vIGA Response:** Response based on achieving Validated Investigator's Global Assessment (vIGA) score of clear (0) or almost clear (1) (on a 5-point scale) and a reduction from baseline of ≥ 2 points at Week 12. 2. **EASI-75 Response:** Response based on achieving $\geq 75\%$ improvement from baseline in the Eczema Area and Severity Index (EASI-75) at Week 12.

7.2 Secondary Endpoints 1. Change from Baseline (CFB) in the Worst Itch Numerical Rating Scale (WI-NRS) at Week 2. 2. Response based on achieving at least a 4-point improvement from baseline in the WI-NRS at Week 12. 3. Response based on achieving WI-NRS < 2 at Week 12.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Collected continuously via site visits and an electronic diary (resembling a cellular phone) provided to patients/caregivers to report daily symptom tracking, dosing reminders, and AEs. **Serious Adverse Events (SAEs):** Specific monitoring for serious systemic or localized infections (e.g., disseminated herpes zoster/simplex), thrombocytopenia, and psychiatric events. **Data Monitoring Committee (DMC):** Overseen by the sponsor (Pfizer) in accordance with Phase 3 pediatric study guidelines.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy endpoint comparisons (abrocitinib vs. placebo); Safety Set for adverse events. **Sample Size Justification:** Sized for a Phase 3 pediatric trial with 2:1 randomization to provide sufficient power to detect a significant difference in the proportion of vIGA and EASI-75 responders among a target enrollment of 150 patients. **Handling of Missing Data:** Standard handling for binary primary endpoints like vIGA 0/1 and EASI-75.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite monitoring visits combined with daily compliance checks of the electronic diary entries for dosing and symptom questionnaires. **Audit Trail:** Electronic diaries automatically log time-stamped questionnaire answers and dosing reminders; eCRF locking utilized for site data.

11. Study Identifiers & Governance

Primary ID: B7451023 **Registry Number:** NCT06807268 **Posting ID:** 1075556539970939144 **Trial Status:** RECRUITING (Currently recruiting participants) **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study of the Medicine Called Abrocitinib in Children 6 to Less Than 12 Years of Age With Moderate-to-Severe Eczema **Official Regulatory Title:** A 16-WEEK, MULTICENTER, INTERVENTIONAL, PHASE 3, RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED, PARALLEL GROUP STUDY TO INVESTIGATE EFFICACY AND SAFETY OF ABROCITINIB IN CHILDREN 6 TO LESS THAN 12 YEARS OF AGE WITH MODERATE-TO-SEVERE ATOPIC DERMATITIS

12. Clinical Framework (Atopic Dermatitis Specific)

Primary Condition / Disease Area: Eczema (Moderate-to-Severe Atopic Dermatitis) **Preferred UMLS Name:** Atopic Dermatitis **Semantic Type:** Disease or Syndrome **Ontology Classifications:** * **MeSH Code:** D004485 * **HPO Code:** HP:0000964 * **SNOMED ID:** 43116000 * **SNOMED Hierarchy:** 703938007 | Inflammatory dermatosis | Inflammatory dermatosis (disorder); 64572001 | Disease | Disease (disorder) **Source Level:** 5 **Diagnosis Threshold:** BSA $\geq 10\%$, vIGA ≥ 3 , EASI ≥ 16 , WI-NRS ≥ 4 **Medical Methodology:** Systemic therapy via oral JAK inhibitor + Background topical medications **Optimization Target:** vIGA of clear (0) or almost clear (1) and EASI-75 achievement

13. Study Procedures & Methodology

Management Pathway: Daily oral systemic targeted therapy administration integrated with standard topical therapy routines. **Scratch Tracking:** Participants will be asked to wear a tracking device on their wrist (that looks like a children's wristwatch) so investigators can identify when they are scratching their skin because of their atopic dermatitis. **Education Protocol:** Investigator training for standardized vIGA, EASI, and BSA scoring. Caregiver/patient training on utilizing the electronic diary and consistent application of background topical therapies. **Quality Audit Mechanism:** Daily e-diary compliance tracking and adherence alerts.

14. Observation & Follow-Up Schedule

Total Duration: ~24 weeks **Onsite Visit Cadence:** Baseline, regular interim intervals, Week 16 (End of Treatment), and Week 24 (End of Study) **Remote Monitoring Frequency:** Daily questionnaire and dosing tracking via cellular-phone-like e-diary, alongside continuous wrist-worn scratch tracking. **Checklist Review Interval:** Every clinical visit

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				