

1. Administrative & Study Identification

Full Study Title: A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Different Anticancer Agents in Advanced Cancers **Short Title/Acronym:** PF-08634404 Combination Study in Advanced Cancers **Protocol Number:** C6461020 **NCT Number:** NCT07227298 **Sponsor Name:** Pfizer **Investigational Product:** PF-08634404 in combination with Sigvotatug Vedotin or other agents **Phase of Development:** Phase 1b/2 **Medical Monitor:** Pfizer CT.gov Call Center (1-800-718-1021 / ClinicalTrials.gov_Inquiries@pfizer.com)

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To evaluate the safety, pharmacokinetics, and preliminary efficacy of PF-08634404 in combination with different anticancer agents in participants with advanced solid tumors, specifically assessing Dose Limiting Toxicities (DLTs) and adverse events. * **Secondary Objectives:** To evaluate the Objective Response Rate (ORR), characterize the pharmacokinetics (PK), and assess the immunogenicity of PF-08634404.

2.2 Background & Rationale This study is being done to learn more about a new medicine called PF-08634404 and how it works when used with other cancer medicines in people who have advanced solid tumors. An advanced solid tumor is a type of cancer that has spread beyond its original location and cannot be removed by surgery or cured with standard treatments. The combination therapy (e.g., with sigvotatug vedotin) is being explored to identify safe dosages and potential therapeutic benefits.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Single-arm (Open-label dose escalation and expansion) * **Blinding:** Open-label * **Randomization:** N/A

3.2 Planned Enrollment & Duration * **Target \$N\$:** ~138 participants * **Number of Sites:** Global multi-center (including locations in the US, Puerto Rico, and Japan) * **Estimated Study Start:** 17-Nov-2025 * **Estimated Primary Completion:** 08-Mar-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age $\geq$ 18 years with an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1. 2. Pathologically confirmed locally advanced (Stage IIIB/IIIC) or metastatic (Stage IV) squamous or non-squamous non-small cell

lung cancer (NSCLC) and are not a candidate for complete surgical resection and curative concurrent/sequential chemoradiotherapy. 3. Measurable disease based on RECIST v1.1 per investigator, and PD-L1 status available (Part B only: PD-L1 $\geq$ TPS 1%).

4.2 Exclusion Criteria 1. Known active CNS lesions, including brainstem, meningeal, or spinal cord metastases or compression; or Leptomeningeal disease. 2. Participants with known actionable genomic alterations (AGAs) including EGFR, ALK, ROS1, NTRK, BRAF, and MET, or active autoimmune diseases requiring systemic treatment within the past 2 years. 3. History of another malignancy within 3 years before the first dose of study intervention, or previous systemic anti-tumor therapy for locally advanced, unresectable, or metastatic NSCLC (with specified exceptions).

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Dosage/Frequency:** PF-08634404 in combination with Sigvotatug Vedotin (Part A) or other specific combination agents (Part B) according to dose escalation rules. * **Route of Administration:** Intravenous (IV) infusion at the study clinic. * **Duration of Treatment:** Cycles continue until disease progression, unacceptable toxicity, or study withdrawal.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Standard supportive care medications. * **Prohibited:** Concurrent systemic anti-tumor therapies or investigational agents outside the study protocol.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Tumor Assessments, Biomarker testing
Baseline	Day 0	Safety Baseline, First IV Dose
Interim	Per Cycle	IV Infusions, Safety Labs, DLT Observation, Efficacy Assessment (RECIST), PK/ADA
End of Study	Follow-up	End of Treatment Visit, Follow-up for Survival and adverse effects

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Adverse Events (AEs) as characterized by type, frequency, severity, timing, and seriousness. The number of participants who experienced Dose Limiting Toxicities (DLTs) during the DLT observation period. * **Measurement Method:** Severity graded by NCI CTCAE version 5.0 and clinical observation.

7.2 Secondary & Exploratory Endpoints * Objective Response Rate (ORR) defined as the proportion of participants with a Best Overall Response (BOR) of confirmed Complete Response (CR) or Partial Response (PR) per RECIST v1.1. * Pharmacokinetics (PK) of

PF-08634404 with anticancer agents. * Immunogenicity of PF-08634404 with anticancer agents.

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Continuous monitoring by trained medical staff during and after each visit, with grading per NCI CTCAE version 5.0.
- **Serious Adverse Events (SAEs):** Standard protocol collection with immediate reporting to the sponsor.
- **Data Monitoring Committee (DMC):** Safety Review Committee manages dose escalation milestones.

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Safety Set (all participants who receive at least one dose) and DLT-evaluable set.
- **Sample Size Justification:** Adaptive Phase 1b/2 dose escalation and expansion statistics governing DLT windows.
- **Handling of Missing Data:** No formal imputation for safety analyses; specific strategies detailed per protocol for efficacy (ORR).

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
- **Monitoring Plan:** Regular onsite and remote monitoring by the sponsor (Pfizer) at the clinical trial sites.
- **Audit Trail:** Robust Electronic Data Capture (EDC) utilizing data clarification procedures.

11. Study Identifiers & Governance

- **Primary ID:** C6461020
- **Registry Number:** NCT07227298
- **Sponsor/Lead Organization:** Pfizer
- **Study Title:** A Study to Learn About the Study Medicine Called PF-08634404 in Combination With Different Anticancer Agents in Advanced Cancers
- **Official Regulatory Title:** AN INTERVENTIONAL OPEN-LABEL PHASE 1B/2 STUDY TO EVALUATE THE SAFETY, PHARMACOKINETICS, AND PRELIMINARY EFFICACY OF PF-08634404 IN COMBINATION WITH DIFFERENT ANTICANCER AGENTS IN PARTICIPANTS WITH ADVANCED SOLID TUMORS

12. Clinical Framework (Oncology Specific)

- **Primary Condition:** Advanced/Metastatic Non-Small Cell Lung Cancer (NSCLC) and Advanced Solid Tumors

- **Diagnosis Threshold:** Pathologically confirmed Stage IIIB/IIIC or Stage IV NSCLC
- **Medical Methodology:** Interventional biologic combinations via Intravenous (IV) Infusion
- **Optimization Target:** Identifying Maximum Tolerated Dose / Recommended Phase 2 Dose

13. Study Procedures & Methodology

- **Management Pathway:** Interventional Oncology Management (Phase 1b/2)
- **Education Protocol (HCP):** Site Initiation Visits, comprehensive DLT reporting training
- **Education Protocol (Patient):** Comprehensive informed consent regarding early phase combinations and toxicity profiles
- **Quality Audit Mechanism:** Source Data Verification by sponsor monitors

14. Observation & Follow-Up Schedule

- **Total Duration:** Projected length from FPI to Study Completion is approx. 8 years (Estimated August 23, 2033)
- **Onsite Visit Cadence:** Based on specific cycle dosing schedules
- **Remote Monitoring Frequency:** Between treatment cycles, as clinically indicated
- **Checklist Review Interval:** Regular tumor assessment scans required per RECIST v1.1 schedules

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					