

1. Administrative & Study Identification

Full Study Title: An Interventional Phase 1b, Randomized, Double-Blind, Sponsor-Open, Placebo-Controlled Study to Investigate the Safety, Tolerability, and Pharmacodynamic Effect of PF-07258669 Compared to Placebo in Older Participants Including Those at Risk of Malnutrition **Short Title/Acronym:** Safety and Effects of PF-07258669 vs Placebo in Older Adults at Risk of Malnutrition **Protocol Number:** C4541011 / 2025-520763-41-00 **NCT Number:** NCT07086664 **Sponsor Name:** Pfizer Inc. **Investigational Product:** PF-07258669 (Oral tablet; MC4R antagonist) **Phase of Development:** Phase 1b **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety, tolerability, and pharmacodynamic effects (how the drug affects the body) of PF-07258669 over 20 weeks in older participants.

Secondary Objectives: To measure changes in body weight from the start of the study to week 16, which serves as an indicator of whether the medication positively affects nutrition or metabolism.

2.2 Background & Rationale To address unintended weight loss and malnutrition risk in older adults. People with this condition often experience decreased appetite and food intake, which leads to poor nutrition and compromised health. PF-07258669 is being developed as an antagonist of the melanocortin-4 receptor (MC4R) to boost appetite and increase food intake. This Phase 1b study aimed to compare the experimental medication against a placebo to determine safety, side effects, and early efficacy signals.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled, Parallel Group **Blinding:** Double-blind (Quadruple masking: Participant, Care Provider, Investigator, Outcomes Assessor) **Randomization:** Randomized

3.2 Planned Enrollment & Duration Target \$N\$: Planned for 60 participants. *(Note: The study was prematurely discontinued due to a business decision after enrolling 7 participants. There were no safety concerns related to the termination).* **Number of Sites:** Multiple global sites **Estimated Study Start:** August 2025 **Estimated Primary Completion:** October 2026 (Discontinued early in 2026)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Age ≥ 60 years to < 90 years. 2. Males or females who can no longer have children. 3. Body weight ≥ 40 kg (about 88 pounds) and Body mass index (BMI) ≤ 25 kg/m².

4.2 Exclusion Criteria 1. Evidence or history of clinically significant medical conditions. 2. Positive for active, untreated HIV, hepatitis B, and/or hepatitis C. 3. History of alcohol abuse or binge drinking and/or any other illicit drug use or dependence within 6 months of Screening.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: PF-07258669 or matching placebo given as oral tablets. **Route of Administration:** Oral **Duration of Treatment:** 16 weeks of treatment.

5.2 Concomitant & Prohibited Therapy Permitted: Standard daily medications suited for an older population without interfering with the trial's drug absorption. **Prohibited:** Illicit drugs, binge drinking, and medications that may significantly alter metabolism or appetite.

6. Study Timeline & Visit Schedule

Visit ID	Window	Key Activities/Procedures
1	Up to 4 weeks	Informed Consent, Medical History, BMI / Weight Check (1 in-person visit)
2	2 weeks	Baseline safety evaluations, pre-treatment preparation (1 in-person visit)
3-17	16 weeks	Randomization, Study Drug/Placebo administration, Safety & Weight Monitoring (8 in-person visits, 4 telehealth visits)
18	4 weeks	Final safety checks, Post-treatment observation (1 in-person visit)

(Total study duration per participant: ~26 weeks)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence of adverse events, side effects, and overall safety and tolerability over 20 weeks of observation. **Measurement Method:** Clinical monitoring, vital signs, and safety laboratory tests.

7.2 Secondary & Exploratory Endpoints Definition: Change in body weight from baseline to Week 16.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Patient monitoring during the 11 scheduled in-person visits and 4 remote telehealth consultations. **Serious Adverse Events (SAEs):** Standard protocol collection and reporting within 24 hours. **Data Monitoring Committee (DMC):** Internal safety monitoring by the Sponsor. *(Note: The study's early termination was a Pfizer business decision, independent of any AE/SAE or safety signals).*

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all randomized participants who received at least one dose of the study drug/placebo). **Sample Size Justification:** A target $N = 60$ was calculated to be sufficient to provide preliminary safety, tolerability, and early pharmacodynamic trends in a Phase 1b setting. **Handling of Missing Data:** Last Observation Carried Forward (LOCF) for early dropouts or standard mixed-model repeated measures (MMRM) for continuous variables like body weight.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study was conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Combination of onsite visits and telehealth assessments. **Audit Trail:** Standard electronic Case Report Form (eCRF) locks and queries per Pfizer clinical operational procedures.

11. Study Identifiers & Governance

Primary ID: C4541011 **Registry Number:** NCT07086664 **Sponsor/Lead Organization:** Pfizer Inc. **Study Title:** Safety and Effects of PF-07258669 vs Placebo in Older Adults at Risk of Malnutrition **Official Regulatory Title:** An Interventional Phase 1b, Randomized, Double-Blind, Sponsor-Open, Placebo-Controlled Study to Investigate the Safety, Tolerability, and Pharmacodynamic Effect of PF-07258669 Compared to Placebo in Older Participants Including Those at Risk of Malnutrition

12. Clinical Framework

Primary Condition: Unintended weight loss / risk of malnutrition **Diagnosis Threshold:** BMI ≤ 25 kg/m², Weight ≥ 40 kg, Ages 60-89 **Medical Methodology:** Pharmacological Intervention (MC4R antagonist) **Optimization Target:** Improvement in appetite, food intake, and body weight

13. Study Procedures & Methodology

Management Pathway: 16-week oral medication period with close clinical monitoring. **Education Protocol (HCP):** Clinical trial site training on dispensing study medicine and placebo. **Education Protocol (Patient):** Nutritional counseling and adherence tracking for oral tablets. **Quality Audit Mechanism:** Clinical site monitoring and safety reviews across the full 26-week timeline.

14. Observation & Follow-Up Schedule

Total Duration: 26 weeks (including screening and follow-up)
Onsite Visit Cadence: 11 total in-person visits (1 screening, 1 pre-treatment, 8 during the 16-week treatment, 1 at follow-up)
Remote Monitoring Frequency: 4 telehealth visits during the treatment phase
Checklist Review Interval: Regular assessments during the 16-week treatment phase.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				