

1. Administrative & Study Identification

Official Study Title: A Phase 2 Study of Brentuximab Vedotin in Combination With Pembrolizumab in Participants With Metastatic Solid Malignancies **Short Title/Acronym:** Brentuximab Vedotin With Pembrolizumab in Metastatic Solid Tumors (SGN35-033) **Protocol Number:** SGN35-033 **NCT Number:** NCT04609566 **Trial Status:** Active, not recruiting **Sponsor Name:** Pfizer **Investigational Product:** Brentuximab Vedotin (Adcetris) + Pembrolizumab (PD-1 Inhibitor) **Phase of Development:** Phase 2 **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the confirmed objective response rate (ORR) of the combination of brentuximab vedotin with pembrolizumab in participants with metastatic solid tumors. **Secondary Objectives:** To evaluate Progression-Free Survival (PFS), Duration of Response (DOR), and Objective Response Rate (ORR) per iRECIST, as well as to assess the overall safety and tolerability of the regimen.

2.2 Background & Rationale Brentuximab vedotin (BV) is a CD30-directed antibody-drug conjugate (ADC). It is hypothesized that BV may selectively deplete immunosuppressive T regulatory cells (Tregs) that express CD30 and modulate the tumor microenvironment through the induction of immunogenic cell death (ICD). This mechanism has the potential to resensitize tumors to anti-PD-1 therapy, enhancing the antitumor immunity of pembrolizumab in patients with PD-1-refractory metastatic solid tumors (such as NSCLC and cutaneous melanoma) or treatment-naïve tumors (such as HNSCC).

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Multi-cohort) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target \$N\$: 161 participants **Number of Sites:** Multicenter (United States) **Estimated Study Start:** 23-Oct-2020 **Current Status:** Active, not recruiting

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Participants must have one of the following: * Metastatic squamous or nonsquamous non-small cell lung cancer (NSCLC) (without known targetable EGFR, ALK, ROS1, or BRAF mutations) who either a) have not yet received frontline therapy for metastatic disease and without prior exposure to anti PD-1/PD-L1, or b) are relapsed/refractory with progression on anti PD-1/PD therapy. * Relapsed/refractory metastatic cutaneous melanoma (regardless of mutation status) with progression on a PD-1 inhibitor. * Metastatic head and neck squamous cell carcinoma (HNSCC) who have not yet received frontline therapy for metastatic disease and without prior exposure to a PD-1/PD-L1 inhibitor. 2. Cohorts 1-4 only: Melanoma participants must be currently on PD-1 checkpoint inhibitor (CPI) therapy (e.g., nivolumab or pembrolizumab) or had their last dose of PD-1 CPI containing therapy as the last previous line of therapy within 90 days prior to enrollment; PD-1 CPI therapy must be the immediate prior line of treatment. 3. Cohorts 1-4 only: Participants must have progressed on treatment with an anti-PD-1 monoclonal antibody (mAb) administered either as monotherapy, or in combination with other CPIs or other therapies. PD-1 treatment progression is defined by meeting all of the following criteria: * Have received at least 2 doses of an approved PD-1 inhibitor. * Have demonstrated disease progression (PD) after a PD-1 inhibitor as defined by RECIST v1.1. * Progressive disease has been documented within 90 days from the last dose of PD-1 inhibitor. * Participants with melanoma will need iRECIST confirmation of progression with a second assessment at least four weeks after the initial date of progressive disease. * NSCLC participants on PD-1 inhibitor containing therapy for less than 90 days will need iRECIST confirmation of progression at least 4 weeks after the initial date of progressive disease. 4. Tumor tissue sample obtained within 3 months prior to enrollment is required, and no systemic anticancer therapy given after the sample was obtained. 5. An Eastern Cooperative Oncology Group (ECOG) Performance Status score of equal to or less than 1.

4.2 Exclusion Criteria 1. Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis. 2. Prior immunosuppressive chemotherapy, any immunotherapy other than a PD-1 inhibitor within 4 weeks of first study drug dose. 3. History of another malignancy within 3 years before the first dose of study drug or any evidence of residual disease from a previously diagnosed malignancy.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Brentuximab vedotin (Adcetris; ATC Code: L01FX05) 1.8 mg/kg every 3 weeks in combination with pembrolizumab 200 mg every 3 weeks. **Route of Administration:** Intravenous (IV). Brentuximab vedotin is

administered at least 30 minutes after the end of the pembrolizumab infusion. **Duration of Treatment:** Up to a maximum of 35 cycles (each cycle is 21 days) or until disease progression/unacceptable toxicity.

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care for managing infusion reactions and routine adverse events. **Prohibited:** Other systemic anticancer therapies, targeted therapies, or investigational agents concurrent with the study regimen.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Tumor Imaging, Biopsy
Baseline	Day 1	Cycle 1 Day 1: First Dose (Pembrolizumab followed by BV)
Interim	Every 3 Weeks	Cycles 2-35: IV Drug Administration, Safety Labs, AE Assessment
End of Study	Follow-up	Final Vital Signs, Exit Interview, Survival Follow-up

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Confirmed objective response rate (ORR) based on investigator assessment using Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST v1.1) criteria. **Measurement Method:** Proportion of participants whose best overall response is a confirmed complete response (CR) or partial response (PR) per RECIST v1.1.

7.2 Secondary Endpoints * **Duration of response (DOR)** based on investigator assessment using RECIST v1.1 criteria. * **Progression-free survival (PFS)** based on investigator assessment using RECIST v1.1 criteria. * **ORR per iRECIST** by investigator assessment.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring for Treatment-Emergent Adverse Events (TEAEs) graded by CTCAE. **Serious Adverse Events (SAEs):** Standard reporting timelines for drug-related SAEs. **Data Monitoring Committee (DMC):** Internal/External safety monitoring mechanisms managed by the sponsor to ensure tolerability.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: All enrolled patients who received at least one dose of study treatment (Safety and Efficacy sets). **Sample Size Justification:** Target enrollment is 161 participants. The multi-cohort design evaluates early response rates to determine potential cohort expansions. **Handling of Missing Data:** Time-to-event endpoints (PFS, DOR) are analyzed utilizing Kaplan-Meier methodologies.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Standard sponsor clinical oversight and onsite/remote monitoring for adherence to the study protocol. **Audit Trail:** eCRF locking, source data verification, and secure data clarification tracking.

11. Study Identifiers & Governance

Sponsor/Company: Pfizer **Study Title:** Brentuximab Vedotin With Pembrolizumab in Metastatic Solid Tumors **Official Regulatory Title:** A Phase 2 Study of Brentuximab Vedotin in Combination With Pembrolizumab in Participants With Metastatic Solid Malignancies **Primary ID:** SGN35-033 **Registry Number:** NCT04609566 **Posting ID:** 1109176357543606839

12. Clinical Framework (Oncology Specific)

Disease Areas / Conditions: Melanoma, Non-small Cell Lung Cancer, Squamous Cell Carcinoma of the Head and Neck. **Diagnosis Threshold:** Histologically confirmed metastatic disease; meeting disease-specific prior treatment criteria (relapsed/refractory or treatment-naïve depending on cohort). **Medical Methodology:** Combination of Antibody-Drug Conjugate (ADC) and PD-1 Checkpoint Inhibitor. **Optimization Target:** Objective Tumor Response, Immunogenic Cell Death, and T regulatory cell (Treg) depletion.

13. Study Procedures & Methodology

Management Pathway: 21-day intravenous treatment cycles. **Education Protocol (HCP):** Site initiation visits and toxicity/AE management training. **Education Protocol (Patient):** Informed consent processes and immunotherapy toxicity awareness. **Quality Audit Mechanism:** Independent imaging review capabilities and robust source data verification.

14. Observation & Follow-Up Schedule

Total Duration: Up to 35 treatment cycles plus survival follow-up. **Onsite Visit Cadence:** Every 3 weeks (Day 1 of each 21-day cycle). **Remote Monitoring Frequency:** As required per patient condition between infusions. **Checklist Review Interval:** Tumor assessments and imaging scans typically occurring every 9 to 12 weeks.

15. Sign-off & Approval

Role	Name	Signature	Date		:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]					
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]					Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]					