

1. Administrative & Study Identification

- **Full Study Title:** A PHASE 1, RANDOMIZED, DOUBLE BLIND, SPONSOR-OPEN, PLACEBOCONTROLLED, SINGLE DOSE ESCALATION STUDY TO EVALUATE THE SAFETY, TOLERABILITY, AND PHARMACOKINETICS OF SUBCUTANEOUS ADMINISTRATION OF PF-07999415 IN HEALTHY ADULT PARTICIPANTS
- **Short Title/Acronym:** A Study to Learn How Different Amounts of the Study Medicine Called PF-07999415 Are Tolerated and Act in the Body in Healthy Adults
- **Protocol Number:** C5671001
- **NCT Number:** NCT06965465
- **Sponsor Name:** Pfizer
- **Investigational Product:** PF-07999415
- **Phase of Development:** Phase I
- **Medical Monitor:** Pfizer Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To evaluate the safety, tolerability, and pharmacokinetics of a single subcutaneous administration of PF-07999415 in healthy adult participants. * **Secondary Objectives:** To further characterize the pharmacokinetic profile and evaluate any initial pharmacodynamic markers associated with the administration of the study medicine.

2.2 Background & Rationale PF-07999415 is a new investigational molecular entity currently in early-stage development (in pipeline for Obesity). This Phase I study is required to establish the basic safety, tolerability, and pharmacokinetic profile of single escalating doses of the medication in healthy adult volunteers. These insights are critically necessary to determine the appropriate and safe dosing parameters for future clinical trials involving patient populations.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Placebo-controlled * **Blinding:** Double-blind, Sponsor-open * **Randomization:** Randomized

3.2 Planned Enrollment & Duration * **Target N:** 53 participants * **Number of Sites:** Global (United States, Belgium) * **Estimated Study Start:** May-2025 * **Estimated Primary Completion:** Late 2025 / Early 2026 (estimated)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male and female participants of non-childbearing potential aged 18 to 65 years, inclusive, at screening who are overtly healthy as determined by medical evaluation. 2. Body Mass Index (BMI) of 18-33 kg/m². 3. Total body weight >50 kg (110 lb).

4.2 Exclusion Criteria 1. Evidence or history of clinically significant hematological, renal, endocrine, pulmonary, dermatological, gastrointestinal, cardiovascular, hepatic, psychiatric, neurological, or allergic disease. 2. Any medical or psychiatric condition including recent (within the past year) or active suicidal ideation/behavior. 3. Previous administration of an investigational product within 30 days or 5 half-lives preceding the first dose of study intervention.

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Dosage/Frequency:** Single Dose Escalation. * **Route of Administration:** Subcutaneous (SC) injection (administered in the thigh, abdomen, or arm). * **Duration of Treatment:** Single injection, followed by an inpatient stay of 2 weeks in a controlled hospital-like environment.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Standard medical care that does not interfere with the PK and safety evaluations. * **Prohibited:** Other investigational products or vaccines at any time during participation in this study.

6. Study Timeline & Visit Schedule

- **Screening (Day -28 to 0):** Informed Consent, Medical History, Physical Examination, Lab tests, Cardiac monitoring.
- **Baseline (Day 0):** Confinement admission, Randomization.
- **Intervention & Interim (Weeks 1 to 2):** Single dose SC administration, followed by 2 weeks of continuous inpatient monitoring for reactions, side effects, and PK sampling.
- **End of Study (Months 3 to 7):** Outpatient follow-up visits every 2 weeks.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Incidence of treatment-emergent adverse events (TEAEs) and serious adverse events (SAEs) to evaluate the safety and tolerability of PF-07999415. * **Measurement Method:** Clinical observation during the 2-week confinement, laboratory tests, cardiac monitoring, and outpatient evaluations.

7.2 Secondary & Exploratory Endpoints * Pharmacokinetics of PF-07999415 (Maximum observed concentration [C_{max}], Area under

the concentration-time curve [AUC]). * Characterization of localized injection site reactions.

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Intensive monitoring by clinical staff during the 2-week inpatient stay, followed by HCP interview during subsequent bi-weekly visits.
- **Serious Adverse Events (SAEs):** Standard protocol, typically 24 hours reporting requirement.
- **Data Monitoring Committee (DMC):** Internal safety review committee (sponsor-open).

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Safety Analysis Set.
- **Sample Size Justification:** $N = 53$ provides sufficient power to identify initial safety signals and define single-dose PK parameters across the escalation cohorts.
- **Handling of Missing Data:** Standard handling for PK analyses (e.g., assigning zeros to concentrations below the limit of quantification).

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
- **Monitoring Plan:** Continuous onsite clinical monitoring during the 2-week inpatient confinement; onsite/remote checks during the follow-up period.
- **Audit Trail:** Data verification, eCRF locking, and rigorous data tracking as per Pfizer's clinical standards.

11. Study Identifiers & Governance

- **Primary ID:** C5671001
- **Registry Number:** NCT06965465
- **Sponsor/Lead Organization:** Pfizer
- **Study Title:** A Study to Learn How Different Amounts of the Study Medicine Called PF-07999415 Are Tolerated and Act in the Body in Healthy Adults
- **Official Regulatory Title:** A PHASE 1, RANDOMIZED, DOUBLE BLIND, SPONSOR-OPEN, PLACEBOCONTROLLED, SINGLE DOSE ESCALATION STUDY TO EVALUATE THE SAFETY, TOLERABILITY, AND PHARMACOKINETICS OF SUBCUTANEOUS ADMINISTRATION OF PF-07999415 IN HEALTHY ADULT PARTICIPANTS

12. Clinical Framework (Study Specific)

- **Primary Condition:** Healthy Volunteers (Development pipeline target: Obesity)

- **Diagnosis Threshold:** Overtly healthy; BMI 18-33 kg/m2.
- **Medical Methodology:** Single Ascending Dose Escalation
- **Optimization Target:** Establishment of safety, tolerability, and PK profile in humans.

13. Study Procedures & Methodology

- **Management Pathway:** Phase 1 confined clinical pharmacology unit pathway.
- **Education Protocol (HCP):** Protocol training on subcutaneous dose administration and rigorous AE tracking.
- **Education Protocol (Patient):** Inpatient facility orientation and post-discharge symptom/AE reporting training.
- **Quality Audit Mechanism:** Standard site monitoring, source data verification, and clinical quality checks.

14. Observation & Follow-Up Schedule

- **Total Duration:** 3 to 7 months per participant
- **Onsite Visit Cadence:** 2-week inpatient confinement, followed by outpatient facility visits every 2 weeks.
- **Remote Monitoring Frequency:** N/A (Follow-ups are conducted in person).
- **Checklist Review Interval:** During every scheduled visit.

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				