

1. Administrative & Study Identification

Full Study Title: A Phase II, Randomized, Open-label, Parallel Group, Multicenter Study to Assess Bioequivalence of Two Subcutaneous Formulations of Ocrelizumab in Patients With Multiple Sclerosis **Short Title/Acronym:** PORTAMENTO **Protocol Number:** CN45320 / 2024-517980-22-00 **NCT Number:** NCT07074886 **Trial Status:** Not Yet Recruiting **Sponsor Name:** Hoffmann-La Roche (Roche) **Investigational Product:** Ocrelizumab (Trade Name: Ocrevus) SC Test Formulation vs. Marketed Ocrelizumab SC Reference Formulation **Phase of Development:** Phase II **Medical Monitor:** Sponsor Medical Monitor, Hoffmann-La Roche

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To assess the bioequivalence of the ocrelizumab subcutaneous (SC) test formulation compared to the marketed ocrelizumab SC reference formulation in participants with relapsing multiple sclerosis (RMS) or primary progressive multiple sclerosis (PPMS). **Secondary Objectives:** To evaluate the safety, tolerability, pharmacodynamics, and immunogenicity (how the immune system reacts) of the two different ocrelizumab SC formulations.

2.2 Background & Rationale There are limitations to currently available treatments for multiple sclerosis (MS), as some treatments can take several hours to be given via intravenous infusion. More options for how treatment is given are needed to improve patient convenience and reduce clinic time. This study tests a new experimental SC formulation of ocrelizumab to determine if the body absorbs and responds to it in the same way as the already approved SC reference formulation.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Interventional **Control Type:** Active Comparator (Marketed Ocrelizumab SC Reference Formulation) **Blinding:** Open-label **Randomization:** Randomized, Parallel Group

3.2 Planned Enrollment & Duration **Target Enrollment (\$N\$):** 182 participants **Number of Sites:** Multicenter (Global: including sites in the US, Europe, Argentina, Mexico, Brazil, and South Africa) **Estimated Study Start:** 13-Nov-2025 **Estimated Primary Completion:** 31-Jan-2027

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Adults aged 18 to 65 years, inclusive. 2. Diagnosis of relapsing multiple sclerosis (RMS) or primary progressive multiple sclerosis (PPMS) according to the revised McDonald 2017 criteria (Thompson et al. 2018) or the most current McDonald criteria at the time of study start. 3. Expanded Disability Status Scale (EDSS) score of 0 to 6.5, inclusive, at screening.

4.2 Exclusion Criteria 1. Participants who have previously received anti-CD20s (including ocrelizumab) less than 2 years before screening. 2. Participants who have previously received anti-CD20s (including ocrelizumab) more than 2 years before screening if one of the following conditions is met: B-cell count is below lower limit of normal (LLN), or the discontinuation of the treatment was due to safety reasons. 3. History of confirmed or suspected progressive multifocal leukoencephalopathy (PML). 4. History of cancer, including hematologic malignancy and solid tumors, within 10 years of screening. 5. Immunocompromised state. 6. Sensitivity or intolerance to any ingredient (including excipients) of ocrelizumab. 7. History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies. 8. Any concomitant disease that may require chronic treatment with systemic corticosteroids or immunosuppressants during the course of the study. 9. Significant, uncontrolled disease, such as cardiovascular, pulmonary, renal, hepatic, endocrine, or gastrointestinal, or any other significant disease that may preclude participation in the study. 10. Lack of peripheral venous access. 11. Previous treatment with cladribine, atacicept, and alemtuzumab. 12. Any previous treatment with bone marrow transplantation and hematopoietic stem cell transplantation. 13. Any previous history of transplantation or anti-rejection therapy. 14. Positive screening tests for active, latent, or inadequately treated hepatitis B virus (HBV).

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: The study consists of 2 phases. In the *controlled phase*, participants receive a single injection (Day 1) of either the SC test formulation or the SC reference formulation. After 6 months, in the *continuation phase*, all participants in both groups receive the SC test formulation every 6 months for 2.5 years. **Route of Administration:** Subcutaneous (SC) injection. **Duration of Treatment:** Total study duration per participant is approximately 3.5 years.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant medications for symptomatic management of MS that do not interfere with immune response. **Prohibited:** Systemic

corticosteroids, immunosuppressants, cladribine, atacicept, and alemtuzumab.

6. Study Timeline & Visit Schedule

| Visit ID | Window (Days) | Key Activities/Procedures | | :---
| :--- | :--- | | **Screening** | Day -28 to 0 | Informed Consent, Medical History, EDSS Scoring, Eligibility Labs | | **Baseline (Controlled Phase)** | Day 1 | Randomization, Single SC Dose of Test or Reference Formulation, PK/PD Sampling | | **Interim (Continuation Phase)** | Month 6 onwards | Administration of SC Test Formulation every 6 months, Safety Labs, Efficacy Assessment | | **End of Study** | Year 3.5 | Final Vital Signs, Exit Interview, Follow-up for Safety and Immunogenicity |

(Note: Participants will have a total of 27 study visits over the 3.5-year period)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints 1. Area Under the Serum Concentration-time Curve (AUC) of Ocrelizumab. 2. Maximum Serum Concentration (C_{max}) of Ocrelizumab. *(Measurement Method: Central Laboratory Analysis evaluating serum concentrations to determine bioequivalence of the ocrelizumab SC test formulation compared to the marketed SC reference formulation).*

7.2 Secondary & Exploratory Endpoints 1. Number of Participants With Adverse Events (AEs) to evaluate the safety and tolerability profile of the test formulation versus reference. 2. Immunogenicity (anti-drug antibodies) and Pharmacodynamics (e.g., B-cell depletion levels).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring across the 27 study visits via physical examinations, lab tests, and participant reporting. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal/External safety monitoring mechanisms managed by Hoffmann-La Roche.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Pharmacokinetic (PK) Analysis Set for primary bioequivalence outcomes; Safety Set for all AE/SAE evaluations. **Sample Size Justification:** \$N=182\$, appropriately powered to establish statistical bioequivalence between the two

subcutaneous formulations. **Handling of Missing Data:** Standard handling for PK studies (e.g., imputation rules for values below the limit of quantification) per study protocol.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Onsite and remote monitoring visits according to a risk-based monitoring plan by the sponsor. **Audit Trail:** Electronic Case Report Form (eCRF) tracking, continuous PK sample handling documentation, and query resolution via standard quality audit mechanisms.

11. Study Identifiers & Governance

Primary ID: CN45320 **Posting ID:** 1137933043976371606 **Registry Number:** NCT07074886 **Sponsor/Lead Organization:** Hoffmann-La Roche (Roche) **Study Title:** PORTAMENTO / A Study to Assess Bioequivalence of Two Subcutaneous (SC) Formulations of Ocrelizumab in Participants With Multiple Sclerosis (MS) **Official Regulatory Title:** A Phase II, Randomized, Open-label, Parallel Group, Multicenter Study to Assess Bioequivalence of Two Subcutaneous Formulations of Ocrelizumab in Patients With Multiple Sclerosis

12. Clinical Framework (Multiple Sclerosis Specific)

Primary Condition: Relapsing Multiple Sclerosis (RMS) or Primary Progressive Multiple Sclerosis (PPMS) **Diagnosis Threshold:**

Confirmed MS diagnosis with EDSS score of 0 to 6.5 **Medical**

Methodology: Anti-CD20 Monoclonal Antibody Therapy **ATC Code:**

L04AA36 **EMA Product Information:** [Ocrevus EPAR](#) **Optimization**

Target: Bioequivalence and improved patient administration via SC formulation

13. Study Procedures & Methodology

Management Pathway: Standard clinical pathway for Subcutaneous (SC) MS Treatments **Education Protocol (HCP):** Subcutaneous injection administration training and handling of test vs.

reference IP **Education Protocol (Patient):** Counseling on

injection site reactions and long-term (3.5 year) study

compliance **Quality Audit Mechanism:** Strict monitoring of PK/PD sampling windows to ensure accurate bioequivalence calculations

14. Observation & Follow-Up Schedule

Total Duration: 3.5 Years **Onsite Visit Cadence:** Total of 27 study visits spanning the 3.5-year duration **Remote Monitoring Frequency:** Intermittent monitoring between the 6-month dosing intervals **Checklist Review Interval:** Dosing occurs every 6 months during the 2.5-year continuation phase

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				