

1. Administrative & Study Identification

Full Study Title A Phase 3, Randomized, Two-part Study with a 5-week Double-blind Part (Randomized, Parallel-group, Placebo-controlled) followed by a 52-week Open-label Extension Part to Evaluate the Efficacy and Safety of KarXT in Acutely Psychotic Japanese Adult Patients with Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (DSM-5) Schizophrenia (J-EMERGENT) **Short Title/Acronym** A Study to Evaluate the Efficacy and Safety of KarXT in Acutely Psychotic Japanese Adult Participants With Schizophrenia (J-EMERGENT) **Protocol Number** CN012-0019 **NCT Number** NCT06882785 **Sponsor Name** Bristol-Myers Squibb **Investigational Product** KarXT (xanomeline and trospium chloride) **Phase of Development** Phase III **Medical Monitor** Clinical Operations Lead / Zhang Changliang

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To evaluate the efficacy and safety of KarXT in acutely psychotic Japanese adult participants with schizophrenia. **Secondary Objectives:** To evaluate the long-term safety and tolerability profile of KarXT during a 52-week open-label extension.

2.2 Background & Rationale To address the critical need for effective and well-tolerated treatments for acute psychotic symptoms in adult patients diagnosed with schizophrenia, evaluating the therapeutic potential and long-term safety of the novel agent KarXT.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind (5-week initial phase) and Open-label (52-week extension phase) **Randomization:** Randomized, Parallel-group

3.2 Planned Enrollment & Duration **Target** \$N\$: 250 participants **Number of Sites:** 55 locations in Japan **Estimated Study Start:** 10-Jun-2025 **Estimated Primary Completion:** 26-Mar-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Participants must have a primary diagnosis of schizophrenia established by a comprehensive

psychiatric evaluation based on the Diagnostic and Statistical Manual of Mental Disorders–Fifth Edition (DSM-5) criteria and confirmed by Mini International Neuropsychiatric Interview (MINI). 2. Participants must have a PANSS total score between 80 and 120, inclusive. 3. Participants must have a CGI-S score of ≥ 4 .

4.2 Exclusion Criteria 1. Participants must not have any primary DSM-5 disorder other than schizophrenia within 12 months before screening. 2. Participants must not be newly diagnosed or experiencing their first treated episode of schizophrenia. 3. Participants must not have any history or presence of clinically significant medical conditions.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: KarXT capsules or matching placebo. **Route of Administration:** Oral. **Duration of Treatment:** 5-week double-blind part, followed by a 52-week open-label extension part (total 57 weeks).

5.2 Concomitant & Prohibited Therapy Permitted: Standard allowed concomitant medications per protocol specifications. **Prohibited:** Specific prohibited concurrent psychotropic medications and therapies during the evaluation periods.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -14 to 0	Informed Consent, Psychiatric Evaluation, PANSS/CGI-S Baseline, Safety Labs
Baseline	Day 0	Randomization, First Dose
Interim 1	Week 5	Primary Efficacy Assessment (PANSS), Safety Labs, Transition to Open-Label
End of Study	Week 57	Final Efficacy & Safety Evaluation, Exit Interview, IP Accountability

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Change From Baseline in Positive and Negative Syndrome Scale (PANSS) Total Score at Week 5 (for the Double-blind part), and Incidence of Treatment-Emergent Adverse Events (TEAEs) (for the Open-label extension part). **Measurement Method:** Clinical psychiatric assessment scale rating (PANSS) and rigorous adverse event monitoring.

7.2 Secondary & Exploratory Endpoints * Change from baseline in secondary clinical efficacy scales (e.g., CGI-S). * Ongoing monitoring of vital signs, laboratory parameters, and suicidality.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection via regular clinical site visits, psychiatric evaluations, and patient interviews focusing on TEAEs. **Serious Adverse Events (SAEs):** Investigational site reporting to the sponsor within 24 hours. **Data Monitoring Committee (DMC):** External independent safety committee managed by the sponsor to oversee trial data integrity and safety.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) for primary efficacy and Safety Set for the evaluation of adverse events. **Sample Size Justification:** $N = 250$ appropriately powered ($\alpha = 0.05$, $Power = 80\%$) to detect statistically significant differences in PANSS score changes between KarXT and placebo. **Handling of Missing Data:** Standard methodologies (e.g., MMRM - Mixed-Model Repeated Measures) to handle missing values for psychiatric scale endpoints.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring visits by clinical research associates to ensure protocol adherence and patient safety. **Audit Trail:** eCRF locking, tracking of scale administration consistency, and data clarification forms via centralized rating platforms.

11. Study Identifiers & Governance

Primary ID: CN012-0019 **Registry Number:** NCT06882785 **Sponsor/Lead Organization:** Bristol-Myers Squibb **Study Title:** A Study to Evaluate the Efficacy and Safety of KarXT in Acutely Psychotic Japanese Adult Participants With Schizophrenia **Official Regulatory Title:** A Phase 3, Randomized, Two-part Study with a 5-week Double-blind Part (Randomized, Parallel-group, Placebo-controlled) followed by a 52-week Open-label Extension Part to Evaluate the Efficacy and Safety of KarXT in Acutely Psychotic Japanese Adult Patients with Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (DSM-5) Schizophrenia (J-EMERGENT)

12. Clinical Framework (Study Specific)

Primary Condition: Schizophrenia **Diagnosis Threshold:** DSM-5 diagnosis, PANSS total score ≥ 80 and ≤ 120 , CGI-S score

Medical Methodology: Pharmacological Treatment (KarXT vs. Placebo) **Optimization Target:** Significant reduction in psychotic symptoms per PANSS

13. Study Procedures & Methodology

Management Pathway: Acute psychiatric assessment and clinical management
Education Protocol (HCP): Protocol-specific clinical scale training and certification (PANSS/MINI administration)
Education Protocol (Patient): Informed consent process, expectation management, and reporting adverse events
Quality Audit Mechanism: Centralized rater review and adherence monitoring

14. Observation & Follow-Up Schedule

Total Duration: 57 weeks **Onsite Visit Cadence:** Regular scheduled visits throughout the 5-week double-blind period and periodically during the 52-week open-label extension **Remote Monitoring Frequency:** Between physical visits as specified by protocol **Checklist Review Interval:** Continuous tracking of medication adherence and tolerability

15. Sign-off & Approval

Role	Name	Signature	Date	_____	_____	_____	_____
Principal Investigator	_____	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Clinical Operations Lead	Zhang Changliang	_____	[DD-MMM-YYYY]	_____	_____	_____	_____
Lead Statistician	_____	_____	[DD-MMM-YYYY]	_____	_____	_____	_____