

1. Administrative & Study Identification

Full Study Title: AZD6244 (ARRY-142886) Solid Oral Dosage Formulation in Participants With Advanced Solid Malignancies
Short Title/Acronym: AZD6244 Solid Malignancies Phase 1
Protocol Number: D1532C000005
NCT Number: NCT00463814
Sponsor Name: AstraZeneca
Investigational Product: AZD6244 (selumetinib, ARRY-142886)
Phase of Development: Phase 1
Medical Monitor: AstraZeneca Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective: To assess the safety, tolerability, and pharmacokinetics of a capsule of AZD6244 in participants with advanced solid malignancies.
Secondary Objectives: To evaluate preliminary anti-tumor activity, objective response, and the relative bioavailability of the capsule versus a free-base suspension.

2.2 Background & Rationale

To evaluate the clinical pharmacokinetics and safety profile of an oral dosage formulation of AZD6244, a selective, non-ATP-competitive inhibitor of MEK1/2. This targeted therapy aims to arrest the mitogen-activated protein kinase (MAPK) signaling cascade in patients with advanced solid cancer that is refractory to standard therapies, or for whom no conventional therapies exist.

3. Study Design & Methodology

3.1 Design Framework

Study Type: Interventional
Control Type: Single-arm (with varying dose sequences for relative bioavailability)
Blinding: Open-label
Randomization: N/A (Sequential assignment for bioavailability sequences)

3.2 Planned Enrollment & Duration

Target \$\$\$: 59 participants
Number of Sites: 4 research sites (Aurora, CO, US; Nijmegen, Netherlands; Utrecht, Netherlands; Sutton, UK)

Estimated Study Start: 08-Mar-2007
Estimated Primary Completion: 17-Jun-2008

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Cancer which is refractory to standard therapies.
2. World Health Organization (WHO) performance status 0 to 2.
3. Evidence of post-menopausal status or negative pregnancy test.

4.2 Exclusion Criteria

1. Radiotherapy/chemotherapy within 21 days prior to entry.
 2. Brain metastases/spinal cord compression unless stable off steroids/anticonvulsants.
 3. Evidence of severe/uncontrolled systemic disease, or participation in an investigational drug study within 30 days prior.
-

5. Intervention & Treatment Plan

5.1 Investigational Regimen

Dosage/Frequency: Single oral dose on Day 1 (cohorts included 25 mg, 50 mg, 75 mg, and 100 mg capsules, or 100 mg mix-and-drink free-base suspension). This was followed by continuous twice daily (bd) dosing from Day 2 (or Day 9 for crossover cohorts) onwards.

Route of Administration: Oral

Duration of Treatment: Until disease progression or another protocol-defined discontinuation criterion is met.

5.2 Concomitant & Prohibited Therapy

Permitted: Standard supportive palliative care medications for symptom control.

Prohibited: Radiotherapy or chemotherapy within 21 days prior to study entry; other investigational drugs within 30 days.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -21 to 0 Informed Consent, Medical History, Performance Status evaluation Baseline Day 1 First oral dose of AZD6244, Pharmacokinetic (PK) blood sampling Interim Day 2 to Day 8 Continuous twice-daily dosing (or 7-day washout for relative bioavailability sequence crossovers), Safety Labs End of Study Up to 11.8 months Disease progression assessment, final TEAE / TSEAE documentation, IP Accountability

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint

Definition: Number of Participants With Treatment-Emergent Adverse Events (TEAEs), Treatment-Emergent Serious Adverse Events (TESAEs), and Abnormal Clinical Laboratory Parameters in Part A and Part B measured up to a maximum observed duration of 11.8 months.

Measurement Method: Clinical safety observations, vital signs, and laboratory blood analysis.

7.2 Secondary & Exploratory Endpoints

- Pharmacokinetics of AZD6244 (e.g., C_{max} , AUC , apparent oral clearance).
 - Anti-tumor efficacy and objective tumor response (e.g., evaluated via CT/PET scans).
-

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection with close monitoring for known MEK1/2 inhibitor toxicities, specifically dermatologic toxicity (rash), gastrointestinal toxicities, and fatigue.

Serious Adverse Events (SAEs): Reported for TEAEs and TESAEs throughout the treatment duration up to 11.8 months.

Data Monitoring Committee (DMC): Internal safety review committee managed by AstraZeneca.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants who received at least one dose of the investigational drug).

Sample Size Justification: Sized according to standard Phase 1 dose-escalation and safety assessment principles ($N=59$) to effectively map the PK profile and identify the maximum tolerated dose.

Handling of Missing Data: Last Observation Carried Forward (LOCF) for ongoing AEs and PK parameters where applicable.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Regular on-site and remote monitoring across the 4 global research sites.

Audit Trail: eCRF locking, tracking of treatment deviations, and data clarification forms.

11. Study Identifiers & Governance

Primary ID: D1532C000005

Registry Number: NCT00463814

Sponsor/Lead Organization: AstraZeneca

Study Title: AZD6244 (ARRY-142886) Solid Oral Dosage Formulation in Participants With Advanced Solid Malignancies

Official Regulatory Title: AZD6244 (ARRY-142886) Solid Oral Dosage Formulation in Participants With Advanced Solid Malignancies

12. Clinical Framework (Solid Malignancy Specific)

Primary Condition: Advanced Solid Malignancies

Diagnosis Threshold: Cancer that is refractory to standard therapies

Medical Methodology: Targeted MEK1/2 Inhibitor Therapy

Optimization Target: Safety, Tolerability, and Pharmacokinetic profiling to determine recommended dosing

13. Study Procedures & Methodology

Management Pathway: Phase 1 dose-escalation and relative bioavailability crossover sequences (capsule vs. suspension).

Education Protocol (HCP): AE management (specifically identifying and handling dermatologic/rash toxicities).

Education Protocol (Patient): Onsite instructions and diaries for taking oral capsules versus the mix-and-drink suspension.

Quality Audit Mechanism: Strict PK blood sampling integrity tracking and regular tumor response monitoring.

14. Observation & Follow-Up Schedule

Total Duration: Up to 11.8 months (maximum observed duration)

Onsite Visit Cadence: Baseline Day 1, Day 8 (for sequence crossovers), and continuous cycle evaluations.

Remote Monitoring Frequency: Regular safety follow-ups per protocol guidelines.

Checklist Review Interval: Continuous AE tracking and progression-free survival evaluation.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				