

1. Administrative & Study Identification

Full Study Title: A Phase 3, Randomized, Open-label, Multicenter Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan Maintenance Treatment With or Without Bevacizumab Versus Standard of Care After Second-line Platinum-based Doublet Chemotherapy in Participants With Platinum-sensitive Recurrent Ovarian Cancer (MK-2870-022/TroFuse-022/ENGOT-ov84/GOG-3103)
Short Title/Acronym: TroFuse-022 / GOG-3103 / MK-2870-022
Protocol Number: MK-2870-022 **NCT Number:** NCT06824467 **Sponsor Name:** Merck **Company:** Merck **Investigational Product:** Sacituzumab tirumotecan (MK-2870), Bevacizumab **Phase of Development:** Phase 3
Trial Status: Recruiting **Medical Monitor:** Merck Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: * **Part 1:** To evaluate the safety and tolerability of sacituzumab tirumotecan maintenance therapy combined with bevacizumab (assessing adverse events and discontinuations up to 6 weeks). * **Part 2:** To compare the progression-free survival (PFS) of sacituzumab tirumotecan maintenance therapy with or without bevacizumab versus standard of care.

Secondary Objectives: To evaluate Overall Survival (OS), safety profile over the trial duration, and change from baseline in the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30) Global Health Status-Quality of Life Score.

2.2 Background & Rationale To test the safety, efficacy, and effect on the quality of life of sacituzumab tirumotecan when given alone, with bevacizumab, or compared to standard maintenance treatment. The medical necessity is to determine if patients with platinum-sensitive recurrent ovarian cancer who take sacituzumab tirumotecan (with or without bevacizumab) experience extended time living without cancer progression compared to those receiving the standard of care.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Active Comparator (Standard of care with or without bevacizumab)
Blinding: Open-label **Randomization:** Randomized

3.2 Planned Enrollment & Duration Target **\$N\$:** 770 participants (Estimated) **Number of Sites:** Approximately 152 locations globally
Estimated Study Start: 09-Apr-2025 **Estimated Primary Completion:** 27-Apr-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Female, Age ≥ 18 years. 2. Histologically confirmed Federation of Gynecology and Obstetrics (FIGO) Stage III or IV epithelial ovarian, primary peritoneal, or fallopian tube carcinoma of certain histologies. 3. Has received 4 or more cycles of platinum-based doublet chemotherapy in first-line and a total of 6 cycles of carboplatin-based doublet chemotherapy in second-line setting for ovarian cancer (OC); and has platinum-sensitive epithelial OC.

4.2 Exclusion Criteria 1. Has nonepithelial cancers (germ cell tumors and sex cord-stromal tumors), low-grade serous tumors, low-grade endometrioid tumors, borderline tumors (low malignant potential), mucinous, seromucinous, malignant Brenner's tumor and undifferentiated carcinoma. 2. Has platinum-resistant OC or platinum-refractory OC. 3. Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Sacituzumab tirumotecan 4 mg/kg Q2W. At the physician's discretion, bevacizumab 15 mg/kg Q3W. **Route of Administration:** Intravenous (IV) infusion. **Duration of Treatment:** Until progressive disease, unacceptable toxicity, or study discontinuation.

5.2 Concomitant & Prohibited Therapy Permitted: Pre-medications including H1/H2 receptor antagonists, paracetamol (acetaminophen, ATC Code: N02BE01), and dexamethasone/steroid mouthwash. Local standard of care (including bevacizumab alone) is permitted for the active comparator arm. **Prohibited:** Previous treatment with ≥ 2 prior lines of systemic therapy for OC, or any active immunosuppressive medication for inflammatory bowel disease.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, ECOG assessment (within 7 days), Hepatitis B/C and HIV viral load checks, Tumor tissue sample collection.
Baseline	Day 0	Randomization, First Dose of Sacituzumab Tirumotecan $\pm$ Bevacizumab or Standard of Care.
Interim	Q2W / Q3W	IV Infusions, EORTC QLQ-C30 assessments, Safety Labs,

Efficacy Assessment (RECIST 1.1). | | **End of Study** | Up to 4 Years | Final Vital Signs, Exit Interview, RECIST 1.1 Confirmation of Disease Progression or study conclusion. |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: * Part 1: Number of participants with one or more adverse events (AEs) and discontinuations due to AEs up to 6 weeks. * Part 2: Progression-free Survival (PFS) – Time from randomization to the first recorded disease progression or death from any cause up to approximately 4 years. **Measurement Method:** Assessed by a blinded independent central review (BICR) according to RECIST 1.1 criteria.

7.2 Secondary & Exploratory Endpoints * Overall Survival (OS). * Change from Baseline in EORTC QLQ-C30 Global Health Status-Quality of Life Score.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Method of collection via routine clinical and laboratory assessments prior to each infusion cycle. Specifically focusing on ocular/corneal adverse events and pneumonitis/interstitial lung disease. **Serious Adverse Events (SAEs):** Reported continuously throughout the study timeline, typically within 24 hours of site awareness. **Data Monitoring Committee (DMC):** An independent Data Monitoring Committee is involved to assess safety data (especially for Part 1) and efficacy bounds.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) population for primary efficacy endpoints (PFS, OS). Safety Set for AE/SAE evaluations. **Sample Size Justification:** $N=770$ appropriately powered ($\alpha = 0.05$, $\text{Power} \geq 80\%$) to evaluate the superiority of the investigational regimens in prolonging PFS and OS compared to standard of care. **Handling of Missing Data:** Standard oncology guidelines; subjects without documented progression or death will be right-censored at the date of their last evaluable tumor assessment.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.

Monitoring Plan: Regular onsite and centralized data monitoring to ensure fidelity to BICR and RECIST 1.1 criteria. **Audit Trail:**

eCRF locking, tracking of infusion administrations, and data clarification forms via standard quality audit mechanisms.

11. Study Identifiers & Governance

Primary ID: {{MK-2870-022}} **Posting ID:** {{139068158510465436}}
Registry Number: {{NCT06824467}} **Sponsor/Lead Organization:** {{Merck}}
Study Title: {{A Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan (MK-2870) Treatment Versus Standard of Care in Participants With Platinum-sensitive Recurrent Ovarian Cancer (MK-2870-022/TroFuse-022/ENGOT-ov84/GOG-3103)}} **Official Regulatory Title:** {{A Phase 3, Randomized, Open-label, Multicenter Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan Maintenance Treatment With or Without Bevacizumab Versus Standard of Care After Second-line Platinum-based Doublet Chemotherapy in Participants With Platinum-sensitive Recurrent Ovarian Cancer}}

12. Clinical Framework (Oncology Specific)

Primary Condition: {{Ovarian Cancer, Fallopian Tube Cancer, Primary Peritoneal Cancer}} **Diagnosis Threshold:** {{Platinum-sensitive recurrent ovarian cancer}} **Medical Methodology:** {{Maintenance Therapy Protocol}} **Optimization Target:** {{Progression-Free Survival (PFS) Extension}}

13. Study Procedures & Methodology

Management Pathway: {{Sacituzumab Tirumotecan \$\\pm\$ Bevacizumab vs. Standard of Care}} **Education Protocol (HCP):** {{Investigator training on adverse event monitoring, notably corneal toxicity and ILD}} **Education Protocol (Patient):** {{Onsite training/consent detailing Q2W/Q3W infusion schedules}} **Quality Audit Mechanism:** {{Blinded Independent Central Review (BICR) tracking}}

14. Observation & Follow-Up Schedule

Total Duration: {{Up to approximately 4 years}} **Onsite Visit Cadence:** {{Q2W (Sacituzumab Tirumotecan) and Q3W (Bevacizumab)}} **Remote Monitoring Frequency:** {{Continuous via central ePRO collection for EORTC QLQ-C30}} **Checklist Review Interval:** {{Prior to each infusion cycle}}

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	____	[DD-MMM-YYYY]				

Clinical Operations Lead | [Name] | ____ | [DD-MMM-YYYY] | | Lead
Statistician | [Name] | _____ | [DD-MMM-YYYY] |