

1. Administrative & Study Identification

Full Study Title: A Study to Learn How Different Forms of the Study Medicine Called PF 08049820 Are Taken up Into the Bloodstream in Healthy Adults **Short Title/Acronym:** PF-08049820 PK and Food Effect Study **Protocol Number:** C6231007 **NCT Number:** NCT07172321 **Sponsor Name:** Pfizer **Investigational Product:** PF-08049820 (STAT6 transcription factor inhibitor) **Phase of Development:** Phase 1 **Medical Monitor:** Patrizia Cavazzoni, MD (Chief Medical Officer, Pfizer) / Pfizer Clinical Operations

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate how different oral formulations (immediate-release, short-release, and long-release tablets) of PF-08049820 are absorbed into the bloodstream when taken by healthy adults. **Secondary Objectives:** To evaluate if eating a high-fat meal changes the absorption and movement of PF-08049820 through the body, and to evaluate overall safety and tolerability.

2.2 Background & Rationale PF-08049820 is a novel STAT6 transcription factor inhibitor being developed as an oral treatment for moderate to severe atopic dermatitis (eczema). Before advancing to larger efficacy trials, it is necessary to determine the optimal formulation of the drug (fast vs. slow release) and to understand whether food impacts its bioavailability. The results of this Phase 1 study will help decide the best version of the medicine and the correct dosing conditions for future studies.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional (Pharmacokinetic/Bioavailability) **Control Type:** Active Comparator / Self-controlled (crossover) **Blinding:** Open-label **Randomization:** Sequence randomization for formulation crossover

3.2 Planned Enrollment & Duration Target \$N\$: 12 Number of Sites: Pfizer Clinical Research Unit - New Haven, CT, USA **Estimated Study Start:** 05-Sep-2025 **Estimated Primary Completion:** 29-Oct-2025

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male and female participants of non-childbearing potential ≥ 18 years of age. 2. BMI of 16-32 kg/m^2 ; and a total body weight >45 kg (99 lbs). 3. Generally healthy with no serious medical problems, willing and able to comply with all scheduled visits (including an overnight clinic stay of ~ 17 days), treatment plan, and laboratory tests.

4.2 Exclusion Criteria 1. Evidence or history of clinically significant hematological, renal, endocrine, pulmonary, gastrointestinal, cardiovascular, hepatic, psychiatric, neurological, or allergic disease. 2. History of alcohol abuse, repeated binge drinking, or illicit drug use within 6 months of screening; positive testing for HIV, Hepatitis B, or Hepatitis C. 3. Use of certain prescription or nonprescription drugs, dietary supplements (e.g., grapefruit), and herbal supplements within 14 days or 5 half-lives prior to the first dose. 4. Recent exposure to live/attenuated vaccines within 28 days of screening or previous administration of an investigational product within 30 days.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Each participant will receive 5 different single-dose versions of the medicine, tested one at a time, with a 3-day washout period between each dose to ensure the drug clears the system. Forms tested include: immediate-release tablet (fast), short-release tablet (moderate), and long-release tablet (slow). **Route of Administration:** Oral (administered under fasted conditions or after eating a high-fat meal). **Duration of Treatment:** 5 isolated doses administered during an approximate 17-day inpatient clinic stay.

5.2 Concomitant & Prohibited Therapy Permitted: None specified (strictly controlled healthy volunteer cohort). **Prohibited:** Use of prescription/nonprescription drugs, herbal supplements, and dietary items like grapefruit for 14 days prior to dosing.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Days -28 to -1	Informed Consent, Medical History, Vitals, BMI calculation, Lab screening.
Baseline	Day 1	Admission to clinical research unit, baseline vitals, first dose.
Interim	Days 1 to 17	Sequential dosing of the 5 formulations (with 3-day washouts); intensive PK blood sampling up to 72 hours post-dose; continuous safety and AE monitoring.
End of Study	Weeks 6 to 11	Follow-up telephone call approximately one month after the final dosing period to monitor safety.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Pharmacokinetic (PK) profile of PF-08049820, tracking the rate and extent of absorption (including Maximum Observed Concentration [$C_{\max}$] and Area Under the Curve [AUC]) for immediate, short, and long-release formulations. **Measurement Method:** Serial blood sampling collected for up to 72 hours after each dose, analyzed via central laboratory.

7.2 Secondary & Exploratory Endpoints Effect of a high-fat meal on the absorption and PK parameters of PF-08049820. Incidence, severity, and characterization of treatment-emergent adverse events (TEAEs) to assess safety and tolerability.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous clinical observation and physiological assessments by medical staff during the 17-day inpatient stay, followed by a safety phone call one month post-discharge. **Serious Adverse Events (SAEs):** Reported to the sponsor within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal safety review by Pfizer Clinical Operations.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Pharmacokinetic (PK) Concentration Population (all subjects providing evaluable blood samples) and Safety Set (all subjects receiving at least one dose). **Sample Size Justification:** Appropriately powered using a crossover design to estimate pharmacokinetic parameters and food effect ratios within standard Phase 1 confidence intervals. **Handling of Missing Data:** Concentrations below the lower limit of quantification (BLQ) will be treated as zero for summary statistics.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Continuous onsite medical monitoring within the dedicated Pfizer Clinical Research Unit. **Audit Trail:** Strict custody tracking of biological PK samples and protocol deviations; eCRF locking upon completion.

11. Study Identifiers & Governance

Primary ID: C6231007 **Registry Number:** NCT07172321 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn How Different Forms of the Study Medicine Called PF 08049820 Are Taken up Into the Bloodstream in Healthy Adults **Official Regulatory Title:** A Phase 1, Open-Label, Crossover Study to Evaluate the Pharmacokinetics of Different Formulations of PF-08049820 in Healthy Adults

12. Clinical Framework (Phase 1 Specific)

Primary Condition: Healthy Adults (Investigational product targeting Atopic Dermatitis) **Diagnosis Threshold:** BMI 16-32 \$kg/m^2\$ and total body weight \$>45\$ kg **Medical Methodology:** Pharmacokinetic and Bioavailability Assessment **Optimization Target:** Selection of optimal oral formulation and dosing conditions (fasted vs. fed)

13. Study Procedures & Methodology

Management Pathway: Phase 1 Inpatient Clinical Research Unit protocol **Education Protocol (HCP):** Protocol training for clinical staff on exact dosing timings and PK sampling schedules. **Education Protocol (Patient):** Review of protocol rules, dietary restrictions, and inpatient clinic rules. **Quality Audit Mechanism:** Direct observation of dosing and biological sampling in the clinical research unit.

14. Observation & Follow-Up Schedule

Total Duration: 6 to 11 weeks per participant **Onsite Visit Cadence:** Approximately 17 days of continuous overnight inpatient stay **Remote Monitoring Frequency:** Telephone follow-up roughly 1 month after the final dose **Checklist Review Interval:** Daily assessment and monitoring during the clinic stay

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	Pfizer Investigator	____	03-Apr-2026	Clinical Operations Lead	Patrizia Cavazzoni (or proxy)	____	03-Apr-2026
Lead Statistician	Pfizer Statistician	____	03-Apr-2026				