

1. Administrative & Study Identification

Full Study Title: A Study to Evaluate Efficacy and Safety of Deucravacitinib in Participants With Active Discoid and/or Subacute Cutaneous Lupus Erythematosus (DLE/SCLE) **Official Title:** A Phase 2 Randomized, Double-Blind, Placebo-Controlled Study to Evaluate Efficacy and Safety of Deucravacitinib (BMS-986165) in Participants With Active Discoid and/or Subacute Cutaneous Lupus Erythematosus (DLE/SCLE) **Short Title/Acronym:** Deucravacitinib in Active DLE/SCLE **Posting ID:** 215931690249756012 **Protocol Number:** IM011-132 **NCT Number:** NCT04857034 **Sponsor/Company Name:** Bristol Myers Squibb **Investigational Product:** Deucravacitinib (BMS-986165) [Trade Name: Sotyktu, ATC Code: L04AA56] **Phase of Development:** Phase 2 **Trial Status:** Active, Not Recruiting (ACTIVE_NOT_RECRUITING) **Medical Monitor:** Bristol Myers Squibb Clinical Operations Lead / Medical Monitor

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To assess the safety, efficacy, and tolerability of deucravacitinib (BMS-986165) compared with placebo in participants with active discoid and/or subacute cutaneous lupus erythematosus (DLE/SCLE) by evaluating the percentage change from baseline in CLASI Activity Score at Week 16. **Secondary Objectives:** To evaluate the percentage of participants with an improvement of $\geq 50\%$ from baseline in the CLASI-A Score (CLASI-50), the percentage of participants who have disease improvement as defined by a reduction in CLASI-A of ≥ 4 points from baseline, and the mean change from baseline in CLASI-A score. This study will also assess if deucravacitinib is biologically active and potentially effective in the treatment of participants with moderate to severe DLE/SCLE with or without systemic lupus erythematosus (SLE) that is not well controlled with standard of care therapy.

2.2 Background & Rationale To address the gap in the treatment of cutaneous lupus erythematosus (CLE), specifically DLE and SCLE, for participants whose disease is not well controlled with standard of care therapy. Mechanism-based treatments targeting JAK-STAT pathways, such as the selective TYK2 inhibitor deucravacitinib, have the potential to provide positive clinical outcomes by minimizing the use of glucocorticoids and/or non-specific immunosuppressants in systemic immune-mediated inflammatory diseases.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** Randomized

3.2 Planned Enrollment & Duration Target **\$N\$:** 74 participants
Number of Sites: Multicenter **Estimated Study Start:** 12-Jul-2021
Estimated Primary Completion: 22-Aug-2024 (Actual)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Diagnosis of discoid/subacute cutaneous lupus erythematosus (DLE/SCLE) for at least 3 months prior to the screening visit. 2. Meets both clinical and histopathological diagnostic cutaneous lupus erythematosus (CLE) criteria per protocol. 3. Currently receiving treatment for DLE/SCLE with a stable regimen of at least one of the following medications: oral corticosteroid, and/or antimalarial, and/or immunosuppressant. 4. Participant could be with or without concurrent systemic lupus erythematosus (SLE). 5. If participant receives nonsteroidal anti-inflammatory drugs (NSAIDs) or analgesics treatment then the participant must be on a stable dose 2 weeks prior to screening.

4.2 Exclusion Criteria 1. Women who are pregnant, lactating, breastfeeding, or planning pregnancy during the study period. 2. Any of the following specific CLE subtypes in isolation: acute cutaneous lupus erythematosus (ACLE), lupus tumidus, lupus (profundus) panniculitis, chilblains. 3. Drug-induced CLE and/or drug-induced systemic lupus erythematosus (SLE). 4. Antiphospholipid antibody syndrome, serious thrombotic event, or unexplained pregnancy loss within 1 year before the screening visit; or a history of 3 or more unexplained consecutive pregnancy losses. 5. Active severe or unstable neuropsychiatric SLE. 6. Other autoimmune diseases or non-SLE driven inflammatory joint or skin disease or overlap syndromes as primary disease that in the opinion of the investigator will significantly impact the assessment of CLE/SLE disease manifestations and activity.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Deucravacitinib (BMS-986165) [Sotyktu] or matched Placebo. **Route of Administration:** Oral **Duration of Treatment:** Continuous study treatment until disease progression, unacceptable toxicity, or study completion (estimated up to Feb 2028).

5.2 Concomitant & Prohibited Therapy Permitted: Stable regimens of background DLE/SCLE/SLE therapies including oral corticosteroids, antimalarials, immunosuppressants, and NSAIDs/analgesics (if stable 2 weeks prior to screening). **Prohibited:**

Live vaccines within 60 days prior to the first dose. Non-SLE driven inflammatory joint or skin disease therapies that could significantly impact the assessment of CLE/SLE disease manifestations.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Day -14 to 0 Informed Consent, Medical/CLE/SLE History, FSH testing (if applicable) Baseline
Day 0	Randomization, First Dose, Baseline CLASI evaluation	Interim	Periodic Monitor for AEs/SAEs, Safety Labs, Efficacy Assessment (CLASI-A Score at Week 16) End of Study
Safety Follow-up	Final Vital Signs, Exit Interview, IP Accountability, Final Safety Lab Assessments		

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Percentage Change From Baseline in CLASI Activity Score at Week 16. **Measurement Method:** Evaluated using the Cutaneous Lupus Erythematosus Disease Area and Severity Index (CLASI), a validated clinical tool designed to assess skin involvement in CLE by separately scoring disease activity (e.g., erythema, scale, mucous membrane involvement, alopecia) and damage (e.g., dyspigmentation, scarring). CLASI enables classification of disease severity as Mild (Activity score 0-9), Moderate (10-20), or Severe (21-70).

7.2 Secondary & Exploratory Endpoints 1. Percentage of Participants With an Improvement of $\geq 50\%$ From Baseline in the CLASI-A Score (CLASI-50). 2. Percentage of Participants Who Have Disease Improvement as Defined by a Reduction in CLASI-A of ≥ 4 Points From Baseline. 3. Mean Change From Baseline in CLASI-A Score. 4. Ongoing safety, tolerability, and laboratory evaluations.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard collection of non-serious AEs and Serious Adverse Events (SAEs) beginning after the informed consent is completed through the safety follow-up visit. **Serious Adverse Events (SAEs):** Continuously monitored and evaluated per regulatory reporting guidelines. **Data Monitoring Committee (DMC):** Internal/External safety monitoring mechanisms per Bristol Myers Squibb guidelines.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Intent-to-Treat (ITT) population for primary efficacy outcomes; Safety Set for all AE/SAE evaluations. **Sample Size Justification:** Target $N=74$ participants, adequately powered ($\alpha = 0.05$, $\text{Power} = 80\%$) to detect a meaningful clinical improvement between the investigational and placebo arms. **Handling of Missing Data:** Multiple Imputation (MI) and Last Observation Carried Forward (LOCF) for clinical scoring and continuous endpoints.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Routine onsite and remote monitoring to ensure protocol adherence and safety. **Audit Trail:** Data collection via eCRF locking and data clarification forms using standard quality audit mechanisms.

11. Study Identifiers & Governance

Primary ID: IM011-132 **Registry Number:** NCT04857034 **Posting ID:** 215931690249756012 **Sponsor/Lead Organization:** Bristol Myers Squibb **Study Title:** A Study to Evaluate Efficacy and Safety of Deucravacitinib in Participants With Active Discoid and/or Subacute Cutaneous Lupus Erythematosus (DLE/SCLE) **Official Regulatory Title:** A Phase 2 Randomized, Double-Blind, Placebo-Controlled Study to Evaluate Efficacy and Safety of Deucravacitinib (BMS-986165) in Participants With Active Discoid and/or Subacute Cutaneous Lupus Erythematosus (DLE/SCLE)

12. Clinical Framework (Study Specific)

Primary Conditions/Disease Areas: Lupus Erythematosus, Discoid; Lupus Erythematosus, Subacute Cutaneous **Preferred UMLS Name:** Cutaneous Lupus Erythematosus **Systemic Disease Context (Lupus Erythematosus, Systemic):** A chronic, relapsing, inflammatory, and often febrile multisystemic disorder of connective tissue (Semantic Type: Disease or Syndrome), characterized principally by involvement of the skin, joints, kidneys, and serosal membranes. It is of unknown etiology, but is thought to represent a failure of the regulatory mechanisms of the autoimmune system. The disease is marked by a wide range of system dysfunctions, an elevated erythrocyte sedimentation rate, and the formation of LE cells in the blood or bone marrow. **Diagnosis Threshold:** Clinically and histopathologically confirmed DLE/SCLE active for ≥ 3 months **Medical Methodology:** Targeted TYK2/JAK-STAT pathway inhibition (Deucravacitinib)

13. Study Procedures & Methodology

Management Pathway: Standard of care therapy + Investigational Product (Deucravacitinib or Placebo) **Education Protocol (HCP):** CLASI score assessment training and GCP compliance training **Education Protocol (Patient):** Drug administration guidelines and AE/SAE reporting mechanisms. **Quality Audit Mechanism:** Site monitoring, source data verification, and clinical safety laboratory assessments.

14. Observation & Follow-Up Schedule

Total Duration: Follow-up until study completion (Estimated Feb 2028) **Onsite Visit Cadence:** Pre-defined periodic clinical visits for CLASI scoring and safety monitoring. **Remote Monitoring Frequency:** Standard interval remote monitoring for site compliance. **Checklist Review Interval:** Regular safety and concomitant medication reviews throughout the study duration.

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]		Lead		
Statistician	[Name]	_____	[DD-MMM-YYYY]				