

1. Administrative & Study Identification

Full Study Title: A PHASE 1, OPEN-LABEL, RANDOMIZED CROSSOVER STUDY TO COMPARE THE SINGLE-DOSE PHARMACOKINETICS OF TWO FORMULATIONS OF PF-07976016 ADMINISTERED ORALLY TO HEALTHY ADULT PARTICIPANTS **Short Title/Acronym:** A Study to Learn How Different Forms of the Study Medicine Called PF-07976016 Are Taken Up Into the Blood in Healthy Adults **Protocol Number:** C5541005 **NCT Number:** NCT06670170 **Sponsor Name:** Pfizer **Investigational Product:** PF-07976016 (Formulation A and Formulation B) **Phase of Development:** Phase I **Medical Monitor:** Pfizer Clinical Research Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To measure and compare the amount of study drug in the blood after a single dose of two formulations of the study drug (PF-07976016 Formulation A and PF-07976016 Formulation B). **Secondary Objectives:** To evaluate the safety, tolerability, and other overall pharmacokinetic (PK) parameters of the different oral formulations in healthy adults.

2.2 Background & Rationale To evaluate how different physical forms (Formulation A vs. Formulation B) of the investigational drug PF-07976016 are absorbed into the bloodstream. This crossover pharmacokinetic assessment is medically necessary to select the optimal formulation for subsequent later-phase clinical efficacy and safety trials.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Active Comparator (Crossover of formulations) **Blinding:** Open-label **Randomization:** Randomized (Sequence 1: Formulation A then B; Sequence 2: Formulation B then A)

3.2 Planned Enrollment & Duration Target \$N\$: 12 Number of Sites: 1 (Pfizer Clinical Research Unit - New Haven, Connecticut, USA) **Estimated Study Start:** 05-Nov-2024 **Estimated Primary Completion:** 24-Dec-2024

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Overtly healthy adults \$>18\$ years of age, as determined by medical evaluation including medical

history, physical examination, laboratory tests, vital signs, and ECG assessments. 2. Willing to provide consent and follow study requirements. 3. Total body weight ≥ 110 lbs (50 kg).

4.2 Exclusion Criteria 1. Use of prescription or non-prescription drugs within a specified timeframe prior to the first dose of study intervention. 2. Positive urine drug test at screening/admission. 3. History of alcohol abuse or repeated binge drinking and/or any other illicit drug use or dependence within 6 months of screening. 4. Use of tobacco/nicotine-containing products more than the equivalent of 5 cigarettes/day or 2 chews of tobacco/day.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Single dose of PF-07976016 administered in up to 3 study periods. **Route of Administration:** Oral **Duration of Treatment:** Single-dose administration per period, with a 7-day washout period between each dose.

5.2 Concomitant & Prohibited Therapy Permitted: Only standard medications pre-approved by the clinical investigator for the management of unanticipated adverse events. **Prohibited:** Use of any unapproved prescription or non-prescription drugs within a specified timeframe prior to dosing; illicit drugs, and excessive alcohol or nicotine/tobacco.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures	:---
:---	:---	Screening	Up to Day -1 Informed Consent, Medical History, Physical Exam, ECG, Lab Tests, Urine Drug Screen
Period 1 (Baseline)	Day 1	Randomization, First Dose (Formulation A or B), PK Blood Draws, Safety Audits	Washout
Day 2 to 7	7-day washout period between doses	Period 2	
Day 8	Second Dose (Alternate Formulation), PK Blood Draws, Safety Assessments	End of Study	Follow-up Final Vital Signs, Exit Interview, IP Accountability, Final Safety Checks

(Note: Study periods repeat up to 3 times depending on sequence assignment)

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Area Under the Curve from time zero to infinity (AUC_{inf}) or to the last quantifiable concentration (AUC_{last}) to measure and compare the amount of study drug in the blood after a single oral dose. **Measurement**

Method: Pharmacokinetic blood sampling and central bioanalytical laboratory quantification.

7.2 Secondary & Exploratory Endpoints * Maximum observed plasma concentration (\$C_{max}\$) of PF-07976016. * Time to reach maximum plasma concentration (\$T_{max}\$) of PF-07976016. * Number and severity of treatment-emergent adverse events (TEAEs).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous monitoring via physical examinations, vital signs, ECGs, and clinical laboratory tests throughout the Phase 1 clinic stays. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal sponsor (Pfizer) review committee.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Pharmacokinetic (PK) Concentration Population, Safety Population. **Sample Size Justification:** Sized to provide adequate statistical power for establishing relative bioavailability and comparing the PK profiles of Formulation A vs. Formulation B in a crossover design. **Handling of Missing Data:** Standard Phase 1 pharmacokinetic handling; subjects with incomplete PK sampling profiles may be excluded from the primary \$AUC\$ comparisons.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Continuous onsite monitoring at the Phase 1 Clinical Research Unit in New Haven, CT. **Audit Trail:** Source document verification, continuous clinical observation, and eCRF locking for safety and bioanalytical data.

11. Study Identifiers & Governance

Primary ID: C5541005 **Registry Number:** NCT06670170 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn How Different Forms of the Study Medicine Called PF-07976016 Are Taken Up Into the Blood in Healthy Adults **Official Regulatory Title:** A PHASE 1, OPEN-LABEL, RANDOMIZED CROSSOVER STUDY TO COMPARE THE SINGLE-DOSE PHARMACOKINETICS OF TWO FORMULATIONS OF PF-07976016 ADMINISTERED ORALLY TO HEALTHY ADULT PARTICIPANTS

12. Clinical Framework (Pharmacokinetics Specific)

Primary Condition: Healthy Volunteers **Diagnosis Threshold:** Overtly healthy via standard clinical/laboratory bounds **Medical Methodology:** Phase 1 Pharmacokinetic (PK) Crossover Assessment **Optimization Target:** Comparative Bioavailability (Formulation A vs Formulation B)

13. Study Procedures & Methodology

Management Pathway: Clinical Pharmacology Unit inpatient/outpatient management **Education Protocol (HCP):** Investigator Brochure (IB) and Study Protocol site initiation training **Education Protocol (Patient):** Strict onsite orientation on dosing restrictions, dietary limits, and reporting AEs **Quality Audit Mechanism:** Bioanalytical sample tracking, Investigational Product (IP) accountability, exit interviews, and subject protocol adherence monitoring

14. Observation & Follow-Up Schedule

Total Duration: Approximately 6-8 weeks from screening to study exit. **Onsite Visit Cadence:** Clinical confinement during dosing periods, separated by a 7-day washout interval. **Remote Monitoring Frequency:** N/A (primarily clinic-based study). **Checklist Review Interval:** Daily during confinement periods (AEs and concomitant medications check), concluding with final vital signs and final safety checks at the end of the study follow-up.

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	Pfizer Clinical Investigator	_____	03-Apr-2026	Clinical Operations Lead	Pfizer Clinical Operations	_____	03-Apr-2026
Lead Statistician	Pfizer Biostatistics Lead	_____	03-Apr-2026				