

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 02-Apr-2026

1. Administrative & Study Identification

Full Study Title: Korea Comirnaty Post-marketing Surveillance
Official Title: A Prospective, Single-arm, Open-label, Non-interventional, Multicenter to Assess the Safety of BNT162b2 in Domestic Post-marketing Surveillance
Short Title/Acronym: Korea Comirnaty PMS
Protocol Number: C4591025
NCT Number: NCT05032976
Posting ID: 226251633705118252
Sponsor Name: Pfizer
Company: Pfizer
Trial Status: ACTIVE_NOT_RECRUITING
Investigational Product: BNT162b2 / Comirnaty (Tozinameran, Riltozinameran, Famtozinameran, Raxtozinameran)
Phase of Development: Not Applicable
Medical Monitor: Pfizer Pharmaceuticals Korea / Participating Site Investigators

2. Study Synopsis & Rationale

2.1 Study Objective
Primary Objective: To collect information on the safety of BNT162b2 products for subjects who have been administered the COVID-19 vaccine in routine clinical practice in Korea.
Secondary Objectives: To monitor adverse events (AEs) and assess the real-world safety profiles of various Comirnaty formulations across different age cohorts.

2.2 Background & Rationale This post-marketing surveillance study is conducted in accordance with the New Drug Re-Examination Guideline of the Ministry of Food and Drug Safety (MFDS) in Korea. It is designed to ensure the ongoing safety and track potential adverse reactions to the vaccine in a real-world, large-scale population setting.

3. Study Design & Methodology

3.1 Design Framework
Study Type: Observational (Post-marketing Surveillance)
Trial Status: ACTIVE_NOT_RECRUITING
Control Type: Single-arm (Non-comparative)
Blinding: Open-label
Randomization: N/A (Non-interventional)

3.2 Planned Enrollment & Duration
Target \$N\$: 12,000 subjects
Number of Sites: Multi-center (Korean health care centers)
Estimated Study Start: 18-Mar-2022
Estimated Primary Completion: 04-Mar-2027

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Korean subjects who are eligible for administration of BNT162b2 products (including bivalent COVID-19 vaccines for Omicron) according to the locally approved and authorized label (indication, age criteria, etc.). 2. Evidence of a personally signed and dated data privacy statement indicating that the subject (or a legally acceptable representative/parent for subjects under 19) has been informed of all pertinent aspects of the study.

4.2 Exclusion Criteria 1. Subjects who involved in the contraindications of use indicated in the locally approved and authorized label. 2. Subjects with a history of hypersensitivity to any ingredients of this product or this product.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: 0.2 mL or 0.3 mL depending on age group and specific formulation cap color (e.g., purple, grey, maroon, or orange caps). Administered as a single dose, a 2-dose primary series (typically 21 days apart), or a 3-dose series for specific pediatric or immunocompromised groups. **Route of Administration:** Intramuscular injection (IM). **Duration of Treatment:** Routine vaccination days (1 to 3 doses) with a 28-day safety follow-up period per administration.

5.2 Concomitant & Prohibited Therapy Permitted: Standard concomitant therapies as per routine clinical practice. **Prohibited:** The interchangeability of BNT162b2 with COVID-19 vaccines from other manufacturers to complete the primary vaccination course has not been established and is not recommended.

6. Study Timeline & Visit Schedule

(Note: There are no study-mandated visits; interactions align with routine clinical practice) | Visit ID | Window (Days) | Key Activities/Procedures | | :--- | :--- | :--- | | **Vaccination** | Day 0 | Vaccine administration, Data Privacy Consent, baseline medical record review | | **Safety Follow-up** | Days 1-27 | Daily safety questionnaire/eDiary entries post-injection | | **End of Follow-up** | Day 28 (Post-dose) | Collection of safety questionnaires 28 days post-second injection or booster dose (via mail or during next visit) |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoints Definition: 1. Number of subjects with Solicited adverse events: Solicited adverse events within 1 week

(Day 1-7) after each dose of BNT162b2 products, and a number of subjects with such events will be collected. 2. Number of subjects with Unsolicited adverse events: Unsolicited adverse events within 28 days by its last dose, and a number of subjects with such events will be collected. **Measurement Method:** Patient (Subject) Reported Outcomes (PRO) collected via electronic diaries (eDiaries) or paper questionnaires, and data extracted from medical records.

7.2 Secondary Endpoints 1. Number of subjects with Adverse Events (AEs). 2. Number of subjects with Serious Adverse Events. 3. Number of subjects with Expected Adverse Events.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Collected via daily case report forms (CRF). If an application is used, it is automatically sent as an eCRF; if a paper questionnaire is used, it is collected at the next vaccination visit or by mail. **Serious Adverse Events (SAEs):** Evaluated and reported according to routine pharmacovigilance practices and the Ministry of Food and Drug Safety (MFDS) guidelines. **Data Monitoring Committee (DMC):** Internal safety monitoring managed by Pfizer Pharmaceuticals.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all enrolled subjects who received at least one dose of the vaccine and provided safety data). **Sample Size Justification:** Target $N=12,000$ to fulfill the requirements of the MFDS New Drug Re-Examination Guideline for robust post-marketing safety evaluation. **Handling of Missing Data:** Descriptive statistics based on observational real-world data without formal imputation for missing PROs.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: Conducted in accordance with Good Pharmacoepidemiology Practices and the New Drug Re-Examination Guideline of the Korean MFDS. **Monitoring Plan:** Post-marketing surveillance agreements are established with investigator sites. Investigators prepare CRFs for all enrolled subjects sequentially. **Audit Trail:** eCRFs automatically transmit data via the study application. Paper questionnaires are tracked and collected systematically 28 days after subsequent doses.

11. Study Identifiers & Governance

Primary ID: C4591025 **Registry Number:** NCT05032976 **Posting ID:** 226251633705118252 **Sponsor/Lead Organization:** Pfizer **Company:** Pfizer **Study Title:** Korea Comirnaty Post-marketing Surveillance **Official Regulatory Title:** A Prospective, Single-arm, Open-label, Non-interventional, Multicenter to Assess the Safety of BNT162b2 in Domestic Post-marketing Surveillance **Trial Status:** ACTIVE_NOT_RECRUITING

12. Clinical Framework (COVID-19 Specific)

Disease Areas / Primary Conditions: COVID-19, Vaccines, Adverse Effects, SARS-CoV-2, Safety **Preferred UMLS Name:** Pharmacovigilance **Diagnosis Threshold:** N/A (Prophylactic Vaccination) **Medical Methodology:** mRNA Vaccine Administration (BNT162b2) **Optimization Target:** Broad population immunization with rigorous safety tracking

13. Study Procedures & Methodology

Management Pathway: Routine COVID-19 vaccination pathway per national Korean guidelines **Education Protocol (HCP):** Site investigators trained on proper vaccine dilution (if applicable), dosing volumes based on cap color, and sequential patient enrollment procedures **Education Protocol (Patient):** Instructions given on completing daily safety diaries via app or paper formats **Quality Audit Mechanism:** Institutional site agreements and automated eCRF tracking to ensure surveillance fidelity

14. Observation & Follow-Up Schedule

Total Duration: Up to 28 days post-final dose per subject (Study umbrella duration spans up to 2027) **Onsite Visit Cadence:** Aligns purely with routine vaccination schedules (e.g., Day 0, Day 21, or 3+ months for boosters) **Remote Monitoring Frequency:** Daily PRO reporting by patients post-vaccination **Checklist Review Interval:** Safety questionnaires are collected 28 days post-second injection and 28 days post-booster

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	____	[DD-MMM-YYYY]				
Statistician	[Name]	____	[DD-MMM-YYYY]				