

1. Administrative & Study Identification

Full Study Title: FIRST-IN-HUMAN (FIH), OPEN-LABEL, PHASE 1 DOSE ESCALATION AND EXPANSION STUDY DESIGNED TO EVALUATE THE SAFETY, TOLERABILITY, PK, PD, AND PRELIMINARY CLINICAL ACTIVITY OF PF-07826390 AS A SINGLE AGENT OR IN COMBINATION TREATMENT FOR PARTICIPANTS WITH ADVANCED SOLID TUMORS **Short Title/Acronym:** A Study to Learn About How Different Amounts of the Study Medicine PF-07826390 Act in the Body of People With Cancer When Taken Alone or With Other Anti-cancer Medicines **Protocol Number:** C5321001 **NCT Number:** NCT06546553 **Sponsor Name:** Pfizer Inc. **Investigational Product:** PF-07826390 (LILRB1/2 antagonist) alone or in combination with Sasanlimab (anti-PD-1) **Phase of Development:** Phase 1 **Medical Monitor:** Pfizer Inc. Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety, tolerability, pharmacokinetics (PK), pharmacodynamics (PD), and preliminary clinical activity (Dose-limiting toxicities [DLTs] and Objective Response Rate [ORR]) of PF-07826390 as a single agent or in combination treatment for participants with advanced solid tumors. **Secondary Objectives:** To evaluate Objective Response (OR), Duration of Response (DoR), Progression-Free Survival (PFS), and Pharmacokinetics.

2.2 Background & Rationale To address the unmet need in patients with advanced or metastatic solid tumors (e.g., Non-Small Cell Lung Cancer [NSCLC], Colorectal Cancer [CRC], Renal Cell Carcinoma [RCC]) who have exhausted standard therapies. The study aims to determine the optimal dose and observe how PF-07826390—a novel bispecific antibody targeting LILRB1 and LILRB2 immunosuppressive receptors—acts in the body alone or when combined with the anti-PD-1 therapy Sasanlimab to overcome immune resistance in the tumor microenvironment.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Dose escalation and expansion cohorts) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target **NS\$:** 190 participants **Number of Sites:** Multinational **Estimated Study Start:** September

2024 **Estimated Primary Completion:** October 2025 (Follow-up continues for up to 4 years)

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Histological or cytological diagnosis of advanced, unresectable, and/or metastatic or relapsed/refractory solid tumor (specifically including NSCLC, CRC, RCC). 2. Prior progression on standard therapies (e.g., prior anti-PD-(L)1 therapy if approved/available; platinum-based chemo for NSCLC; specific regimens for MSS CRC and RCC). 3. At least 1 measurable lesion based on RECIST 1.1 that has not been previously irradiated.

4.2 Exclusion Criteria 1. Treatment with any systemic anticancer therapy within 4 weeks or 5 half-lives (whichever is shorter) prior to planned first dose. 2. Active or history of clinically significant autoimmune disease or other medical condition that required chronic systemic immunosuppressive therapy within recent 2 years. 3. Prior treatment with another LILRB1 (ILT2), LILRB2 (ILT4), and/or LILRB1/2 (B1 and B2) antagonist antibodies or pathway targeting agents, including HLA conformers and HLA-G antibodies.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Dose escalation of PF-07826390 administered alone or in combination with Sasanlimab (administered every 4 weeks). **Route of Administration:** Intravenous / Subcutaneous (Sasanlimab is administered subcutaneously). **Duration of Treatment:** Until disease progression or unacceptable toxicity (up to approximately 2 years, followed by up to 2 years of post-treatment follow-up).

5.2 Concomitant & Prohibited Therapy Permitted: Standard supportive care for advanced solid tumors. **Prohibited:** Systemic anticancer therapy within 4 weeks of the first dose, and chronic systemic immunosuppressive therapy.

6. Study Timeline & Visit Schedule

| Visit ID | Window (Days) | Key Activities/Procedures | | :---
| :--- | :--- | | **Screening** | Day -28 to 0 | Informed Consent, Medical History, Tumor Tissue sample collection, Baseline Scans |
| **Baseline** | Day 0 | First Dose, PK/PD evaluations, Safety assessments | | **Interim** | Weekly | Weekly study visits during active treatment: Safety Labs, DLT monitoring (especially First cycle, Day 1 to 28), Efficacy Scans | | **End of Study** | Follow-up | Stopping medication at approx. 2 years; followed for another 2 years for disease progression/survival tracking |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Number of participants with Dose-Limiting Toxicities (DLTs) within the first cycle (Day 1 up to Day 28); Incidence of Treatment-Emergent Adverse Events (AEs); and Objective Response Rate (ORR). **Measurement Method:** Clinical safety evaluation and Investigator assessment of objective response based on RECIST 1.1 criteria.

7.2 Secondary & Exploratory Endpoints * Objective Response (OR) * Duration of Response (DoR) * Progression-Free Survival (PFS) * Pharmacokinetics (PK) profile

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Standard clinical monitoring including untoward medical occurrences reported from the start of treatment and up to 90 days after the last dose or start of new anticancer therapy. **Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness. **Data Monitoring Committee (DMC):** Internal safety review committee governing dose-escalation decisions.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants receiving ≥ 1 dose) and Efficacy Evaluable Set. **Sample Size Justification:** Target $N=190$; calculated based on standard Phase 1 dose escalation (e.g., modified toxicity probability intervals) and expansion cohort requirements. **Handling of Missing Data:** Standard handling mechanisms for early phase oncology trials; missing tumor assessments censored at the last evaluable scan.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Regular onsite and remote monitoring to ensure protocol adherence and patient safety during dose escalation. **Audit Trail:** Electronic Case Report Form (eCRF) locking and standard source data verification.

11. Study Identifiers & Governance

Primary ID: C5321001 **Registry Number:** NCT06546553 **Sponsor/Lead Organization:** Pfizer Inc. **Study Title:** A Study to Learn About How Different Amounts of the Study Medicine PF-07826390 Act in the

Body of People With Cancer When Taken Alone or With Other Anti-cancer Medicines. **Official Regulatory Title:** FIRST-IN-HUMAN (FIH), OPEN-LABEL, PHASE 1 DOSE ESCALATION AND EXPANSION STUDY DESIGNED TO EVALUATE THE SAFETY, TOLERABILITY, PK, PD, AND PRELIMINARY CLINICAL ACTIVITY OF PF-07826390 AS A SINGLE AGENT OR IN COMBINATION TREATMENT FOR PARTICIPANTS WITH ADVANCED SOLID TUMORS.

12. Clinical Framework

Primary Condition: Advanced or Metastatic Solid Tumors (including NSCLC, CRC, RCC) **Diagnosis Threshold:** Histological or cytological diagnosis of advanced, unresectable, and/or metastatic solid tumor **Medical Methodology:** PF-07826390 administered as a single agent or in combination with Sasanlimab **Optimization Target:** Safety, Tolerability, and anti-tumor efficacy (ORR, PFS)

13. Study Procedures & Methodology

Management Pathway: Experimental phase 1 clinical pathway for advanced solid tumors refractory to standard therapies **Education Protocol (HCP):** Dose escalation safety monitoring and DLT management training **Education Protocol (Patient):** Informed consent, weekly schedule expectations, and safety self-reporting **Quality Audit Mechanism:** Clinical site monitoring, safety data review, and PK/PD sample auditing

14. Observation & Follow-Up Schedule

Total Duration: Up to 4 years **Onsite Visit Cadence:** Weekly study visits during active treatment **Remote Monitoring Frequency:** Continuous safety and survival reporting **Checklist Review Interval:** Every cycle (e.g., 28 days)

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead	
Statistician	[Name]	_____	[DD-MMM-YYYY]				