

1. Administrative & Study Identification

Official Study Title: A Phase IIb Study to Characterize the Efficacy and Safety of Adjuvant Ribociclib Plus Endocrine Therapy in a Close-to-Clinical Practice Patient Population With HR+ HER2-Early Breast Cancer **Study Title:** Phase IIb Study of Ribociclib + ET in Early Breast Cancer **Short Title/Acronym:** Adjuvant WIDER **Protocol Number:** CLEE011012001 **NCT Number:** NCT05827081 **Trial Status:** Recruiting **Sponsor Name:** Novartis Pharmaceuticals **Investigational Product:** Ribociclib + Endocrine Therapy (ET) **Phase of Development:** Phase IIb **Medical Monitor:** Novartis Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To characterize the efficacy and safety of the combination of ribociclib and standard adjuvant endocrine therapy (ET) on invasive breast cancer-free survival (iBCFS) in a close-to-clinical practice patient population. * **Secondary Objectives:** To evaluate Distant Disease-Free Survival (DDFS), Invasive Disease-Free Survival (iDFS), and the incidence and severity of adverse events (AEs) associated with the combination regimen.

2.2 Background & Rationale To evaluate the effectiveness and safety of adding targeted therapy (ribociclib) to standard hormone treatment (endocrine therapy) after surgery for people with early-stage breast cancer (HR-positive, HER2-negative). This study aims to provide real-world insights into how well this treatment combination reduces the risk of cancer recurrence in a wide patient population, including both pre/post-menopausal women and men, in a setting close to standard clinical practice.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Single-arm * **Blinding:** Open-label * **Randomization:** N/A (Single-arm study)

3.2 Planned Enrollment & Duration * **Target \$N\$:** 1,400 participants * **Number of Sites:** Multicenter (United States + 14 other countries) * **Estimated Study Start:** 27-Feb-2024 * **Estimated Primary Completion:** 07-Sep-2029

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Adult, male or female ≥ 18 years of age at the time of informed consent. 2. Histologically and/or cytologically confirmed diagnosis of HR+ (estrogen/progesterone receptor positive) and HER2- Early Breast Cancer (EBC) with surgical resection where tumor was removed completely and microscopic margins are free from tumor. 3. Anatomic Stage Group III, Anatomic Stage Group IIB, or a subset of Anatomic Stage Group IIA. 4. Eastern Cooperative Oncology Group (ECOG) Performance Status of 0, 1 or 2. 5. Adequate bone marrow and organ function. 6. Allowed prior standard neoadjuvant and/or adjuvant ET (including tamoxifen or toremifene); enrollment must occur within 36 months of prior ET start date and participants should have at least 3 years remaining of endocrine adjuvant therapy. 7. ECG values assessed by KardiaMobile-6L or standard 12-lead ECG demonstrating a QTcF interval < 450 msec and mean resting heart rate of 50-99 beats per minute.

4.2 Exclusion Criteria 1. Distant metastases of breast cancer beyond regional lymph nodes (Stage IV according to AJCC 8th edition) and/or evidence of recurrence after curative surgery. 2. Concurrent use of other anti-neoplastic therapy, with the exception of adjuvant ET. 3. Clinically significant, uncontrolled heart disease and/or cardiac repolarization abnormality. 4. Any other concurrent severe and/or uncontrolled medical condition limiting life expectancy to ≤ 5 years. 5. Pregnant or breast-feeding (lactating) women, or women of child-bearing potential (CBP) unless using highly effective methods of contraception while taking study treatment and for 21 days after stopping treatment.

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Dosage/Frequency:** Ribociclib 400 mg orally once daily on Days 1 to 21 of a 28-day cycle, in combination with daily ET. * **Route of Administration:** Oral * **Duration of Treatment:** 36 months (approximately 39 cycles) from the date of the first dose.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Standard adjuvant endocrine therapy (letrozole, anastrozole, exemestane, tamoxifen, or toremifene; combined with goserelin or leuprolide for pre/peri-menopausal women and men). * **Prohibited:** Concurrent anti-neoplastic therapies other than standard adjuvant ET.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	-28 to 0	Informed Consent, Medical History, Confirmation of Eligibility, baseline ECG via

KardiaMobile-6L or 12-lead. | | **Baseline** | Day 0 | Cycle 1 Day 1, First Dose of Ribociclib + ET. | | **Interim** | Cycles 1 - 39 | Regular clinic visits, heart function checks (ECG monitoring), lab assessments, and safety/efficacy monitoring. | | **End of Study** | Follow-Up | 30-day Safety Follow-up call after last dose; long-term follow-up until 48 months after the last participant's first dose (EOS). |

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Invasive Breast Cancer Free Survival (iBCFS) rate at 3 years, defined as the time from the first dose to the first event of invasive ipsilateral breast tumor recurrence, local/regional invasive recurrence, distant recurrence, death, or contralateral invasive BC. * **Measurement Method:** Evaluated by the Investigator using STEEP criteria version 2.0 (Standardized Definitions for Efficacy End Points in Adjuvant Breast Cancer Trials).

7.2 Secondary & Exploratory Endpoints * Incidence and severity of adverse events (AEs) using CTCAE v4.03. * Invasive Disease-Free Survival (iDFS). * Distant Disease-Free Survival (DDFS).

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Standard collection via regular clinic visits, laboratory tests, and continuous heart function (ECG) monitoring graded per CTCAE v4.03.
 - **Serious Adverse Events (SAEs):** Reported within 24 hours of Investigator awareness.
 - **Data Monitoring Committee (DMC):** Internal/External safety committee per Novartis standard protocols.
-

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Intent-to-Treat (ITT) population for efficacy outcomes; Safety Set for all participants receiving at least one dose of study treatment.
 - **Sample Size Justification:** $N = 1,400$ adequately powered to observe long-term survival events in an open-label, close-to-clinical practice setting.
 - **Handling of Missing Data:** Censoring methodologies applied per STEEP 2.0 guidelines for time-to-event survival analysis.
-

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
 - **Monitoring Plan:** Combination of onsite and remote monitoring visits to ensure subject safety and data integrity.
 - **Audit Trail:** Robust clinical data management including eCRF locking and automated data clarification forms.
-

11. Study Identifiers & Governance

- **Primary ID:** CLEE011012001
 - **Registry Number:** NCT05827081
 - **Sponsor/Lead Organization:** Novartis Pharmaceuticals
 - **Trial Status:** Recruiting
 - **Study Title:** Adjuvant WIDER
 - **Official Regulatory Title:** A Phase IIIB Study to Characterize the Efficacy and Safety of Adjuvant Ribociclib Plus Endocrine Therapy in a Close-to-Clinical Practice Patient Population With HR+ HER2- Early Breast Cancer
-

12. Clinical Framework (Breast Cancer Specific)

- **Primary Condition:** Early Breast Cancer (EBC)
 - **Diagnosis Threshold:** HR-positive, HER2-negative, Anatomic Stage Group III, IIB, or a subset of IIA
 - **Medical Methodology:** CDK4/6 Inhibitor (Targeted Therapy) combined with Adjuvant Endocrine Therapy
 - **Optimization Target:** Maximize Invasive Breast Cancer Free Survival (iBCFS)
-

13. Study Procedures & Methodology

- **Management Pathway:** Standard oncology clinical pathway for Adjuvant Breast Cancer.
 - **Education Protocol (HCP):** Site initiation visits covering study drug administration, ET combinations, and ECG monitoring compliance.
 - **Education Protocol (Patient):** Education on oral medication compliance (21 days on, 7 days off) and recognizing potential AEs.
 - **Quality Audit Mechanism:** Continuous safety monitoring and AE/SAE tracking by the sponsor.
-

14. Observation & Follow-Up Schedule

- **Total Duration:** 36 months of active treatment + follow-up (up to 48 months for EOS)
 - **Onsite Visit Cadence:** Regular 28-day cycles (Cycles 1-39)
 - **Remote Monitoring Frequency:** 30-day Safety Follow-up call post-treatment
 - **Checklist Review Interval:** Regular review at each cycle visit for drug accountability and safety.
-

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				