

1. Administrative & Study Identification

Full Study Title A Study to Assess Efficacy, Safety, Tolerability, Pharmacokinetics (PK) and Pharmacodynamics (PD) of R07204239 in Combination With Tirzepatide in Participants With Obesity or Overweight With At Least One Weight-related Comorbidity **Short Title/Acronym** GYMINDA Study **Protocol Number** BC45538 / 2024-519561-22-00 **NCT Number** NCT06965413 **Posting ID** 266493664382974427 **Trial Status** ACTIVE_NOT_RECRUITING (Active but not currently recruiting) **Sponsor Name** Roche **Investigational Product** R07204239 (Emugrobart) in combination with Tirzepatide **Phase of Development** Phase 2 **Medical Monitor** [Name and Contact Information]

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To assess the effect of R07204239 (emugrobart) in combination with tirzepatide, compared to placebo in combination with tirzepatide, on body weight loss after 48 weeks of treatment in adults with obesity or overweight with at least one weight-related comorbidity, but without diabetes mellitus (DM). * **Secondary Objectives:** To assess absolute change from baseline in body weight, Body Mass Index (BMI), waist-to-height ratio, waist circumference, total body fat mass (measured by DXA), and total lean body mass (measured by DXA) at Week 48.

2.2 Background & Rationale To address the clinical need for the preservation of lean body mass during pharmacologically induced weight loss. While incretin therapies such as tirzepatide effectively reduce body weight, they often lead to unintended loss of skeletal muscle. Emugrobart (R07204239) is an anti-myostatin candidate designed to promote skeletal muscle growth and maintenance, potentially preserving lean body mass and improving the overall quality of weight loss when used as combination therapy.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Placebo-controlled * **Blinding:** Double-blind * **Randomization:** Randomized (to one of 4 treatment arms) * **Trial Status:** ACTIVE_NOT_RECRUITING

3.2 Planned Enrollment & Duration * Target SN\$: 285 * Number of Sites: [Global and Regional breakdown] *** Estimated Study Start:** [DD-MMM-YYYY] *** Estimated Primary Completion:** August 2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Male or female subjects ≥ 18 years of age. 2. BMI ≥ 30.0 kg/m² (additional weight-related comorbidities are not required for inclusion) OR BMI ≥ 27.0 kg/m² and < 30.0 kg/m² with at least one weight-related comorbidity such as: hypertension, dyslipidemia, obstructive sleep apnea, and any cardiovascular disease. 3. History of at least one self-reported unsuccessful dietary or exercise effort to lose body weight. 4. Weight stability: self-reported change in body weight < 5 kg (11 lbs) within 3 months prior to screening.

4.2 Exclusion Criteria 1. Prior history or diagnosis of DM. 2. Presence of non-proliferative diabetic retinopathy requiring acute therapy, proliferative diabetic retinopathy, or diabetic macular edema. 3. Have obesity induced by other endocrinologic disorders, participation in unbalanced/extreme diets, or prior or planned surgical treatment for obesity. 4. Endoscopic and/or device-based therapy for obesity or device removal within 6 months prior to screening. 5. Have a known clinically significant gastric emptying abnormality. 6. Have any of the following cardiovascular conditions within 6 months prior to screening: acute myocardial infarction, cerebrovascular accident (stroke), unstable angina, or hospitalization due to congestive heart failure (CHF). 7. Have evidence of significant active, uncontrolled cardiovascular, autoimmune, endocrine, renal, hepatic, dermatological, chronic respiratory or gastrointestinal disease, a neurological or psychiatric condition, or a history of any neuromuscular disorder or autoimmune/inflammatory disorders that may cause muscle wasting or medical condition capable of constituting a risk when taking the study medication or interfering with the interpretation of data, as judged by the investigator at screening. 8. Have evidence of a significant, uncontrolled endocrine abnormality. 9. Have a history of an active or untreated malignancy or are in remission from a clinically significant malignancy. 10. Have evidence of a significant, active autoimmune abnormality. 11. Have anemia. 12. Have signs and symptoms of any other liver disease other than nonalcoholic fatty liver disease. 13. Have an average weekly alcohol intake that exceeds 21 units per week (males) and 14 units per week (females).

5. Intervention & Treatment Plan

5.1 Investigational Regimen * Dosage/Frequency: R07204239 (low, medium, or high dose) or matching placebo administered every 4 weeks (Q4W). Tirzepatide (trade names: Zepbound, Mounjaro)

administered as background therapy weekly (QW) and up-titrated according to approved prescribing information. * **Route of Administration:** Subcutaneous (SC) injection. * **Duration of Treatment:** 48-week core treatment period, followed by a 24-week treatment extension period (where tirzepatide is stopped and participants receive R07204239 or placebo), and a 24-week post-treatment follow-up period.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Tirzepatide (as background therapy during the 48-week core period). * **Prohibited:** Other weight loss medications, unbalanced/extreme diets, and endoscopic/surgical interventions for obesity.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Eligibility Labs, DXA Scan
Baseline	Day 0	Randomization, First Dose of R07204239/Placebo and Tirzepatide
Interim	Core Period	Subcutaneous injections Q4W (Investigational Drug) and QW (Tirzepatide); Safety & Efficacy Assessments
End of Study	Week 96	Final 24-Week Post-Treatment Follow-up, Final Vital Signs, DXA Scan, Exit Interview

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Percent Change From Baseline in Body Weight at Week 48. * **Measurement Method:** Standardized body weight measurements in clinic.

7.2 Secondary & Exploratory Endpoints * Absolute Change From Baseline in Body Weight at Week 48. * Change From Baseline in Body Mass Index (BMI) at Week 48. * Change From Baseline in Waist-to-height Ratio and waist circumference at Week 48. * Change from baseline in total body fat mass and total lean body mass measured by Dual-energy X-ray Absorptiometry (DXA) at Week 48.

8. Safety & Adverse Event Reporting

- Adverse Event (AE) Monitoring:** Standard collection via patient reporting, HCP interview, and physical examinations.
 - Serious Adverse Events (SAEs):** Reported within 24 hours of site awareness.
 - Data Monitoring Committee (DMC):** Internal/External safety monitoring mechanisms as established by Roche.
-

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Intent-to-Treat (ITT) population for primary weight loss outcomes; Safety Set for all AE/SAE evaluations.
 - **Sample Size Justification:** Appropriately powered ($\alpha = 0.05$, $\text{Power} \geq 80\%$) with a targeted enrollment of 285 to detect statistically significant differences in body weight and lean mass changes between the combination and active background (tirzepatide + placebo) arms.
 - **Handling of Missing Data:** Multiple Imputation (MI) and Mixed Models for Repeated Measures (MMRM) where applicable.
-

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki.
 - **Monitoring Plan:** Routine onsite and remote monitoring visits to ensure protocol compliance and accurate DXA image central reading.
 - **Audit Trail:** Electronic Case Report Form (eCRF) locking and standard data clarification procedures.
-

11. Study Identifiers & Governance

Primary ID: BC45538 **Registry Number:** NCT06965413 **Posting ID:** 266493664382974427 **Trial Status:** ACTIVE_NOT_RECRUITING **Sponsor/Lead Organization:** Roche **Study Title:** A Study to Assess Efficacy, Safety, Tolerability, Pharmacokinetics (PK) and Pharmacodynamics (PD) of R07204239 in Combination With Tirzepatide in Participants With Obesity or Overweight With At Least One Weight-related Comorbidity **Official Regulatory Title:** A Randomized, Double-blind, Placebo-controlled Phase 2 Trial to Assess Efficacy, Safety, and Tolerability of R07204239 in Combination With Tirzepatide in Participants With Obesity or Overweight With At Least One Weight-related Comorbidity

12. Clinical Framework (Obesity Specific)

Disease Areas: Obesity, Overweight, Overweight With One Weight Related Comorbidity **Diagnosis Threshold:** BMI ≥ 30.0 kg/m² OR BMI ≥ 27.0 kg/m² with comorbidity **Medical Methodology:** Anti-myostatin + GIP/GLP-1 RA combination therapy **Optimization Target:** Significant body weight loss with lean body mass preservation

13. Study Procedures & Methodology

Management Pathway: Pharmacological weight management coupled with lifestyle modifications **Education Protocol (HCP):** Protocol-specific training for combination therapy dosing, SC injection schedules, and DXA standardization **Education Protocol (Patient):** Standardized nutritional counseling and physical activity guidelines **Quality Audit Mechanism:** Independent central reading for DXA scans and clinical site adherence monitoring

14. Observation & Follow-Up Schedule

Total Duration: 96 weeks (48 weeks core treatment + 24 weeks treatment extension + 24 weeks follow-up) **Onsite Visit Cadence:** Protocol-defined intervals, typically aligned with Q4W investigational drug dosing **Remote Monitoring Frequency:** [As per site specific monitoring plan] **Checklist Review Interval:** [Interval tracking for body weight and dietary adherence]

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]				Lead
Statistician	[Name]	_____	[DD-MMM-YYYY]				