

## **1. Administrative & Study Identification**

**Full Study Title:** AN INTERVENTIONAL EFFICACY AND SAFETY, PHASE 3, DOUBLE-BLIND, PARALLEL GROUP STUDY TO INVESTIGATE INTERMITTENT PREVENTION OF MENSTRUAL MIGRAINE WITH RIMEGEPANT COMPARED WITH PLACEBO IN WOMEN PARTICIPANTS 18 TO 45 YEARS OF AGE **Short Title/ Acronym:** A Study to Learn About the Study Medicine Called Rimegepant in Women When Used for Intermittent Prevention of Menstrual Migraine **Protocol Number:** C4951063 **NCT Number:** NCT06641466 **Posting ID:** 271503695776790138 **Sponsor Name / Company:** Pfizer **Investigational Product:** Rimegepant (Trade Name: Nurtec; ATC Code: N02CD06, 75 mg Orally Disintegrating Tablet [ODT]) **Phase of Development:** Phase 3 **Trial Status:** Recruiting (Currently recruiting participants) **Medical Monitor:** Pfizer Clinical Operations Lead

---

## **2. Study Synopsis & Rationale**

**2.1 Study Objective Primary Objective:** To evaluate the efficacy and safety of rimegepant when administered during the peri-menstrual period (PMP) for the intermittent prevention of migraine in women who experience menstrual migraine attacks.

**Secondary Objectives:** To evaluate the safety, tolerability, and additional efficacy measures of rimegepant compared to placebo in reducing the frequency, severity, and acute medication use associated with menstrual migraines over multiple cycles.

**2.2 Background & Rationale** To address the gap in effective preventative treatments for menstrual migraines. Menstrual migraine is a specific type of migraine headache that is closely associated with the menstrual cycle, typically occurring during the late luteal phase (just before menstruation) or in the first few days of menstruation. It is primarily triggered by the natural drop in estrogen levels experienced at these times. These migraines are often more severe, prolonged, and disabling than non-menstrual migraines, characterized by intense throbbing pain, usually on one side of the head, and frequently accompanied by nausea, vomiting, and heightened sensitivity to light and sound. This study evaluates whether intermittent, targeted dosing of rimegepant during the 7-day peri-menstrual period can safely and effectively prevent these specific migraine attacks.

---

### 3. Study Design & Methodology

**3.1 Design Framework Study Type:** Interventional **Control Type:** Placebo-controlled **Blinding:** Double-blind **Randomization:** 1:1 Parallel Group

**3.2 Planned Enrollment & Duration Targeted Enrollment (N):** 723 participants. Multicenter large-scale Phase 3 enrollment. **Number of Sites:** 121 locations globally (including USA, Canada, Poland, etc.) **Estimated Study Start:** 11-Mar-2025 **Estimated Primary Completion:** 29-Nov-2026

---

### 4. Subject Selection (Eligibility Criteria)

**4.1 Inclusion Criteria** 1. Female participants 18 to 45 years of age. 2. Participant has regular menstrual cycles  $\geq 24$  days and  $\leq 34$  days. 3. A minimum 1-year history of migraine (with or without aura) consistent with a diagnosis according to the International Classification of Headache Disorders, 3rd Edition. 4. A history of menstrual migraine attacks of at least 3 months, experiencing at least 1 migraine attack during the perimenstrual period in at least 2 out of 3 menstrual cycles immediately prior to screening. 5. If the participant is receiving a permitted background continuous prophylactic migraine medication, the medication dose must be stable for at least 3 months prior to the Screening visit and, the dose is not expected to change during the course of the study.

**4.2 Exclusion Criteria** 1. Greater than 6 migraine days per month that are outside of the perimenstrual period in the 3 months prior to Screening. 2. A diagnosis of chronic migraine or a history of more than 14 headache days per month on average, in the 3 months prior to Screening. 3. History of retinal migraine, basilar migraine, or hemiplegic migraine. 4. Current diagnosis of schizophrenia, bipolar, or borderline personality disorder. 5. Other pain syndromes (eg, fibromyalgia, complex regional pain syndrome), or significant neurological disorders (other than migraine), or other medical conditions (including endocrine and gynecological eg, severe dysmenorrhea).

---

### 5. Intervention & Treatment Plan

**5.1 Investigational Regimen Dosage/Frequency:** Rimegepant 75 mg ODT or Placebo taken by mouth once daily for 7 days each month. Dosing begins 3 days before the expected start of the menstrual period. **Route of Administration:** Oral (Orally Disintegrating Tablet) **Duration of Treatment:** 5 months (5 menstrual cycles) in the double-blind phase, with an option to continue in an open-label treatment phase.

**5.2 Concomitant & Prohibited Therapy Permitted:** Permitted background continuous prophylactic migraine medications, provided the dose is stable for at least 3 months prior to the Screening visit and is not expected to change during the study. **Prohibited:** Concurrent investigational drugs and agents strictly conflicting with CGRP receptor antagonists or standard migraine assessments.

---

## 6. Study Timeline & Visit Schedule

| Visit ID     | Window           | Key Activities/Procedures                                                                      |
|--------------|------------------|------------------------------------------------------------------------------------------------|
| Screening    | Prior to Cycle 1 | Informed Consent, Medical History, Verification of Migraine History & Menstrual Cycle length.  |
| Baseline     | Cycle 1 Day 0    | Randomization, dispensing of rimegepant/placebo, instruction on ePRO diary.                    |
| Interim      | Cycles 1 to 4    | Ongoing 7-day monthly dosing, daily ePRO diary for headache days/intensity, safety monitoring. |
| End of Study | Cycle 5          | Final efficacy assessment, AE monitoring, option to transition into the open-label phase.      |

---

## 7. Outcome Measures (Endpoints)

**7.1 Primary Endpoint Definition:** Mean change from the Observation Period in the number of migraine days per 5-day perimenstrual period across each cycle of the Double-Blind Treatment Phase.  
**Measurement Method:** Electronic Patient-Reported Outcome (ePRO) Diary.

**7.2 Secondary & Exploratory Endpoints**

1. Mean change from the Observation Period in number of headache days per 5-day perimenstrual period across each cycle of the Double-Blind Treatment Phase.
2. Percentage of participants with  $\geq 50\%$  reduction from the Observation Period in number of migraine days per 5-day perimenstrual period across each cycle of the Double-Blind Treatment Phase (50% responder).
3. Mean change from the Observation Period in number of acute migraine medication days per 5-day perimenstrual period across each cycle of the Double-Blind Treatment Phase.
4. Overall safety and tolerability profile of intermittent rimegepant administration.

---

## 8. Safety & Adverse Event Reporting

**Adverse Event (AE) Monitoring:** Collected via daily ePRO diary entries and site visits/interviews. **Serious Adverse Events (SAEs):** Reported by site personnel to the sponsor within 24 hours of awareness. **Data Monitoring Committee (DMC):** Internal and external safety review committees managed by Pfizer.

---

## 9. Statistical Analysis Plan (SAP) Summary

**Analysis Populations:** Intent-to-Treat (ITT) population for primary efficacy analysis; Safety Set for all tolerability and AE summaries. **Sample Size Justification:** Powered to detect a statistically significant mean change in perimenstrual migraine days between the rimegepant 75 mg and placebo groups ( $\alpha = 0.05$ ). **Handling of Missing Data:** Mixed-Effect Model Repeated Measure (MMRM) and/or Multiple Imputation for continuous efficacy endpoints across the 5 menstrual cycles.

---

## 10. Quality Assurance & Regulatory Compliance

**GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Regular onsite and remote monitoring visits by Clinical Research Associates (CRAs) to verify source data and protocol adherence. **Audit Trail:** Validated electronic data capture (EDC) system with full audit trails for ePRO diaries and electronic Case Report Forms (eCRFs).

---

## 11. Study Identifiers & Governance

**Primary ID:** C4951063 **Registry Number:** NCT06641466 **Posting ID:** 271503695776790138 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Study to Learn About the Study Medicine Called Rimegepant in Women When Used for Intermittent Prevention of Menstrual Migraine **Official Regulatory Title:** AN INTERVENTIONAL EFFICACY AND SAFETY, PHASE 3, DOUBLE-BLIND, PARALLEL GROUP STUDY TO INVESTIGATE INTERMITTENT PREVENTION OF MENSTRUAL MIGRAINE WITH RIMEGEPANT COMPARED WITH PLACEBO IN WOMEN PARTICIPANTS 18 TO 45 YEARS OF AGE **Data Source Level:** 5

---

## 12. Clinical Framework (Disease Specific)

**Primary Condition / Disease Area:** Menstrual Migraine (Preferred Name: Menstrual migraine; Preferred UMLS Name: Menstrual Migraine) **Semantic Type:** Pathologic Function **SNOMED ID:** 23186000 (Hierarchy encompasses disorders associated with menstruation and/or menopause, disorders of brain/head/nervous system, vascular headache, etc.) **Diagnosis Threshold:**  $\geq 1$  migraine attack during the perimenstrual period in at least 2 out of 3 menstrual cycles **Medical Methodology:** Intermittent prevention during the peri-menstrual period (PMP) **Optimization Target:** Reduction in the number of migraine days per 5-day perimenstrual period

---

### 13. Study Procedures & Methodology

**Management Pathway:** Intermittent preventive dosing starting 3 days before the expected menstrual period **Education Protocol (HCP):** Investigator training on protocol criteria, ePRO diary administration, and safety reporting. **Education Protocol (Patient):** Instructions on daily ePRO diary completion and adherence to the 7-day monthly dosing schedule. **Quality Audit Mechanism:** ePRO compliance tracking and regular site data monitoring.

---

### 14. Observation & Follow-Up Schedule

**Total Duration:** 5 months (5 menstrual cycles) double-blind phase + optional open-label follow-up **Onsite Visit Cadence:** Screening, Baseline, and periodic visits spanning the 5 treatment cycles. **Remote Monitoring Frequency:** Continuous daily tracking via ePRO diary. **Checklist Review Interval:** Daily (ePRO diary input by the participant).

---

### 15. Sign-off & Approval

|                          |        |           |               |     |      |     |     |
|--------------------------|--------|-----------|---------------|-----|------|-----|-----|
| Role                     | Name   | Signature | Date          | :-- | :--  | :-- | :-- |
| Principal Investigator   | [Name] | _____     | [DD-MMM-YYYY] |     |      |     |     |
| Clinical Operations Lead | [Name] | _____     | [DD-MMM-YYYY] |     | Lead |     |     |
| Statistician             | [Name] | _____     | [DD-MMM-YYYY] |     |      |     |     |