

1. Administrative & Study Identification

Full Study Title: A Study to Learn More About Prevenar 20 Once it is Out in the Korean Market **Official Regulatory Title:** A Study to Learn More About Prevenar 20 Once it is Out in the Korean Market **Short Title/Acronym:** Prevenar 20 Korean PMS (Post-Marketing Surveillance) **Protocol Number:** [Pfizer Internal Study ID] **Primary ID / Registry Number (NCT):** NCT06760208 **Sponsor/Lead Organization:** Pfizer **Investigational Product:** Prevenar 20 (Pneumococcal 20-valent conjugate vaccine) **Primary Condition:** Pneumococcal Immunization **Phase of Development:** Phase 4 / Post-Marketing Observational **Medical Monitor:** Pfizer Clinical Trial Contact Center

2. Study Synopsis & Rationale

2.1 Study Objective **Primary Objective:** To understand the safety and tolerability of the Prevenar 20 vaccine once it is available in the Korean market. **Secondary Objectives:** To collect real-world data regarding the experiences of individuals receiving the study vaccine and to evaluate any adverse events following administration. **Optimization Target:** Safety and Tolerability Tracking

2.2 Background & Rationale Prevenar 20 is a 20-valent pneumococcal conjugate vaccine designed to protect against pneumococcal disease. This study is being conducted to address the regulatory and clinical need for post-marketing surveillance. The study will monitor real-world safety by evaluating the experiences of patients who have been prescribed Prevenar 20 by their physician according to the approved product label in the Republic of Korea.

3. Study Design & Methodology

3.1 Design Framework **Study Type:** Observational (Post-Marketing Surveillance) **Management Pathway:** Post-Marketing Surveillance (PMS) / Real-world tracking **Control Type:** Single-arm **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration **Target N\$:** [TBD - based on post-marketing prescription volume in the Republic of Korea] **Number of Sites:** Multiple sites in the Republic of Korea (including Seoul) **Estimated Study Start:** [2024/2025] **Estimated Primary Completion:** [28 Days post-last patient enrollment]

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Infant, children, and adolescents who are aged 6 weeks to under 18 years, or adults who are 18 years or older (**Diagnosis Threshold:** Preventative Health / Age $\geq$ 6 weeks). 2. Subjects who have been prescribed Prevenar 20 by their physician as per the approved product label. 3. Evidence of a personally signed and dated informed consent document indicating that the subject or their parent(s)/legal guardian have been informed of all pertinent aspects of the study.

4.2 Exclusion Criteria 1. Any known contraindication to the Prevenar 20 vaccine as per the locally approved product label. 2. Unwillingness or inability to participate in the 28-day safety follow-up.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: 1 dose (0.5 mL) of Prevenar 20. **Medical Methodology:** Standard Vaccination Protocol per Local Label. **Route of Administration:** Intramuscular injection, preferably into the shoulder muscle. **Duration of Treatment:** Single injection, followed by a 28-day safety observation period (**Total Duration:** 28 Days).

5.2 Concomitant & Prohibited Therapy Permitted: Routine concomitant medications that are not contraindicated by the Prevenar 20 product label. **Prohibited:** Any therapies strictly conflicting with the vaccine's administration guidelines.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
1	Screening / Baseline	Day 0 Informed Consent, Medical History verification, Administration of Prevenar 20 vaccine. Education Protocol (Patient): Vaccine information briefing prior to administration and follow-up instruction.
2	End of Study	Day 28 Safety information collection. <i>Note: There is no strict fixed visit schedule. If subjects are unable to visit the clinic, safety information will be collected by telephone or e-mail.</i> Onsite Visit Cadence: Day 0 (Administration); Day 28 (Optional onsite). Remote Monitoring Frequency: Day 28 (via phone or e-mail if unable to visit clinic). Checklist Review Interval: End of Study / 28 Days.

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: Incidence of Adverse Events (AEs) and Serious Adverse Events (SAEs) within 28 days following the administration of the Prevenar 20 vaccine. **Measurement**

Method: Patient or caregiver reporting via telephone, e-mail, or follow-up clinic visit at the end of the 28-day observation period.

7.2 Secondary & Exploratory Endpoints General tolerability and real-world experience of the participants receiving the study medicine in the Korean market.

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Method of collection relies on clinical visits, telephone calls, or email communication 28 days after vaccine administration. **Serious Adverse Events (SAEs):** Routine post-marketing pharmacovigilance reporting timelines (typically 24 hours for severe events). **Data Monitoring Committee (DMC):** Pfizer internal pharmacovigilance and safety monitoring boards.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all participants who received the Prevenar 20 vaccine and provided follow-up data). **Sample Size Justification:** Sample size will be driven by the natural real-world prescription volume required by local health authorities for new vaccine surveillance. **Handling of Missing Data:** Missing safety follow-ups will be documented; primary analysis will utilize descriptive statistics for observed events without imputation for missing data.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Pharmacoepidemiology Practices (GPP), Good Clinical Practice (GCP), and the regulatory requirements of the Republic of Korea. **Education Protocol (HCP):** Training on reporting AEs according to the Prevenar 20 product label. **Monitoring Plan:** Centralized monitoring of safety records and follow-up contacts (phone/email). **Quality Audit Mechanism:** Patient follow-up tracking via email or phone at 28 days. **Audit Trail:** Details on eCRF locking and adverse event tracking via Pfizer's centralized data capture systems.

11. Study Identifiers & Governance

Primary ID: {{NCT06760208}} **Registry Number:** {{NCT06760208}}
Sponsor/Lead Organization: {{Pfizer}} **Study Title:** {{A Study to Learn More About Prevenar 20 Once it is Out in the Korean

Market}} **Official Regulatory Title:** {{A Study to Learn More About
Prevenar 20 Once it is Out in the Korean Market}}

12. Clinical Framework (Pneumococcal Immunization Specific)

Primary Condition: {{Pneumococcal Immunization}} **Diagnosis Threshold:** {{Preventative Health / Age \$\\ge\$ 6 weeks}} **Medical Methodology:** {{Standard Vaccination Protocol per Local Label}} **Optimization Target:** {{Safety and Tolerability Tracking}}

13. Study Procedures & Methodology

Management Pathway: {{Post-Marketing Surveillance (PMS) / Real-world tracking}} **Education Protocol (HCP):** {{Training on reporting AEs according to the Prevenar 20 product label}} **Education Protocol (Patient):** {{Vaccine information briefing prior to administration and follow-up instruction}} **Quality Audit Mechanism:** {{Patient follow-up tracking via email or phone at 28 days}}

14. Observation & Follow-Up Schedule

Total Duration: {{28 Days}} **Onsite Visit Cadence:** {{Day 0 (Administration); Day 28 (Optional onsite)}} **Remote Monitoring Frequency:** {{Day 28 (via phone or e-mail if unable to visit clinic)}} **Checklist Review Interval:** {{End of Study / 28 Days}}

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	[Name]	_____	[DD-MMM-YYYY]			Lead	
Statistician	[Name]	_____	[DD-MMM-YYYY]				