

1. Administrative & Study Identification

Full Study Title: A Clinical Trial of PF-08046037 Alone or With Sasanlimab in Patients With Advanced or Metastatic Malignancies
Short Title/Acronym: PF-08046037 +/- Sasanlimab Phase 1 Trial
Protocol Number: PF-08046037-01 **NCT Number:** NCT06974734 **Sponsor Name:** Pfizer **Investigational Product:** PF-08046037 (PD-L1-targeted Immunostimulatory Drug Conjugate [ISAC] with a TLR7 agonist payload) and Sasanlimab (anti-PD-1 monoclonal antibody) **Phase of Development:** Phase 1 **Medical Monitor:** Patrizia Cavazzoni, MD (Chief Medical Officer, Pfizer)

2. Study Synopsis & Rationale

2.1 Study Objective Primary Objective: To evaluate the safety, tolerability, and optimal dose of PF-08046037 administered as monotherapy or in combination with sasanlimab in participants with advanced or metastatic malignancies. **Secondary Objectives:** To evaluate pharmacokinetics (PK), duration of response (DOR), progression-free survival (PFS), overall survival (OS), and immunogenicity (Anti-Drug Antibodies [ADAs]).

2.2 Background & Rationale To address the need for more effective treatments in advanced solid tumors, PF-08046037 was developed to target PD-L1-expressing immunosuppressive macrophages and dendritic cells. It aims to repolarize the suppressive tumor microenvironment, enhance antigen presentation, and establish anti-tumor immunity. Combining it with sasanlimab—a PD-1 inhibitor—offers a novel dual-approach to boost the immune system against tough-to-treat cancers like NSCLC, HNSCC, melanoma, and PDAC.

3. Study Design & Methodology

3.1 Design Framework Study Type: Interventional **Control Type:** Single-arm (Dose-escalation and dose-expansion cohorts) **Blinding:** Open-label **Randomization:** N/A

3.2 Planned Enrollment & Duration Target N: Standard phase 1 dose-escalation and expansion cohorts (Specific N not publicly released) **Number of Sites:** Multicenter **Estimated Study Start:** 06-May-2025 **Estimated Primary Completion:** 31-Mar-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Diagnosis of advanced or metastatic non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), melanoma, or pancreatic ductal adenocarcinoma (PDAC). 2. Must have progressive disease following at least 1 prior approved systemic therapy. 3. Measurable disease per RECIST v1.1. 4. Ability to provide sufficient tumor tissue samples (archival tissue within 6-12 months or de novo baseline/on-treatment tumor biopsies depending on the cohort).

4.2 Exclusion Criteria 1. History of autoimmune disease or active, severe infections. 2. Unacceptable toxicity or hypersensitivity to prior immunotherapy. 3. Untreated or actively progressing central nervous system (CNS) metastases or inadequate organ function.

5. Intervention & Treatment Plan

5.1 Investigational Regimen Dosage/Frequency: Administered once every 3 weeks (Q3W). **Route of Administration:** PF-08046037 via Intravenous (IV) infusion alone, or in combination with Sasanlimab via Subcutaneous (SQ) injection. **Duration of Treatment:** Through disease progression, unacceptable toxicity, or up to approximately 2 years.

5.2 Concomitant & Prohibited Therapy Permitted: Standard palliative care and symptom management. **Prohibited:** Concurrent systemic anti-cancer therapies, strong immunosuppressive medications, or live vaccines.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, RECIST Evaluation, Tumor Tissue Submission
Baseline	Day 0	First Dose (IV/SQ), Baseline PK/PD and Safety Labs
Interim Visits	Q3W (every 3 weeks)	Dosing, Safety Labs, AE Monitoring, Tumor Imaging (per protocol schedule)
End of Study	Up to ~2 Years	Final Vital Signs, Safety Follow-up (30-37 days post-last dose), Survival Follow-up

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint Definition: - Number of participants with adverse events (AEs) and laboratory abnormalities. - Percent change of cells within tumors based on multiplex immunofluorescence. - Number of participants with antidrug antibodies (ADAs). **Measurement Method:** Clinical and laboratory

safety assessments; central laboratory multiplex immunofluorescence.

7.2 Secondary & Exploratory Endpoints - PK parameters (Tmax, AUC, t1/2). - Duration of response (DOR) per RECIST v1.1. - Progression-free survival (PFS) per RECIST v1.1. - Overall survival (OS).

8. Safety & Adverse Event Reporting

Adverse Event (AE) Monitoring: Continuous collection of AEs and laboratory abnormalities through 30-37 days after the last study treatment. **Serious Adverse Events (SAEs):** Evaluated per standard Phase 1 oncology safety reporting timelines. **Data Monitoring Committee (DMC):** Internal safety and dose-escalation review committees to assess Dose-Limiting Toxicities (DLTs) and optimize dosing.

9. Statistical Analysis Plan (SAP) Summary

Analysis Populations: Safety Set (all patients receiving at least one dose) and Efficacy Evaluable Set. **Sample Size Justification:** Scaled for Phase 1 dose-escalation (Part 1a/1b) and expansion phases (Part 2/3) to robustly evaluate safety profiles and preliminary efficacy signals. **Handling of Missing Data:** Efficacy analyses will utilize standard censoring rules for time-to-event endpoints (PFS, DOR, OS) without imputation for safety events.

10. Quality Assurance & Regulatory Compliance

GCP Compliance: This study will be conducted in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. **Monitoring Plan:** Central laboratory evaluation for biomarker/tissue status and central radiographic review for RECIST v1.1 compliance. **Audit Trail:** eCRF locking and routine data clarification/audit procedures.

11. Study Identifiers & Governance

Primary ID: NCT06974734 **Registry Number:** NCT06974734 **Sponsor/Lead Organization:** Pfizer **Study Title:** A Clinical Trial of PF-08046037 Alone or With Sasanlimab in Patients With Advanced or Metastatic Malignancies **Official Regulatory Title:** An open-label Phase 1 study to evaluate PF-08046037 as monotherapy and part of combination therapy in participants with advanced malignancies.

12. Clinical Framework (Oncology Specific)

Primary Condition: Advanced/Metastatic NSCLC, HNSCC, Melanoma, and PDAC **Diagnosis Threshold:** Measurable disease per RECIST v1.1 and progressive disease after ≥ 1 prior therapy **Medical Methodology:** Immuno-oncology (Targeted ISAC + PD-1 inhibition) **Optimization Target:** Repolarization of the suppressive tumor microenvironment and restoration of anti-tumor T-cell immunity

13. Study Procedures & Methodology

Management Pathway: Intravenous and subcutaneous immunotherapy infusions **Education Protocol (HCP):** Management of immune-related Adverse Events (irAEs) and study-specific DLT guidelines **Education Protocol (Patient):** Informed consent, safety reporting instructions, and visit schedule adherence **Quality Audit Mechanism:** Central review of tumor biopsies (multiplex immunofluorescence) and independent RECIST evaluation

14. Observation & Follow-Up Schedule

Total Duration: Up to approximately 2 years **Onsite Visit Cadence:** Every 3 weeks (Q3W) **Remote Monitoring Frequency:** Routine survival and safety status tracking **Checklist Review Interval:** Pre-infusion assessments every 3 weeks

15. Sign-off & Approval

Role	Name	Signature	Date	:--	:--	:--	:--
Principal Investigator	[Name]	_____	[DD-MMM-YYYY]				
Clinical Operations Lead	Patrizia Cavazzoni	_____	[DD-MMM-YYYY]				
Lead Statistician	[Name]	_____	[DD-MMM-YYYY]				