

1. Administrative & Study Identification

Brief Study Title: A Study to Determine the Efficacy, Safety, and Durability of Faricimab in Participants With Neovascular Age-Related Macular Degeneration **Full Study Title:** A Phase IIIb/IV, Multicenter, Randomized, Open-Label, Two-Arm Study to Investigate the Efficacy, Safety, and Durability of Faricimab Administered up to Every 24 Weeks in Patients With Neovascular Age-Related Macular Degeneration **Short Title/Acronym:** CONSTANCE **Protocol Number:** MR45638 / 2024-517545-13-00 **NCT Number:** NCT06795048 **Trial Status:** Active, not recruiting (ACTIVE_NOT_RECRUITING) **Sponsor Name:** Hoffmann-La Roche **Investigational Product:** Faricimab (Trade Name: Vabysmo, ATC Code: S01LA09, 6-mg intravitreal injection) **Phase of Development:** Phase IIIb/IV (Phase 4) **Medical Monitor:** Hoffmann-La Roche Clinical Operations Lead

2. Study Synopsis & Rationale

2.1 Study Objective * **Primary Objective:** To investigate the efficacy, safety, and durability of intravitreal 6-mg faricimab administered at up to 24-week intervals in patients with neovascular age-related macular degeneration (nAMD) that are treatment-naïve in the study eye. * **Secondary Objectives:** To assess long-term visual and anatomic outcomes (e.g., Best-Corrected Visual Acuity [BCVA], subretinal fluid reduction) and evaluate the number of injections required over the 100-week treatment duration.

2.2 Background & Rationale Neovascular (or wet) age-related macular degeneration (nAMD) is characterized by the growth of unhealthy blood vessels in the central area at the back of the eye, causing swelling, fluid leakage, and vision loss. Standard treatments require frequent clinic visits for intravitreal injections, creating a high treatment burden. This study investigates whether administering faricimab at extended intervals (up to 24 weeks) can maintain treatment efficacy and safety while significantly reducing the patient and healthcare burden.

3. Study Design & Methodology

3.1 Design Framework * **Study Type:** Interventional * **Control Type:** Active Comparator (Two-arm) * **Blinding:** Open-label * **Randomization:** Randomized

3.2 Planned Enrollment & Duration * **Target N\$:** 274 evaluable patients * **Number of Sites:** Approximately 68 global locations * **Estimated Study Start:** 14-Mar-2025 * **Estimated Primary Completion:** 15-Oct-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria 1. Overtly healthy as determined by medical evaluation that includes medical history and physical examination. 2. Agreement to adhere to the contraception requirements described in the protocol. 3. Active treatment-naïve macular neovascularization (MNV) secondary to age-related macular degeneration (AMD), confirmed by the investigator based on the presence of intraretinal fluid (IRF) or subretinal fluid (SRF) affecting the central subfield on optical coherence tomography (OCT). 4. BCVA of 83 to 24 letters, inclusive (20/25 to 20/320 approximate Snellen equivalent, using the early treatment diabetic retinopathy study [ETDRS] protocol and addressed at the initial testing distance of 4 meters on Day 1). 5. Sufficiently clear ocular media and adequate pupillary dilation to allow acquisition of good quality retinal images to confirm diagnosis.

4.2 Exclusion Criteria 1. MNV due to causes other than nAMD, such as ocular histoplasmosis, trauma, pathological myopia, angioid streaks, choroidal rupture, or uveitis. 2. Retinal pigment epithelial tear involving the macula on Day 1. 3. Current vitreous hemorrhage on Day 1. 4. Prior periocular pharmacological or intravitreal (IVT) treatment (including faricimab, anti-vascular endothelial growth factor [VEGF], or complement inhibitor medication) for other retinal diseases. 5. Participants who have a nonfunctioning fellow (non-study) eye, defined as either BCVA of hand motion or worse, or no physical presence of non-study eye (i.e., monocular), at both the screening and study Day 1 visits. 6. History of idiopathic or autoimmune associated uveitis in either eye, or active ocular inflammation or suspected or active ocular or periocular infection in either eye on study Day 1.

5. Intervention & Treatment Plan

5.1 Investigational Regimen * **Dosage/Frequency:** 6-mg administered at up to 24-week extended intervals. * **Route of Administration:** Intravitreal (IVT) injection. * **Duration of Treatment:** 100 weeks of active treatment.

5.2 Concomitant & Prohibited Therapy * **Permitted:** Standard concomitant systemic medications for other co-morbidities. * **Prohibited:** Concurrent administration of other experimental or approved anti-VEGF treatments or complement inhibitors in the study eye.

6. Study Timeline & Visit Schedule

Visit ID	Window (Days)	Key Activities/Procedures
Screening	Day -28 to 0	Informed Consent, Medical History, Ophthalmic Exam, OCT, BCVA assessment
Baseline	Day 1	Randomization, First IVT Injection of Faricimab, Safety check
Interim	Months 1 to 24	Visual acuity checks, OCT, IVT Injections (every 1 to 6 months depending on randomized arm interval), Safety Monitoring
End of Study	Week 104	Final safety assessment and follow-up visit 1 month after the final dose of study treatment

7. Outcome Measures (Endpoints)

7.1 Primary Endpoint * **Definition:** Change from Baseline in Best-Corrected Visual Acuity (BCVA) Score Averaged Over Weeks 44, 48, and 52. * **Measurement Method:** Standardized BCVA evaluation using the ETDRS protocol and centralized OCT imaging reading.

7.2 Secondary & Exploratory Endpoints * Change from Baseline in BCVA Score Averaged Over Weeks 92, 96, and 100. * Change from Baseline in BCVA Score Over Time. * Percentage of Participants Avoiding a Loss of ≥ 15 Letters in BCVA from Baseline Over Time. * Change from baseline in Central Subfield Thickness (CST) as measured by OCT. * Total number of intravitreal injections received in the study eye over the 100-week period.

8. Safety & Adverse Event Reporting

- **Adverse Event (AE) Monitoring:** Conducted at every scheduled onsite visit (every 1 to 6 months), with specific attention paid to intraocular inflammation and intraocular pressure spikes.
- **Serious Adverse Events (SAEs):** Required to be reported within 24 hours of investigator awareness.
- **Data Monitoring Committee (DMC):** Independent Data Monitoring Committee appointed to periodically review unblinded safety and efficacy data.

9. Statistical Analysis Plan (SAP) Summary

- **Analysis Populations:** Intent-to-Treat (ITT) for primary efficacy evaluation; Safety Set for all participants who received at least one dose of the investigational product.
- **Sample Size Justification:** A sample of $N=274$ provides sufficient statistical power (e.g., $\text{Power} = 80\%$, $\alpha = 0.05$) to evaluate the durability and non-inferiority/efficacy profiles of extended 24-week dosing versus standard historical intervals.

- **Handling of Missing Data:** Mixed-Effect Model Repeated Measure (MMRM) and/or Multiple Imputation methods applied for visual acuity and anatomical continuous variables.
-

10. Quality Assurance & Regulatory Compliance

- **GCP Compliance:** This study will be conducted in accordance with Good Clinical Practice (GCP) guidelines and the Declaration of Helsinki.
 - **Monitoring Plan:** Routine onsite and remote monitoring to verify adherence to protocol, especially injection techniques and imaging acquisition.
 - **Audit Trail:** Continuous tracking and resolution of data clarification forms with final eCRF locking prior to unblinding and database freeze.
-

11. Study Identifiers & Governance

- **Primary ID:** MR45638 / 2024-517545-13-00
 - **Registry Number:** NCT06795048
 - **Posting ID:** 396550255396827043
 - **Sponsor/Lead Organization:** Hoffmann-La Roche
 - **Brief Title:** A Study to Determine the Efficacy, Safety, and Durability of Faricimab in Participants With Neovascular Age-Related Macular Degeneration
 - **Official Regulatory Title:** A Phase IIIb/IV, Multicenter, Randomized, Open-Label, Two-Arm Study to Investigate the Efficacy, Safety, and Durability of Faricimab Administered up to Every 24 Weeks in Patients With Neovascular Age-Related Macular Degeneration
 - **EMA Product Information URL:** https://www.ema.europa.eu/en/documents/product-information/vabysmo-epar-product-information_en.pdf
-

12. Clinical Framework (nAMD Specific)

- **Primary Condition:** {{Neovascular Age-Related Macular Degeneration (nAMD)}}
 - **Diagnosis Threshold:** {{Presence of intraretinal/subretinal fluid (IRF/SRF) on OCT and ETDRS BCVA between 83 and 24 letters}}
 - **Medical Methodology:** {{Anti-VEGF and Angiopoietin-2 (Ang-2) Inhibition Therapy}}
 - **Optimization Target:** {{Maintenance of vision and anatomy on an extended treatment interval up to 24 weeks}}
-

13. Study Procedures & Methodology

- **Management Pathway:** {{Standard clinical pathway for nAMD with extended durability dosing protocol}}
 - **Education Protocol (HCP):** {{Central reading center certification for OCT imaging and standardized BCVA testing training}}
 - **Education Protocol (Patient):** {{Counseling on identifying sudden visual changes or symptoms of ocular inflammation}}
 - **Quality Audit Mechanism:** {{Masked centralized reading center evaluation for anatomic criteria and interval extension decisions}}
-

14. Observation & Follow-Up Schedule

- **Total Duration:** {{104 weeks (100-week treatment duration + 4-week follow-up)}}
 - **Onsite Visit Cadence:** {{Every 1 to 6 months depending on assigned injection interval}}
 - **Remote Monitoring Frequency:** {{Ad hoc telehealth evaluations for interval vision changes}}
 - **Checklist Review Interval:** {{At each scheduled injection and monitoring visit}}
-

15. Sign-off & Approval

Role	Name	Signature	Date	:---	:---	:---	:---
Principal Investigator	[Insert Name]	_____	07-Apr-2026				
Clinical Operations Lead	[Insert Name]	_____	07-Apr-2026				
Lead Statistician	[Insert Name]	_____	07-Apr-2026				